

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : A61K 39/12, C07H 21/04 C07K 7/06, 7/08, 7/10 C07K 13/00, 15/28	A1	(11) International Publication Number: WO 90/11089 (43) International Publication Date: 4 October 1990 (04.10.90)									
(21) International Application Number: PCT/US90/01348 (22) International Filing Date: 15 March 1990 (15.03.90) (30) Priority data: <table border="0" style="width: 100%;"><tr><td style="width: 30%;">325,338</td><td style="width: 40%;">17 March 1989 (17.03.89)</td><td style="width: 30%;">US</td></tr><tr><td>341,334</td><td>20 April 1989 (20.04.89)</td><td>US</td></tr><tr><td>355,002</td><td>18 May 1989 (18.05.89)</td><td>US</td></tr></table> (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US). (72) Inventors: HOUGHTON, Michael ; 53 Rosemead Court, Danville, CA 94526 (US). CHOO, Qui-Lim ; 5700 Fern Street, El Cerrito, CA 94530 (US). KUO, George ; 1370 Sixth Avenue, San Francisco, CA 94122 (US).		325,338	17 March 1989 (17.03.89)	US	341,334	20 April 1989 (20.04.89)	US	355,002	18 May 1989 (18.05.89)	US	(74) Agents: MONROY, Gladys, H. et al.; Irell & Manella, 545 Middlefield Road, Suite 200, Menlo Park, CA 94025 (US). (81) Designated States: AU, FI, HU, JP, KR, NO, SU. Published <i>With international search report.</i>
325,338	17 March 1989 (17.03.89)	US									
341,334	20 April 1989 (20.04.89)	US									
355,002	18 May 1989 (18.05.89)	US									
(54) Title: NANBV DIAGNOSTICS AND VACCINES (57) Abstract <p>A new virus, Hepatitis C virus (HCV), which has proven to be the major etiologic agent of blood-borne NANBH, was discovered by Applicant. The initial work on this virus, which includes a partial genomic sequence of the prototype HCV isolate, is described in EPO Pub. No. 318,216, and PCT Pub. No. WO 89/04669. The present invention, which in part is based on new HCV sequences and polypeptides which are not disclosed in the above-cited publications, includes the application of these new sequences and polypeptides in immunoassays, probe diagnostics, anti-HCV antibody production, PCR technology, and recombinant DNA technology. Included within the invention also are novel, immunogenic polypeptides encoded within clones containing HCV cDNA, novel methods for purifying an immunogenic HCV polypeptide, and antisense polynucleotides derived from HCV cDNA.</p>											

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

5

NANBV DIAGNOSTICS AND VACCINES10 Technical Field

The invention relates to materials and methodologies for managing the spread of non-A, non-B hepatitis virus (NANBV) infection. More specifically, it relates to polynucleotides derived from the genome of an etiological agent of NANBH, hepatitis C virus (HCV), to polypeptides encoded therein, and to antibodies directed to the polypeptides. These reagents are useful as screening agents for HCV and its infection, and as protective agents against the disease.

20

References Cited in the Application

- Barr et al. (1986), *Biotechniques* 4:428.
Botstein (1979), *Gene* 8:17.
25 Brinton, M.A. (1986) in *THE VIRUSES: THE TOGAVIRIDAE AND FLAVIVIRIDAE* (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press), p.327-374.
Broach (1981) in: *Molecular Biology of the Yeast Saccharomyces*, Vol. 1, p.445, Cold Spring Harbor Press.
30 Broach et al. (1983), *Meth. Enz.* 101:307.
Chang et al. (1977), *Nature* 198:1056.
Chirgwin et al. (1979), *Biochemistry* 18:5294.
Chomczynski and Sacchi (1987), *Analytical Biochemistry*
35 162:156.
Clewell et al. (1969), *Proc. Natl. Acad. Sci. USA* 62:1159.

- Clewell (1972), J. Bacteriol. 110:667.
- Cohen (1972), Proc. Natl. Acad. Sci. USA 69:2110.
- Cousens et al. (1987), gene 61:265.
- 5 De Boer et al. (1983), Proc. Natl. Acad. Sci. USA 292:128.
- Dreesman et al. (1985), J. Infect. Disease 151:761.
- Feinstone, S.M. and Hoofnagle, J.H. (1984), New Engl. J. Med. 311:185.
- 10 Fields & Knipe (1986), FUNDAMENTAL VIROLOGY (Raven Press, N.Y.).
- Fiers et al. (1978), Nature 273:113.
- Gerety, R.J. et al., in VIRAL HEPATITIS AND LIVER DISEASE (Vyas, B.N., Dienstag, J.L., and Hoofnagle, J.H., eds, Grune and Stratton, Inc., 1984) pp 23-47.
- 15 Goeddel et al. (1980), Nucleic Acids Res. 8:4057.
- Graham and Van der Eb (1978), Virology 52:546.
- Grunstein and Hogness (1975), Proc. Natl. Acad. Sci. USA 73:3961.
- Grych et al. (1985), Nature 316:74.
- 20 Gubler and Hoffman (1983), Gene 25:263.
- Hahn (1988) Virology 162:167.
- Hammerling et al. (1981), MONOCLONAL ANTIBODIES AND T-CELL HYBRIDOMAS.
- Han (1987), Biochemistry 26:1617.
- 25 Helfman (1983), Proc. Natl. Acad. Sci. USA 80:31.
- Hess et al. (1968), J. Adv. Enzyme Reg 7:149.
- Hinnen et al. (1978), Proc. Natl. Acad. Sci. 75:1929.
- Hitzeman et al. (1980), J. Biol. Chem. 255:2073.
- Holland et al. (1978), Biochemistry 17:4900.
- 30 Holland (1981), J. Biol. Chem. 256: 1385.
- Houghton et al. (1981), Nucleic Acids Res. 9:247
- Hunyh, T.V. et al. (1985) in DNA CLONING TECHNIQUES; A PRACTICAL APPROACH (D. Glover, Ed., IRL Press, Oxford, U.K.) pp. 49-78.
- 35 Immun. Rev. (1982) 62:185.
- Iwarson (1987), British Medical J. 295:946.

- Kennett et al. (1980) MONOCLONAL ANTIBODIES.
Kyte and Doolittle (1982), J. Mol. Biol. 157:105-132.
Laemmli (1970), Nature 227, 680.
5 Lee et al. (1988), Science 239:1288.
Maniatis, T., et al. (1982) MOLECULAR CLONING; A
LABORATORY MANUAL (Cold Spring Harbor Press, Cold Spring
Harbor, N.Y.).
10 Mayer and Walker, eds. (1987), IMMUNOCHEMICAL METHODS IN
CELL AND MOLECULAR BIOLOGY (Academic Press, London).
Maxam et al. (1980), Methods in Enzymology 65:499.
MacNamara et al. (1984), Science 226:1325.
Messing et al. (1981), Nucleic Acids Res. 9:309.
Messing (1983), Methods in Enzymology 101:20-37.
15 METHODS IN ENZYMOLOGY (Academic Press).
Michelle et al., Int. Symposium on Viral Hepatitis.
Monath (1986) in THE VIRUSES: THE TOGAVIRIDAE AND
FLAVIVIRIDAE (Series eds. Fraenkel-Conrat and Wagner, vol.
eds. Schlesinger and Schlesinger, Plenum Press), p.375-
20 440.
Nagahuma et al. (1984), Anal. Biochem. 141:74.
Neurath et al. (1984), Science 224:392.
Nisonoff et al. (1981), Clin. Immunol. Immunopathol.
21:397-406.
25 Overby, L.R. (1985), Curr. Hepatol. 5:49.
Overby, L.R. (1986), Curr. Hepatol. 6:65.
Overby, L.R. (1987), Curr. Hepatol. 7:35.
Peleg (1969), Nature 221:193.
Pfefferkorn and Shapiro (1974), in COMPREHENSIVE VIROLOGY,
30 Vol. 2 (Fraenkel-Conrat & Wagner, eds., Plenum, N.Y.) pp.
171-230.
Prince, A.M. (1983), Annu. Rev. Microbiol. 37:217.
Rice et al. (1985), Science 229:726.

- Rice et al. (1986) in THE VIRUSES: THE TOGAVIRIDAE AND FLAVIVIRIDAE (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press), p.279-328.
- 5 Roehrig (1986) in THE VIRUSES: THE TOGAVIRIDAE AND FLAVIVIRIDAE (Series eds. Fraenkel-Conrat and Wagner, vol. eds. Schlesinger and Schlesinger, Plenum Press)
- Sadler et al. (1980), Gene 8, 279.
- 10 Saiki et al. (1986), Nature 324: 163.
- Saiki et al. (1988), Science 239:487.
- Sanger et al. (1977), Proc. Natl. Acad. Sci. USA 74:5463.
- Schlesinger et al. (1986), J. Virol. 60:1153.
- Schreier, M., et al. (1980) HYBRIDOMA TECHNIQUES
- 15 Scopes (1984), PROTEIN PURIFICATION, PRINCIPLES AND PRACTICE, SECOND EDITION (Springer-Verlag, N.Y.).
- Shimatake et al. (1981), Nature 292:128.
- Sippel (1973), Eur. J. Biochem. 37:31.
- Steimer et al. (1986), J. Virol. 58:9.
- 20 Stollar (1980), in THE TOGAVIRUSES (R.W. Schlesinger, ed., Academic Press, N.Y.), pp. 584-622.
- Sumiyoshi et al. (1987), Virology 161:497.
- Taylor et al. (1976), Biochem. Biophys. Acta 442:324.
- Towbin et al. (1979), Proc. Natl. Acad. Sci. USA 76, 4350.
- 25 Tsu and Herzenberg (1980), in SELECTED METHODS IN CELLULAR IMMUNOLOGY (W.H. Freeman and Co.) pp. 373-391.
- Vytdehaag et al. (1985), J. Immunol. 134:1225.
- Valenzuela, P., et al. (1982), Nature 298:344.
- Valenzuela, P., et al. (1984), in HEPATITIS B (Millman, I., et al., ed, Plenum Press) pp. 225-236.
- 30 Warner (1984), DNA 3:401.
- Wu and Grossman (1987), Methods in Enzymology Vol. 154, RECOMBINANT DNA, Part E.
- Wu (1987), Methods in Enzymology vol 155, RECOMBINANT DNA, part F.
- 35 Zoller (1982), Nucleic Acids Res. 10:6487.

Cited Patents

- 5 EPO Pub. No. 318,216
PCT Pub. No. WO 89/04669
U.S. Patent No. 4,341,761
U.S. Patent No. 4,399,121
U.S. Patent No. 4,427,783
U.S. Patent No. 4,444,887
10 U.S. Patent No. 4,466,917
U.S. Patent No. 4,472,500
U.S. Patent No. 4,491,632
U.S. Patent No. 4,493,890

Background Art

- 15 Non-A, Non-B hepatitis (NANBH) is a transmissible disease or family of diseases that are believed to be viral-induced, and that are distinguishable from other forms of viral-associated liver diseases, including that caused by the known hepatitis viruses,
20 i.e., hepatitis A virus (HAV), hepatitis B virus (HBV), and delta hepatitis virus (HDV), as well as the hepatitis induced by cytomegalovirus (CMV) or Epstein-Barr virus (EBV). NANBH was first identified in transfused individuals. Transmission from man to chimpanzee and se-
25 rial passage in chimpanzees provided evidence that NANBH is due to a transmissible infectious agent or agents.

- Epidemiologic evidence is suggestive that there may be three types of NANBH: the water-borne epidemic type; the blood or needle associated type; and the
30 sporadically occurring (community acquired) type. However, the number of agents which may be the causative of NANBH are unknown.

- Clinical diagnosis and identification of NANBH has been accomplished primarily by exclusion of other
35 viral markers. Among the methods used to detect putative NANBV antigens and antibodies are agar-gel diffusion,

counterimmunoelectrophoresis, immunofluorescence
microscopy, immune electron microscopy, radioimmunoassay,
and enzyme-linked immunosorbent assay. However, none of
5 these assays has proved to be sufficiently sensitive,
specific, and reproducible to be used as a diagnostic test
for NANBH.

Previously there was neither clarity nor agree-
ment as to the identity or specificity of the antigen
10 antibody systems associated with agents of NANBH. This
was due, at least in part, to the prior or co-infection of
HBV with NANBV in individuals, and to the known complexity
of the soluble and particulate antigens associated with
HBV, as well as to the integration of HBV DNA into the
15 genome of liver cells. In addition, there is the pos-
sibility that NANBH is caused by more than one infectious
agent, as well as the possibility that NANBH has been mis-
diagnosed. Moreover, it is unclear what the serological
assays detect in the serum of patients with NANBH. It has
20 been postulated that the agar-gel diffusion and counter-
immunoelectrophoresis assays detect autoimmune responses
or nonspecific protein interactions that sometimes occur
between serum specimens, and that they do not represent
specific NANBV antigen-antibody reactions. The
25 immunofluorescence, and enzyme-linked immunosorbent, and
radioimmunoassays appear to detect low levels of a
rheumatoid-factor-like material that is frequently present
in the serum of patients with NANBH as well as in patients
with other hepatic and nonhepatic diseases. Some of the
30 reactivity detected may represent antibody to host-
determined cytoplasmic antigens.

There have been a number of candidate NANBV.
See, for example the reviews by Prince (1983), Feinstone
and Hoofnagle (1984), and Overby (1985, 1986, 1987) and
35 the article by Iwarson (1987). However, there is no proof

that any of these candidates represent the etiological agent of NANBH.

5 The demand for sensitive, specific methods for screening and identifying carriers of NANBV and NANBV contaminated blood or blood products is significant. Post-transfusion hepatitis (PTH) occurs in approximately 10% of transfused patients, and NANBH accounts for up to 90% of these cases. The major problem in this disease is
10 the frequent progression to chronic liver damage (25-55%).

 Patient care as well as the prevention of transmission of NANBH by blood and blood products or by close personal contact require reliable screening,
15 diagnostic and prognostic tools to detect nucleic acids, antigens and antibodies related to NANBV. In addition, there is also a need for effective vaccines and immunotherapeutic therapeutic agents for the prevention
and/or treatment of the disease.

20 Applicant discovered a new virus, the Hepatitis C virus (HCV), which has proven to be the major etiologic agent of blood-borne NANBH (BB-NANBH). Applicant's initial work, including a partial genomic sequence of the prototype HCV isolate, CDC/HCV1 (also called HCV1), is described in EPO Pub. No. 318,216 (published 31 May 1989)
25 and PCT Pub. No. WO 89/04669 (published 1 June 1989). The disclosures of these patent applications, as well as any corresponding national patent applications, are incorporated herein by reference. These applications teach, inter alia, recombinant DNA methods of cloning and
30 expressing HCV sequences, HCV polypeptides, HCV immunodiagnostic techniques, HCV probe diagnostic techniques, anti-HCV antibodies, and methods of isolating new hCV sequences, including sequences of new HCV
isolates.

35

Disclosure of the Invention

5 The present invention is based, in part, on new HCV sequences and polypeptides that are not disclosed in EPO Pub. No. 318,216, or in PCT Pub. No. WO 89/04669. Included within the invention is the application of these new sequences and polypeptides in, inter alia, immunodiagnos-
10 tics, probe diagnostics, anti-HCV antibody production, PCR technology and recombinant DNA technology. Included within the invention, also, are new immunoassays based upon the immunogenicity of HCV polypeptides disclosed herein. The new subject matter claimed herein, while developed using techniques described in, for
15 example, EPO Pub. No. 318,216, has a priority date which antecedes that publication, or any counterpart thereof. Thus, the invention provides novel compositions and methods useful for screening samples for HCV antigens and antibodies, and useful for treatment of HCV infections.


20 Accordingly, one aspect of the invention is a recombinant polynucleotide comprising a sequence derived from HCV cDNA, wherein the HCV cDNA is in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone pil4a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or
25 clone 16jh, or wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17.

Another aspect of the invention is a purified polypeptide comprising an epitope encoded within HCV cDNA
30 wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17.

Yet another aspect of the invention is an immunogenic polypeptide produced by a cell transformed with
35 a recombinant expression vector comprising an ORF of DNA derived from HCV cDNA, wherein the HCV cDNA is comprised

of a sequence derived from the HCV cDNA sequence in clone CA279a, or clone CA74a, or clone 13i, or clone CA290a, or clone 33C or clone 40b, or clone 33b, or clone 25c, or
 5 clone 14c, or clone 8f, or clone 33f, or clone 33g, or clone 39c, or clone 15e, and wherein the ORF is operably linked to a control sequence compatible with a desired host.

Another aspect of the invention is a peptide
 10 comprising an HCV epitope, wherein the peptide is of the formula



15 wherein x and y designate amino acid numbers shown in Fig. 17, and wherein the peptide is selected from the group consisting of AA1-AA25, AA1-AA50, AA1-AA84, AA9-AA177, AA1-AA10, AA5-AA20, AA20-AA25, AA35-AA45, AA50-AA100, AA40-AA90, AA45-AA65, AA65-AA75, AA80-90, AA99-AA120,
 20 AA95-AA110, AA105-AA120, AA100-AA150, AA150-AA200, AA155-AA170, AA190-AA210, AA200-AA250, AA220-AA240, AA245-AA265, AA250-AA300, AA290-AA330, AA290-305, AA300-AA350, AA310-AA330, AA350-AA400, AA380-AA395, AA405-AA495, AA400-AA450, AA405-AA415, AA415-AA425,
 25 AA425-AA435, AA437-AA582, AA450-AA500, AA440-AA460, AA460-AA470, AA475-AA495, AA500-AA550, AA511-AA690, AA515-AA550, AA550-AA600, AA550-AA625, AA575-AA605, AA585-AA600, AA600-AA650, AA600-AA625, AA635-AA665, AA650-AA700, AA645-AA680, AA700-AA750, AA700-AA725, AA700-AA750,
 30 AA725-AA775, AA770-AA790, AA750-AA800, AA800-AA815, AA825-AA850, AA850-AA875, AA800-AA850, AA920-AA990, AA850-AA900, AA920-AA945, AA940-AA965, AA970-AA990, AA950-AA1000, AA1000-AA1060, AA1000-AA1025, AA1000-AA1050, AA1025-AA1040, AA1040-AA1055, AA1075-AA1175,
 35 AA1050-AA1200, AA1070-AA1100, AA1100-AA1130, AA1140-AA1165, AA1192-AA1457, AA1195-AA1250,

AA1200-AA1225, AA1225-AA1250, AA1250-AA1300,
AA1260-AA1310, AA1260-AA1280, AA1266-AA1428,
AA1300-AA1350, AA1290-AA1310, AA1310-AA1340, AA1345-
5 AA1405, AA1345-AA1365, AA1350-AA1400, AA1365-AA1380,
AA1380-AA1405, AA1400-AA1450, AA1450-AA1500,
AA1460-AA1475, AA1475-AA1515, AA1475-AA1500,
AA1500-AA1550, AA1500-AA1515, AA1515-AA1550,
AA1550-AA1600, AA1545-AA1560, AA1569-AA1931,
10 AA1570-AA1590, AA1595-AA1610, AA1590-AA1650,
AA1610-AA1645, AA1650-AA1690, AA1685-AA1770,
AA1689-AA1805, AA1690-AA1720, AA1694-AA1735,
AA1720-AA1745, AA1745-AA1770, AA1750-AA1800,
AA1775-AA1810, AA1795-AA1850, AA1850-AA1900,
15 AA1900-AA1950, AA1900-AA1920, AA1916-AA2021,
AA1920-AA1940, AA1949-AA2124, AA1950-AA2000,
AA1950-AA1985, AA1980-AA2000, AA2000-AA2050,
AA2005-AA2025, AA2020-AA2045, AA2045-AA2100,
AA2045-AA2070, AA2054-AA2223, AA2070-AA2100,
20 AA2100-AA2150, AA2150-AA2200, AA2200-AA2250,
AA2200-AA2325, AA2250-AA2330, AA2255-AA2270,
AA2265-AA2280, AA2280-AA2290, AA2287-AA2385,
AA2300-AA2350, AA2290-AA2310, AA2310-AA2330,
AA2330-AA2350, AA2350-AA2400, AA2348-AA2464,
25 AA2345-AA2415, AA2345-AA2375, AA2370-AA2410,
AA2371-AA2502, AA2400-AA2450, AA2400-AA2425,
AA2415-AA2450, AA2445-AA2500, AA2445-AA2475,
AA2470-AA2490, AA2500-AA2550, AA2505-AA2540,
AA2535-AA2560, AA2550-AA2600, AA2560-AA2580,
30 AA2600-AA2650, AA2605-AA2620, AA2620-AA2650,
AA2640-AA2660, AA2650-AA2700, AA2655-AA2670,
AA2670-AA2700, AA2700-AA2750, AA2740-AA2760,
AA2750-AA2800, AA2755-AA2780,
AA2780-AA2830, AA2785-AA2810, AA2796-AA2886,
35 AA2810-AA2825, AA2800-AA2850, AA2850-AA2900,

AA2850-AA2865, AA2885-AA2905, AA2900-AA2950,
AA2910-AA2930, AA2925-AA2950, AA2945-end(C' terminal).

5 Still another aspect of the invention is a
monoclonal antibody directed against an epitope encoded in
HCV cDNA, wherein the HCV cDNA is of a sequence indicated
by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig.
17, or is the sequence present in clone 13i, or clone 26j,
or clone 59a, or clone 84a, or clone CA156e, or clone
10 167b, or clone pil4a, or clone CA216a, or clone CA290a, or
clone ag30a, or clone 205a, or clone 18g, or clone 16jh.

Yet another aspect of the invention is a
preparation of purified polyclonal antibodies directed
against a polypeptide comprised of an epitope encoded
15 within HCV cDNA; wherein the HCV cDNA is of a sequence
indicated by nucleotide numbers -319 to 1348 or 8659 to
8866 in Fig. 17, or is the sequence present in in clone
13i, or clone 26j, or clone 59a, or clone 84a, or clone
CA156e, or clone 167b, or clone pil4a, or clone CA216a, or
20 clone CA290a, or clone ag30a, or clone 205a, or clone 18g,
or clone 16jh.

Still another aspect of the invention is a
polynucleotide probe for HCV, wherein the probe is
comprised of an HCV sequence derived from an HCV cDNA
25 sequence indicated by nucleotide numbers -319 to 1348 or
8659 to 8866 in Fig. 17, or from the complement of the HCV
cDNA sequence.

Yet another aspect of the invention is a kit for
analyzing samples for the presence of polynucleotides from
30 HCV comprising a polynucleotide probe containing a
nucleotide sequence of about 8 or more nucleotides,
wherein the nucleotide sequence is derived from HCV cDNA
which is of a sequence indicated by nucleotide numbers -
319 to 1348 or 8659 to 8866 in Fig. 17, wherein the
35 polynucleotide probe is in a suitable container.

Another aspect of the invention is a kit for analyzing samples for the presence of an HCV antigen comprising an antibody which reacts immunologically with an HCV antigen, wherein the antigen contains an epitope encoded within HCV cDNA which is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or wherein the HCV cDNA is in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone p14a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh.

Yet another aspect of the invention is a kit for analyzing samples for the presence of an HCV antibody comprising an antigenic polypeptide containing an HCV epitope encoded within HCV cDNA which is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone p14a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh.

Another aspect of the invention is a kit for analyzing samples for the presence of an HCV antibody comprising an antigenic polypeptide expressed from HCV cDNA in clone CA279a, or clone CA74a, or clone 13i, or clone CA290a, or clone 33C or clone 40b, or clone 33b, or clone 25c, or clone 14c, or clone 8f, or clone 33f, or clone 33g, or clone 39c, or clone 15e, wherein the antigenic polypeptide is present in a suitable container.

Still another aspect of the invention is a method for detecting HCV nucleic acids in a sample comprising:

(a) reacting nucleic acids of the sample with a polynucleotide probe for HCV, wherein the probe is comprised of an HCV sequence derived from an HCV cDNA sequence is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, and wherein the

reacting is under conditions which allow the formation of a polynucleotide duplex between the probe and the HCV nucleic acid from the sample; and (b) detecting a
5 polynucleotide duplex which contains the probe, formed in step (a).

Yet another aspect of the invention is an immunoassay for detecting an HCV antigen comprising:

(a) incubating a sample suspected of containing
10 an HCV antigen with an antibody directed against an HCV epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is the sequence present in clone 13i, or clone 26j, or clone 59a, or clone 84a, or
15 clone CA156e, or clone 167b, or clone p114a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, and wherein the incubating is under conditions which allow formation of an antigen-antibody complex; and (b) detecting an antibody-antigen
20 complex formed in step (a) which contains the antibody.

Still another aspect of the invention is an immunoassay for detecting antibodies directed against an HCV antigen comprising:

(a) incubating a sample suspected of containing
25 anti-HCV antibodies with an antigen polypeptide containing an epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is the sequence present in clone 13i, or clone 26j, or clone 59a, or clone 84a, or
30 clone CA156e, or clone 167b, or clone p114a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, and wherein the incubating is under conditions which allow formation of an antigen-antibody complex; and detecting an antibody-antigen
35 complex formed in step (a) which contains the antigen polypeptide.

Another aspect of the invention is a vaccine for treatment of HCV infection comprising an immunogenic polypeptide containing an HCV epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17 or is the sequence present in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone p114a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, and wherein the immunogenic polypeptide is present in a pharmacologically effective dose in a pharmaceutically acceptable excipient.

Yet another aspect of the invention is a method for producing antibodies to HCV comprising administering to an individual an isolated immunogenic polypeptide containing an HCV epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is of the sequence present in clone CA279a, or clone CA74a, or clone 13i, or clone CA290a, or clone 33C or clone 40b, or clone 33b, or clone 25c, or clone 14c, or clone 8f, or clone 33f, or clone 33g, or clone 39c, or clone 15e, and wherein the immunogenic polypeptide is present in a pharmacologically effective dose in a pharmaceutically acceptable excipient.

Still another aspect of the invention is an antisense polynucleotide derived from HCV cDNA, wherein the HCV cDNA is that shown in Fig. 17.

Yet another aspect of the invention is a method for preparing purified fusion polypeptide C100-3 comprising:

(a) providing a crude cell lysate containing polypeptide C100-3,

- (b) treating the crude cell lysate with an amount of acetone which causes the polypeptide to precipitate,
- 5 (c) isolating and solubilizing the precipitated material,
- (d) isolating the C100-3 polypeptide by anion exchange chromatography, and
- 10 (e) further isolating the C100-3 polypeptide of step (d) by gel filtration.

Brief Description of the Drawings

15 Fig. 1 shows the sequence of the HCV cDNA in clone 12f, and the amino acids encoded therein.

Fig. 2 shows the HCV cDNA sequence in clone k9-1, and the amino acids encoded therein.

20 Fig. 3 shows the sequence of clone 15e, and the amino acids encoded therein.

Fig. 4 shows the nucleotide sequence of HCV cDNA in clone 13i, the amino acids encoded therein, and the sequences which overlap with clone 12f.

25 Fig. 5 shows the nucleotide sequence of HCV cDNA in clone 26j, the amino acids encoded therein, and the sequences which overlap clone 13i.

Fig. 6 shows the nucleotide sequence of HCV cDNA in clone CA59a, the amino acids encoded therein, and the sequences which overlap with clones 26j and K9-1.

30 Fig. 7 shows the nucleotide sequence of HCV cDNA in clone CA84a, the amino acids encoded therein, and the sequences which overlap with clone CA59a.

35 Fig. 8 shows the nucleotide sequence of HCV cDNA in clone CA156e, the amino acids encoded therein, and the sequences which overlap with CA84a.

Fig. 9 shows the nucleotide sequence of HCV cDNA in clone CA167b, the amino acids encoded therein, and the sequences which overlap CA156e.

Fig. 10 shows the nucleotide sequence of HCV cDNA in clone CA216a, the amino acids encoded therein, and the overlap with clone CA167b.

Fig. 11 shows the nucleotide sequence of HCV cDNA in clone CA290a, the amino acids encoded therein, and the overlap with clone CA216a.

Fig. 12 shows the nucleotide sequence of HCV cDNA in clone ag30a and the overlap with clone CA290a.

Fig. 13 shows the nucleotide sequence of HCV cDNA in clone CA205a, and the overlap with the HCV cDNA sequence in clone CA290a.

Fig. 14 shows the nucleotide sequence of HCV cDNA in clone 18g, and the overlap with the HCV cDNA sequence in clone ag30a.

Fig. 15 shows the nucleotide sequence of HCV cDNA in clone 16jh, the amino acids encoded therein, and the overlap of nucleotides with the HCV cDNA sequence in clone 15e.

Fig. 16 shows the ORF of HCV cDNA derived from clones p114a, CA167b, CA156e, CA84a, CA59a, K9-1, 12f, 14i, 11b, 7f, 7e, 8h, 33c, 40b, 37b, 35, 36, 81, 32, 33b, 25c, 14c, 8f, 33f, 33g, 39c, 35f, 19g, 26g, and 15e.

Fig. 17 shows the sense strand of the compiled HCV cDNA sequence derived from the above-described clones and the compiled HCV cDNA sequence published in EPO Pub. No. 318,216. The clones from which the sequence was derived are b114a, 18g, ag30a, CA205a, CA290a, CA216a, p114a, CA167b, CA156e, CA84a, CA59a, K9-1 (also called k9-1), 26j, 13i, 12f, 14i, 11b, 7f, 7e, 8h, 33c, 40b, 37b, 35, 36, 81, 32, 33b, 25c, 14c, 8f, 33f, 33g, 39c, 35f, 19g, 26g, 15e, b5a, and 16jh. In the figure the three horizontal dashes above the sequence indicate the position

of the putative initiator methionine codon; the two vertical dashes indicate the first and last nucleotides of the published sequence. Also shown in the figure is the amino acid sequence of the putative polyprotein encoded in the HCV cDNA.

Fig. 18 is a diagram of the immunological colony screening method used in antigenic mapping studies.

Fig. 19 shows the hydrophobicity profiles of polyproteins encoded in HCV and in West Nile virus.

Fig. 20 is a tracing of the hydrophilicity/hydrophobicity profile and of the antigenic index of the putative HCV polyprotein.

Fig. 21 shows the conserved co-linear peptides in HCV and Flaviviruses.

Modes for Carrying Out the Invention

I. Definitions

The term "hepatitis C virus" has been reserved by workers in the field for an heretofore unknown etiologic agent of NANBH. Accordingly, as used herein, "hepatitis C virus" (HCV) refers to an agent causative of NANBH, which was formerly referred to as NANBV and/or BB-NANBV. The terms HCV, NANBV, and BB-NANBV are used interchangeably herein. As an extension of this terminology, the disease caused by HCV, formerly called NANB hepatitis (NANBH), is called hepatitis C. The terms NANBH and hepatitis C may be used interchangeably herein.

The term "HCV", as used herein, denotes a viral species of which pathogenic strains cause NANBH, and attenuated strains or defective interfering particles derived therefrom. As shown infra., the HCV genome is comprised of RNA. It is known that RNA containing viruses have relatively high rates of spontaneous mutation, i.e., reportedly on the order of 10^{-3} to 10^{-4} per incorporated

nucleotide (Fields & Knipe (1986)). Therefore, there are multiple strains, which may be virulent or avirulent, within the HCV species described infra. The compositions and methods described herein, enable the propagation, identification, detection, and isolation of the various HCV strains or isolates. Moreover, the disclosure herein allows the preparation of diagnostics and vaccines for the various strains, as well as compositions and methods that have utility in screening procedures for anti-viral agents for pharmacologic use, such as agents that inhibit replication of HCV.

The information provided herein, although derived from the prototype strain or isolate of HCV, hereinafter referred to as CDC/HCV1 (also called HCV1), is sufficient to allow a viral taxonomist to identify other strains which fall within the species. The information provided herein allows the belief that HCV is a Flavi-like virus. The morphology and composition of Flavivirus particles are known, and are discussed in Brinton (1986). Generally, with respect to morphology, Flaviviruses contain a central nucleocapsid surrounded by a lipid bilayer. Virions are spherical and have a diameter of about 40-50 nm. Their cores are about 25-30 nm in diameter. Along the outer surface of the virion envelope are projections that are about 5-10 nm long with terminal knobs about 2 nm in diameter.

Different strains or isolates of HCV are expected to contain variations at the amino acid and nucleic acids compared with the prototype isolate, HCV1. Many isolates are expected to show much (i.e. more than about 40%) homology in the total amino acid sequence compared with HCV1. However, it may also be found that other less homologous HCV isolates. These would be defined as HCV strains according to various criteria such as an ORF of approximately 9,000 nucleotides to ap-

proximately 12,000 nucleotides, encoding a polyprotein similar in size to that of HCV1, an encoded polyprotein of similar hydrophobic and antigenic character to that of HCV1, and the presence of co-linear peptide sequences that are conserved with HCV1. In addition, the genome would be a positive-stranded RNA.

HCV encodes at least one epitope which is immunologically identifiable with an epitope in the HCV genome from which the cDNAs described herein are derived; preferably the epitope is contained an amino acid sequence described herein. The epitope is unique to HCV when compared to other known Flaviviruses. The uniqueness of the epitope may be determined by its immunological reactivity with anti-HCV antibodies and lack of immunological reactivity with antibodies to other Flavivirus species. Methods for determining immunological reactivity are known in the art, for example, by radioimmunoassay, by Elisa assay, by hemagglutination, and several examples of suitable techniques for assays are provided herein.

In addition to the above, the following parameters of nucleic acid homology and amino acid homology are applicable, either alone or in combination, in identifying a strain or isolate as HCV. Since HCV strains and isolates are evolutionarily related, it is expected that the overall homology of the genomes at the nucleotide level probably will be about 40% or greater, probably about 60% or greater, and even more probably about 80% or greater; and in addition that there will be corresponding contiguous sequences of at least about 13 nucleotides. The correspondence between the putative HCV strain genomic sequence and the CDC/HCV1 cDNA sequence can be determined by techniques known in the art. For example, they can be determined by a direct comparison of the sequence information of the polynucleotide from the putative HCV, and the HCV cDNA sequence(s) described herein. For example, also,

they can be determined by hybridization of the polynucleotides under conditions which form stable duplexes between homologous regions (for example, those
5 which would be used prior to S_1 digestion), followed by digestion with single stranded specific nuclease(s), followed by size determination of the digested fragments.

Because of the evolutionary relationship of the strains or isolates of HCV, putative HCV strains or
10 isolates are identifiable by their homology at the polypeptide level. Generally, HCV strains or isolates are expected to be more than about 40% homologous, probably more than about 70% homologous, and even more probably more than about 80% homologous, and some may even be more
15 than about 90% homologous at the polypeptide level. The techniques for determining amino acid sequence homology are known in the art. For example, the amino acid sequence may be determined directly and compared to the sequences provided herein. Alternatively the nucleotide
20 sequence of the genomic material of the putative HCV may be determined (usually via a cDNA intermediate), the amino acid sequence encoded therein can be determined, and the corresponding regions compared.

As used herein, a polynucleotide "derived from"
25 a designated sequence refers to a polynucleotide sequence which is comprised of a sequence of approximately at least about 6 nucleotides, preferably at least about 8 nucleotides, more preferably at least about 10-12
30 nucleotides, and even more preferably at least about 15-20 nucleotides corresponding to a region of the designated nucleotide sequence. "Corresponding" means homologous to or complementary to the designated sequence. Preferably, the sequence of the region from which the polynucleotide
35 is derived is homologous to or complementary to a sequence which is unique to an HCV genome. Whether or not a sequence is unique to the HCV genome can be determined by

techniques known to those of skill in the art. For example, the sequence can be compared to sequences in databanks, e.g., Genbank, to determine whether it is present in the uninfected host or other organisms. The sequence can also be compared to the known sequences of other viral agents, including those which are known to induce hepatitis, e.g., HAV, HBV, and HDV, and to other members of the Flaviviridae. The correspondence or non-correspondence of the derived sequence to other sequences can also be determined by hybridization under the appropriate stringency conditions. Hybridization techniques for determining the complementarity of nucleic acid sequences are known in the art, and are discussed infra. See also, for example, Maniatis et al. (1982). In addition, mismatches of duplex polynucleotides formed by hybridization can be determined by known techniques, including for example, digestion with a nuclease such as S1 that specifically digests single-stranded areas in duplex polynucleotides. Regions from which typical DNA sequences may be "derived" include but are not limited to, for example, regions encoding specific epitopes, as well as non-transcribed and/or non-translated regions.

The derived polynucleotide is not necessarily physically derived from the nucleotide sequence shown, but may be generated in any manner, including for example, chemical synthesis or DNA replication or reverse transcription or transcription. In addition, combinations of regions corresponding to that of the designated sequence may be modified in ways known in the art to be consistent with an intended use.

Similarly, a polypeptide or amino acid sequence "derived from" a designated nucleic acid sequence refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least

3-5 amino acids, and more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence.

A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence, for example, the HCV cDNA sequences described herein, or from an HCV genome; it may be generated in any manner, including for example, chemical synthesis, or expression of a recombinant expression system, or isolation from mutated HCV. A recombinant or derived polypeptide may include one or more analogs of amino acids or unnatural amino acids in its sequence. Methods of inserting analogs of amino acids into a sequence are known in the art. It also may include one or more labels, which are known to those of skill in the art.

The term "recombinant polynucleotide" as used herein intends a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation which: (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature.

The term "polynucleotide" as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule. Thus, this term includes double- and single-stranded DNA, as well as double- and single stranded RNA. It also includes known types of modifications, for example, labels which are known in the art, methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl

phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those
5 containing pendant moieties, such as, for example proteins (including for e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing
10 chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide.

The term "purified viral polynucleotide" refers to
15 an HCV genome or fragment thereof which is essentially free, i.e., contains less than about 50%, preferably less than about 70%, and even more preferably less than about 90% of polypeptides with which the viral polynucleotide is naturally associated. Techniques for purifying viral
20 polynucleotides from viral particles are known in the art, and include for example, disruption of the particle with a chaotropic agent, differential extraction and separation of the polynucleotide(s) and polypeptides by ion-exchange chromatography, affinity chromatography, and sedimentation
25 according to density.

The term "purified viral polypeptide" refers to an HCV polypeptide or fragment thereof which is essentially free, i.e., contains less than about 50%,
30 preferably less than about 70%, and even more preferably less than about 90%, of cellular components with which the viral polypeptide is naturally associated. Techniques for purifying viral polypeptides are known in the art, and examples of these techniques are discussed infra. The term
35 "purified viral polynucleotide" refers to an HCV genome or fragment thereof which is essentially free, i.e., contains less than about 20%, preferably less than about 50%, and

even more preferably less than about 70% of polypeptides with which the viral polynucleotide is naturally associated. Techniques for purifying viral polynucleotides from viral particles are known in the art, and include for example, disruption of the particle with a chaotropic agent, and separation of the polynucleotide(s) and polypeptides by ion-exchange chromatography, affinity chromatography, and sedimentation according to density.

"Recombinant host cells", "host cells", "cells", "cell lines", "cell cultures", and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be, or have been, used as recipients for recombinant vector or other transfer DNA, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

A "replicon" is any genetic element, e.g., a plasmid, a chromosome, a virus, a cosmid, etc. that behaves as an autonomous unit of polynucleotide replication within a cell; i.e., capable of replication under its own control.

A "vector" is a replicon in which another polynucleotide segment is attached, so as to bring about the replication and/or expression of the attached segment.

"Control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and terminators; in eukaryotes, generally, such control sequences include promoters, terminators and, in some instances, enhancers.

The term "control sequences" is intended to include, at a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences.

"Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

An "open reading frame" (ORF) is a region of a polynucleotide sequence which encodes a polypeptide; this region may represent a portion of a coding sequence or a total coding sequence.

A "coding sequence" is a polynucleotide sequence which is transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. A coding sequence can include, but is not limited to mRNA, cDNA, and recombinant polynucleotide sequences.

"Immunologically identifiable with/as" refers to the presence of epitope(s) and polypeptides(s) which are also present in the designated polypeptide(s), usually HCV proteins. Immunological identity may be determined by antibody binding and/or competition in binding; these techniques are known to those of average skill in the art, and are also illustrated infra.

As used herein, "epitope" refers to an antigenic determinant of a polypeptide; an epitope could comprise 3 amino acids in a spatial conformation which is unique to

the epitope, generally an epitope consists of at least 5 such amino acids, and more usually, consists of at least 8-10 such amino acids. Methods of determining the spatial
5 conformation of amino acids are known in the art, and include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance.

A polypeptide is "immunologically reactive" with an antibody when it binds to an antibody due to antibody
10 recognition of a specific epitope contained within the polypeptide. Immunological reactivity may be determined by antibody binding, more particularly by the kinetics of antibody binding, and/or by competition in binding using as competitor(s) a known polypeptide(s) containing an
15 epitope against which the antibody is directed. The techniques for determining whether a polypeptide is immunologically reactive with an antibody are known in the art.

As used herein, the term "immunogenic
20 polypeptide" is a polypeptide that elicits a cellular and/or humoral response, whether alone or linked to a carrier in the presence or absence of an adjuvant.

The term "polypeptide" refers to a polymer of amino acids and does not refer to a specific length of the
25 product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the
30 like. Included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.
35

"Transformation", as used herein, refers to the insertion of an exogenous polynucleotide into a host cell, irrespective of the method used for the insertion, for example, direct uptake, transduction, f-mating or electroporation. The exogenous polynucleotide may be maintained as a non-integrated vector, for example, a plasmid, or alternatively, may be integrated into the host genome.

"Treatment" as used herein refers to prophylaxis and/or therapy.

An "individual", as used herein, refers to vertebrates, particularly members of the mammalian species, and includes but is not limited to domestic animals, sports animals, and primates, including humans.

As used herein, the "sense strand" of a nucleic acid contains the sequence that has sequence homology to that of mRNA. The "anti-sense strand" contains a sequence which is complementary to that of the "sense strand".

As used herein, a "positive stranded genome" of a virus is one in which the genome, whether RNA or DNA, is single-stranded and which encodes a viral polypeptide(s). Examples of positive stranded RNA viruses include Togaviridae, Coronaviridae, Retroviridae, Picornaviridae, and Caliciviridae. Included also, are the Flaviviridae, which were formerly classified as Togaviridae. See Fields & Knipe (1986).

As used herein, "antibody-containing body component" refers to a component of an individual's body which is a source of the antibodies of interest. Antibody containing body components are known in the art, and include but are not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, white blood cells, and myelomas.

As used herein, "purified HCV" refers to a preparation of HCV which has been isolated from the cellular constituents with which the virus is normally associated, and from other types of viruses which may be present in the infected tissue. The techniques for isolating viruses are known to those of skill in the art, and include, for example, centrifugation and affinity chromatography; a method of preparing purified HCV is discussed infra.

The term "HCV particles" as used herein include entire virion as well as particles which are intermediates in virion formation. HCV particles generally have one or more HCV proteins associated with the HCV nucleic acid.

As used herein, the term "probe" refers to a polynucleotide which forms a hybrid structure with a sequence in a target region, due to complementarity of at least one sequence in the probe with a sequence in the target region. The probe, however, does not contain a sequence complementary to sequence(s) used to prime the polymerase chain reaction.

As used herein, the term "target region" refers to a region of the nucleic acid which is to be amplified and/or detected.

As used herein, the term "viral RNA", which includes HCV RNA, refers to RNA from the viral genome, fragments thereof, transcripts thereof, and mutant sequences derived therefrom.

As used herein, a "biological sample" refers to a sample of tissue or fluid isolated from an individual, including but not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, tumors, organs, and also samples of in vitro cell culture constituents (including but not limited to conditioned medium resulting from the

growth of cells in cell culture medium, putatively virally infected cells, recombinant cells, and cell components).

5 II. Description of the Invention

 The practice of the present invention will
employ, unless otherwise indicated, conventional
techniques of molecular biology, microbiology, recombinant
DNA, and immunology, which are within the skill of the
10 art. Such techniques are explained fully in the
literature. See e.g., Maniatis, Fitch & Sambrook,
MOLECULAR CLONING; A LABORATORY MANUAL (1982); DNA CLON-
ING, VOLUMES I AND II (D.N Glover ed. 1985);
15 OLIGONUCLEOTIDE SYNTHESIS (M.J. Gait ed, 1984); NUCLEIC
ACID HYBRIDIZATION (B.D. Hames & S.J. Higgins eds. 1984);
TRANSCRIPTION AND TRANSLATION (B.D. Hames & S.J. Higgins
eds. 1984); ANIMAL CELL CULTURE (R.I. Freshney ed. 1986);
IMMOBILIZED CELLS AND ENZYMES (IRL Press, 1986); B.
20 Perbal, A PRACTICAL GUIDE TO MOLECULAR CLONING (1984);
the series, METHODS IN ENZYMOLOGY (Academic Press, Inc.);
GENE TRANSFER VECTORS FOR MAMMALIAN CELLS (J.H. Miller and
M.P. Calos eds. 1987, Cold Spring Harbor Laboratory),
Methods in Enzymology Vol. 154 and Vol. 155 (Wu and
25 Grossman, and Wu, eds., respectively), Mayer and Walker,
eds. (1987), IMMUNOCHEMICAL METHODS IN CELL AND MOLECULAR
BIOLOGY (Academic Press, London), Scopes, (1987), PROTEIN
PURIFICATION: PRINCIPLES AND PRACTICE, Second Edition
(Springer-Verlag, N.Y.), and HANDBOOK OF EXPERIMENTAL IM-
30 MUNOLOGY, VOLUMES I-IV (D.M. Weir and C. C. Blackwell eds
1986). All patents, patent applications, and publications
mentioned herein, both supra and infra, are hereby in-
corporated herein by reference.

 The useful materials and processes of the
35 present invention are made possible by the provision of a
family of nucleotide sequences isolated from cDNA librar-
ies which contain HCV cDNA sequences. These cDNA librar-

ies were derived from nucleic acid sequences present in the plasma of an HCV-infected chimpanzee. The construction of one of these libraries, the "c" library (ATCC No. 40394), was reported in EPO Pub. No. 318,216. Several of the clones containing HCV cDNA reported herein were obtained from the "c" library. Although other clones reported herein were obtained from other HCV cDNA libraries, the presence of clones containing the sequences in the "c" library was confirmed. As discussed in EPO Pub. No. 318,216, the family of HCV cDNA sequences isolated from the "c" library are not of human or chimpanzee origin, and show no significant homology to sequences contained within the HBV genome.

The availability of the HCV cDNAs described herein permits the construction of polynucleotide probes which are reagents useful for detecting viral polynucleotides in biological samples, including donated blood. For example, from the sequences it is possible to synthesize DNA oligomers of about 8-10 nucleotides, or larger, which are useful as hybridization probes to detect the presence of HCV RNA in, for example, donated blood, sera of subjects suspected of harboring the virus, or cell culture systems in which the virus is replicating. In addition, the cDNA sequences also allow the design and production of HCV specific polypeptides which are useful as diagnostic reagents for the presence of antibodies raised during HCV infection. Antibodies to purified polypeptides derived from the cDNAs may also be used to detect viral antigens in biological samples, including, for example, donated blood samples, sera from patients with NANBH, and in tissue culture systems being used for HCV replication. Moreover, the immunogenic polypeptides disclosed herein, which are encoded in portions of the ORF of HCV cDNA shown in Fig. 17, are also useful for HCV

screening, diagnosis, and treatment, and for raising antibodies which are also useful for these purposes.

5 In addition, the novel cDNA sequences described herein enable further characterization of the HCV genome. Polynucleotide probes and primers derived from these sequences may be used to amplify sequences present in cDNA libraries, and/or to screen cDNA libraries for additional
10 overlapping cDNA sequences, which, in turn, may be used to obtain more overlapping sequences. As indicated infra. and in EPO Pub. No. 318,216, the genome of HCV appears to be RNA comprised primarily of a large open reading frame (ORF) which encodes a large polyprotein.

15 The HCV cDNA sequences provided herein, the polypeptides derived from these sequences, and the immunogenic polypeptides described herein, as well as antibodies directed against these polypeptides are also useful in the isolation and identification of the blood-borne NABV (BB-NANBV) agent(s). For example, antibodies
20 directed against HCV epitopes contained in polypeptides derived from the cDNAs may be used in processes based upon affinity chromatography to isolate the virus. Alternatively, the antibodies may be used to identify
25 viral particles isolated by other techniques. The viral antigens and the genomic material within the isolated viral particles may then be further characterized.

In addition to the above, the information provided infra allows the identification of additional HCV strains or isolates. The isolation and characterization
30 of the additional HCV strains or isolates may be accomplished by isolating the nucleic acids from body components which contain viral particles and/or viral RNA, creating cDNA libraries using polynucleotide probes based on the HCV cDNA probes described infra., screening the
35 libraries for clones containing HCV cDNA sequences described infra., and comparing the HCV cDNAs from the new

isolates with the cDNAs described infra. The polypeptides encoded therein, or in the viral genome, may be monitored for immunological cross-reactivity utilizing the
5 polypeptides and antibodies described supra. Strains or isolates which fit within the parameters of HCV, as described in the Definitions section, supra., are readily identifiable. Other methods for identifying HCV strains
10 will be obvious to those of skill in the art, based upon the information provided herein.

Isolation of the HCV cDNA Sequences

The novel HCV cDNA sequences described infra. extend the sequence of the cDNA to the HCV genome reported
15 in EPO Pub. No. 318,216. The sequences which are present in clones b114a, 18g, ag30a, CA205a, CA290a, CA216a, p114a, CA167b, CA156e, CA84a, and CA59a lie upstream of the reported sequence, and when compiled, yield
20 nucleotides nos. -319 to 1348 of the composite HCV cDNA sequence. (The negative number on a nucleotide indicates its distance upstream of the nucleotide which starts the putative initiator MET codon.) The sequences which are present in clones b5a and 16jh lie downstream of the
25 reported sequence, and yield nucleotides nos. 8659 to 8866 of the composite sequence. The composite HCV cDNA sequence which includes the sequences in the aforementioned clones, is shown in Fig. 17.

The novel HCV cDNAs described herein were isolated from a number of HCV cDNA libraries, including
30 the "c" library present in lambda gt11 (ATCC No. 40394). The HCV cDNA libraries were constructed using pooled serum from a chimpanzee with chronic HCV infection and containing a high titer of the virus, i.e., at least 10^6 chimp infectious doses/ml (CID/ml). The pooled serum was used
35 to isolate viral particles; nucleic acids isolated from these particles was used as the template in the construc-

tion of cDNA libraries to the viral genome. The procedures for isolation of putative HCV particles and for constructing the "c" HCV cDNA library is described in EPO Pub. No. 318,216. Other methods for constructing HCV cDNA libraries are known in the art, and some of these methods are described infra., in the Examples. Isolation of the sequences was by screening the libraries using synthetic polynucleotide probes, the sequences of which were derived from the 5'-region and the 3'-region of the known HCV cDNA sequence. The description of the method to retrieve the cDNA sequences is mostly of historical interest. The resultant sequences (and their complements) are provided herein, and the sequences, or any portion thereof, could be prepared using synthetic methods, or by a combination of synthetic methods with retrieval of partial sequences using methods similar to those described herein.

Preparation of Viral Polypeptides and Fragments

The availability of HCV cDNA sequences, or nucleotide sequences derived therefrom (including segments and modifications of the sequence), permits the construction of expression vectors encoding antigenically active regions of the polypeptide encoded in either strand. These antigenically active regions may be derived from coat or envelope antigens or from core antigens, or from antigens which are non-structural including, for example, polynucleotide binding proteins, polynucleotide polymerase(s), and other viral proteins required for the replication and/or assembly of the virus particle. Fragments encoding the desired polypeptides are derived from the cDNA clones using conventional restriction digestion or by synthetic methods, and are ligated into vectors which may, for example, contain portions of fusion sequences such as beta-galactosidase or superoxide dismutase (SOD), preferably SOD. Methods and vectors

which are useful for the production of polypeptides which contain fusion sequences of SOD are described in European Patent Office Publication number 0196056, published
5 October 1, 1986. Vectors for the expression of fusion polypeptides of SOD and HCV polypeptides encoded in a number of HCV clones are described infra., in the Examples. Any desired portion of the HCV cDNA containing an open reading frame, in either sense strand, can be
10 obtained as a recombinant polypeptide, such as a mature or fusion protein; alternatively, a polypeptide encoded in the cDNA can be provided by chemical synthesis.

The DNA encoding the desired polypeptide, whether in fused or mature form, and whether or not
15 containing a signal sequence to permit secretion, may be ligated into expression vectors suitable for any convenient host. Both eukaryotic and prokaryotic host systems are presently used in forming recombinant polypeptides, and a summary of some of the more common
20 control systems and host cell lines is given infra. The polypeptide is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use. Purification may be by techniques known in the art, for example, differential extraction, salt
25 fractionation, chromatography on ion exchange resins, affinity chromatography, centrifugation, and the like. See, for example, Methods in Enzymology for a variety of methods for purifying proteins. Such polypeptides can be used as diagnostics, or those which give rise to neutral-
30 izing antibodies may be formulated into vaccines. Antibodies raised against these polypeptides can also be used as diagnostics, or for passive immunotherapy. In addition, as discussed infra., antibodies to these
35 polypeptides are useful for isolating and identifying HCV particles.

Preparation of Antigenic Polypeptides and Conjugation with Carrier

5 An antigenic region of a polypeptide is generally relatively small--typically 8 to 10 amino acids or less in length. Fragments of as few as 5 amino acids may characterize an antigenic region. These segments may correspond to regions of HCV antigen. Accordingly, using the
10 cDNAs of HCV as a basis, DNAs encoding short segments of HCV polypeptides can be expressed recombinantly either as fusion proteins, or as isolated polypeptides. In addition, short amino acid sequences can be conveniently obtained by chemical synthesis. In instances wherein the
15 synthesized polypeptide is correctly configured so as to provide the correct epitope, but is too small to be immunogenic, the polypeptide may be linked to a suitable carrier.

A number of techniques for obtaining such linkage are known in the art, including the formation of
20 disulfide linkages using N-succinimidyl-3-(2-pyridylthio)propionate (SPDP) and succinimidyl 4-(N-maleimido-methyl)cyclohexane-1-carboxylate (SMCC) obtained from Pierce Company, Rockford, Illinois, (if the peptide lacks a
25 sulfhydryl group, this can be provided by addition of a cysteine residue.) These reagents create a disulfide linkage between themselves and peptide cysteine residues on one protein and an amide linkage through the epsilon-amino on a lysine, or other free amino group in the other. A variety of such disulfide/amide-forming agents are
30 known. See, for example, Immun. Rev. (1982) 62:185. Other bifunctional coupling agents form a thioether rather than a disulfide linkage. Many of these thio-ether-forming agents are commercially available and include reactive esters of 6-maleimidocaproic acid, 2-bromoacetic
35 acid, 2-iodoacetic acid, 4-(N-maleimido-methyl)cyclohexane-1-carboxylic acid, and the like. The

carboxyl groups can be activated by combining them with succinimide or 1-hydroxyl-2-nitro-4-sulfonic acid, sodium salt. Additional methods of coupling antigens employs the
5 rotavirus/"binding peptide" system described in EPO Pub. No. 259,149, the disclosure of which is incorporated herein by reference. The foregoing list is not meant to be exhaustive, and modifications of the named compounds can clearly be used.

10 Any carrier may be used which does not itself induce the production of antibodies harmful to the host. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins; polysaccharides, such as
15 latex functionalized sepharose, agarose, cellulose, cellulose beads and the like; polymeric amino acids, such as polyglutamic acid, polylysine, and the like; amino acid copolymers; and inactive virus particles. Especially useful protein substrates are serum albumins, keyhole limpet
20 hemocyanin, immunoglobulin molecules, thyroglobulin, ovalbumin, tetanus toxoid, and other proteins well known to those skilled in the art.

In addition to full-length viral proteins, polypeptides comprising truncated HCV amino acid sequences encoding at least one viral epitope are useful im-
25 munological reagents. For example, polypeptides comprising such truncated sequences can be used as reagents in an immunoassay. These polypeptides also are candidate subunit antigens in compositions for antiserum production or vaccines. While these truncated sequences can be
30 produced by various known treatments of native viral protein, it is generally preferred to make synthetic or recombinant polypeptides comprising an HCV sequence. Polypeptides comprising these truncated HCV sequences can
35 be made up entirely of HCV sequences (one or more epitopes, either contiguous or noncontiguous), or HCV sequences and heterologous sequences in a fusion protein.

Useful heterologous sequences include sequences that provide for secretion from a recombinant host, enhance the immunological reactivity of the HCV epitope(s), or facilitate the coupling of the polypeptide to an immunoassay support or a vaccine carrier. See, e.g., EPO Pub. No. 116,201; U.S. Pat. No. 4,722,840; EPO Pub. No. 259,149; U.S. Pat. No. 4,629,783, the disclosures of which are incorporated herein by reference.

The size of polypeptides comprising the truncated HCV sequences can vary widely, the minimum size being a sequence of sufficient size to provide an HCV epitope, while the maximum size is not critical. For convenience, the maximum size usually is not substantially greater than that required to provide the desired HCV epitopes and function(s) of the heterologous sequence, if any. Typically, the truncated HCV amino acid sequence will range from about 5 to about 100 amino acids in length. More typically, however, the HCV sequence will be a maximum of about 50 amino acids in length, preferably a maximum of about 30 amino acids. It is usually desirable to select HCV sequences of at least about 10, 12 or 15 amino acids, up to a maximum of about 20 or 25 amino acids.

Truncated HCV amino acid sequences comprising epitopes can be identified in a number of ways. For example, the entire viral protein sequence can be screened by preparing a series of short peptides that together span the entire protein sequence. An example of antigenic screening of the regions of the HCV polyprotein is shown infra. In addition, by starting with, for example, 100mer polypeptides, it would be routine to test each polypeptide for the presence of epitope(s) showing a desired reactivity, and then testing progressively smaller and overlapping fragments from an identified 100mer to map the epitope of interest. Screening such peptides in an

immunoassay is within the skill of the art. It is also known to carry out a computer analysis of a protein sequence to identify potential epitopes, and then prepare oligopeptides comprising the identified regions for screening. Such a computer analysis of the HCV amino acid sequence is shown in Fig. 20, where the hydrophilic/hydrophobic character is displayed above the antigen index. The amino acids are numbered from the starting MET (position 1) as shown in Fig. 17. It is appreciated by those of skill in the art that such computer analysis of antigenicity does not always identify an epitope that actually exists, and can also incorrectly identify a region of the protein as containing an epitope.

Examples of HCV amino acid sequences that may be useful, which are expressed from expression vectors comprised of clones 5-1-1, 81, CA74a, 35f, 279a, C36, C33b, CA290a, C8f, C12f, 14c, 15e, C25c, C33c, C33f, 33g, C39c, C40b, CA167b are described infra. Other examples of HCV amino acid sequences that may be useful as described herein are set forth below. It is to be understood that these peptides do not necessarily precisely map one epitope, and may also contain HCV sequence that is not immunogenic. These non-immunogenic portions of the sequence can be defined as described above using conventional techniques and deleted from the described sequences. Further, additional truncated HCV amino acid sequences that comprise an epitope or are immunogenic can be identified as described above. The following sequences are given by amino acid number (i.e., "AAn") where n is the amino acid number as shown in Fig. 17:

AA1-AA25; AA1-AA50; AA1-AA84; AA9-AA177; AA1-AA10;
AA5-AA20; AA20-AA25; AA35-AA45; AA50-AA100;
AA40-AA90; AA45-AA65; AA65-AA75; AA80-90; AA99-AA120;
AA95-AA110; AA105-AA120; AA100-AA150; AA150-AA200;
AA155-AA170; AA190-AA210; AA200-AA250; AA220-AA240;

5 AA245-AA265; AA250-AA300; AA290-AA330; AA290-305;
AA300-AA350; AA310-AA330; AA350-AA400; AA380-AA395;
AA405-AA495; AA400-AA450; AA405-AA415; AA415-AA425;
AA425-AA435; AA437-AA582; AA450-AA500; AA440-AA460;
AA460-AA470; AA475-AA495; AA500-AA550; AA511-AA690;
AA515-AA550; AA550-AA600; AA550-AA625; AA575-AA605;
AA585-AA600; AA600-AA650; AA600-AA625; AA635-AA665;
AA650-AA700; AA645-AA680; AA700-AA750; AA700-AA725;
AA700-AA750; AA725-AA775; AA770-AA790; AA750-AA800;
AA800-AA815; AA825-AA850; AA850-AA875; AA800-AA850;
AA920-AA990; AA850-AA900; AA920-AA945; AA940-AA965;
10 AA970-AA990; AA950-AA1000; AA1000-AA1060;
AA1000-AA1025; AA1000-AA1050; AA1025-AA1040;
AA1040-AA1055; AA1075-AA1175; AA1050-AA1200;
AA1070-AA1100; AA1100-AA1130; AA1140-AA1165;
AA1192-AA1457; AA1195-AA1250; AA1200-AA1225;
AA1225-AA1250; AA1250-AA1300; AA1260-AA1310;
AA1260-AA1280; AA1266-AA1428; AA1300-AA1350;
AA1290-AA1310; AA1310-AA1340; AA1345-AA1405;
15 AA1345-AA1365; AA1350-AA1400; AA1365-AA1380;
AA1380-AA1405; AA1400-AA1450; AA1450-AA1500;
AA1460-AA1475; AA1475-AA1515; AA1475-AA1500;
AA1500-AA1550; AA1500-AA1515; AA1515-AA1550;
AA1550-AA1600; AA1545-AA1560; AA1569-AA1931;
AA1570-AA1590; AA1595-AA1610; AA1590-AA1650;
AA1610-AA1645; AA1650-AA1690; AA1685-AA1770;
20 AA1689-AA1805; AA1690-AA1720; AA1694-AA1735;
AA1720-AA1745; AA1745-AA1770; AA1750-AA1800;
AA1775-AA1810; AA1795-AA1850; AA1850-AA1900;
AA1900-AA1950; AA1900-AA1920; AA1916-AA2021;
AA1920-AA1940; AA1949-AA2124; AA1950-AA2000;
AA1950-AA1985; AA1980-AA2000; AA2000-AA2050;
AA2005-AA2025; AA2020-AA2045; AA2045-AA2100;
AA2045-AA2070; AA2054-AA2223; AA2070-AA2100;
25 AA2100-AA2150; AA2150-AA2200; AA2200-AA2250;
AA2200-AA2325; AA2250-AA2330; AA2255-AA2270;
AA2265-AA2280; AA2280-AA2290; AA2287-AA2385;
AA2300-AA2350; AA2290-AA2310; AA2310-AA2330;
AA2330-AA2350; AA2350-AA2400; AA2348-AA2464;
AA2345-AA2415; AA2345-AA2375; AA2370-AA2410;
AA2371-AA2502; AA2400-AA2450; AA2400-AA2425;
30 AA2415-AA2450; AA2445-AA2500; AA2445-AA2475;
AA2470-AA2490; AA2500-AA2550; AA2505-AA2540;
AA2535-AA2560; AA2550-AA2600; AA2560-AA2580;
AA2600-AA2650; AA2605-AA2620; AA2620-AA2650;
AA2640-AA2660; AA2650-AA2700; AA2655-AA2670;
AA2670-AA2700; AA2700-AA2750; AA2740-AA2760;
AA2750-AA2800; AA2755-AA2780;
AA2780-AA2830; AA2785-AA2810; AA2796-AA2886;
35 AA2810-AA2825; AA2800-AA2850; AA2850-AA2900;
AA2850-AA2865; AA2885-AA2905; AA2900-AA2950;

AA2910-AA2930; AA2925-AA2950; AA2945-end(C' terminal).

5 The above HCV amino acid sequences can be prepared as discrete peptides or incorporated into a larger polypeptide, and may find use as described herein. Additional polypeptides comprising truncated HCV sequences are described in the examples.

10 The observed relationship of the putative polyproteins of HCV and the Flaviviruses allows some prediction of the putative domains of the HCV "non-structural" (NS) proteins. The locations of the individual NS proteins in the putative Flavivirus precursor polyprotein are fairly well-known. Moreover, these
15 also coincide with observed gross fluctuations in the hydrophobicity profile of the polyprotein. It is established that NS5 of Flaviviruses encodes the virion polymerase, and that NS1 corresponds with a complement fixation antigen which has been shown to be an effective
20 vaccine in animals. Recently, it has been shown that a flaviviral protease function resides in NS3. Due to the observed similarities between HCV and the Flaviviruses, described infra., deductions concerning the approximate locations of the corresponding protein domains and func-
25 tions in the HCV polyprotein are possible. The expression of polypeptides containing these domains in a variety of recombinant host cells, including, for example, bacteria, yeast, insect, and vertebrate cells, should give rise to important immunological reagents which can be used for
30 diagnosis, detection, and vaccines.

Although the non-structural protein regions of the putative polyproteins of the HCV isolate described herein and of Flaviviruses appear to have some similarity,
35 there is less similarity between the putative structural regions which are towards the N-terminus. In this region, there is a greater divergence in sequence, and in addi-

tion, the hydrophobic profile of the two regions show less similarity. This "divergence" begins in the N-terminal region of the putative NS1 domain in HCV, and extends to the presumed N-terminus. Nevertheless, it may still be possible to predict the approximate locations of the putative nucleocapsid (N-terminal basic domain) and E (generally hydrophobic) domains within the HCV polyprotein. In the Examples the predictions are based on the changes observed in the hydrophobic profile of the HCV polyprotein, and on a knowledge of the location and character of the flaviviral proteins. From these predictions it may be possible to identify approximate regions of the HCV polyprotein that could correspond with useful immunological reagents. For example, the E and NS1 proteins of Flaviviruses are known to have efficacy as protective vaccines. These regions, as well as some which are shown to be antigenic in the HCV isolate described herein, for example those within putative NS3, C, and NS5, etc., should also provide diagnostic reagents. Moreover, the location and expression of viral-encoded enzymes may also allow the evaluation of anti-viral enzyme inhibitors, i.e., for example, inhibitors which prevent enzyme activity by virtue of an interaction with the enzyme itself, or substances which may prevent expression of the enzyme, (for example, anti-sense RNA, or other drugs which interfere with expression).

Preparation of Hybrid Particle Immunogens Containing HCV Epitopes

The immunogenicity of the epitopes of HCV may also be enhanced by preparing them in mammalian or yeast systems fused with or assembled with particle-forming proteins such as, for example, that associated with hepatitis B surface antigen. Constructs wherein the NANBV epitope is linked directly to the particle-forming protein

coding sequences produce hybrids which are immunogenic with respect to the HCV epitope. In addition, all of the vectors prepared include epitopes specific to HBV, having various degrees of immunogenicity, such as, for example, the pre-S peptide. Thus, particles constructed from particle forming protein which include HCV sequences are immunogenic with respect to HCV and HBV.

Hepatitis surface antigen (HBSAg) has been shown to be formed and assembled into particles in S. cerevisiae (Valenzuela et al. (1982)), as well as in, for example, mammalian cells (Valenzuela, P., et al. (1984)). The formation of such particles has been shown to enhance the immunogenicity of the monomer subunit. The constructs may also include the immunodominant epitope of HBSAg, comprising the 55 amino acids of the presurface (pre-S) region. Neurath et al. (1984). Constructs of the pre-S-HBSAg particle expressible in yeast are disclosed in EPO 174,444, published March 19, 1986; hybrids including heterologous viral sequences for yeast expression are disclosed in EPO 175,261, published March 26, 1986. These constructs may also be expressed in mammalian cells such as Chinese hamster ovary (CHO) cells using an SV40-dihydrofolate reductase vector (Michelle et al. (1984)).

In addition, portions of the particle-forming protein coding sequence may be replaced with codons encoding an HCV epitope. In this replacement, regions which are not required to mediate the aggregation of the units to form immunogenic particles in yeast or mammals can be deleted, thus eliminating additional HBV antigenic sites from competition with the HCV epitope.

Preparation of Vaccines

Vaccines may be prepared from one or more immunogenic polypeptides derived from HCV cDNA, including the cDNA sequences described in the Examples. The

observed homology between HCV and Flaviviruses provides information concerning the polypeptides which may be most effective as vaccines, as well as the regions of the genome in which they are encoded. The general structure of the Flavivirus genome is discussed in Rice et al (1986). The flavivirus genomic RNA is believed to be the only virus-specific mRNA species, and it is translated into the three viral structural proteins, i.e., C, M, and E, as well as two large nonstructural proteins, NS4 and NS5, and a complex set of smaller nonstructural proteins. It is known that major neutralizing epitopes for Flaviviruses reside in the E (envelope) protein (Roehrig (1986)). Thus, vaccines may be comprised of recombinant polypeptides containing epitopes of HCV E. These polypeptides may be expressed in bacteria, yeast, or mammalian cells, or alternatively may be isolated from viral preparations. It is also anticipated that the other structural proteins may also contain epitopes which give rise to protective anti-HCV antibodies. Thus, polypeptides containing the epitopes of E, C, and M may also be used, whether singly or in combination, in HCV vaccines.

In addition to the above, it has been shown that immunization with NS1 (nonstructural protein 1), results in protection against yellow fever (Schlesinger et al (1986)). This is true even though the immunization does not give rise to neutralizing antibodies. Thus, particularly since this protein appears to be highly conserved among Flaviviruses, it is likely that HCV NS1 will also be protective against HCV infection. Moreover, it also shows that nonstructural proteins may provide protection against viral pathogenicity, even if they do not cause the production of neutralizing antibodies.

The information provided in the Examples concerning the immunogenicity of the polypeptides

expressed from cloned HCV cDNAs which span the various regions of the HCV ORF also allows predictions concerning their use in vaccines.

5 In view of the above, multivalent vaccines against HCV may be comprised of one or more epitopes from one or more structural proteins, and/or one or more epitopes from one or more nonstructural proteins. These vaccines may be comprised of, for example, recombinant HCV
10 polypeptides and/or polypeptides isolated from the virions. In particular, vaccines are contemplated comprising one or more of the following HCV proteins, or subunit antigens derived therefrom: E, NS1, C, NS2, NS3, NS4 and NS5. Particularly preferred are vaccines compris-
15 ing E and/or NS1, or subunits thereof.

The preparation of vaccines which contain an immunogenic polypeptide(s) as active ingredients, is known to one skilled in the art. Typically, such vaccines are
20 prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The preparation may also be emulsified, or the protein encapsulated in liposomes. The active immunogenic
25 ingredients are often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vac-
30 cine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine. Examples of adjuvants which may be effective include but are not limited to: aluminum hydroxide,
35 N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-

isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components
5 extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. The effectiveness of an adjuvant may be determined by measuring the amount of antibodies directed against an immunogenic polypeptide
10 containing an HCV antigenic sequence resulting from administration of this polypeptide in vaccines which are also comprised of the various adjuvants.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which
15 are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or
20 triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1%-2%. Oral formulations include such normally employed excipients as, for example,
25 pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and
30 contain 10%-95% of active ingredient, preferably 25%-70%.

The proteins may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with
35 inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids such as acetic, oxalic, tartaric, maleic, and the like. Salts formed with

the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic
5 bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

Dosage and Administration of Vaccines

10 The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be prophylactically and/or therapeutically effective. The quantity to be administered, which is generally
15 in the range of 5 micrograms to 250 micrograms of antigen per dose, depends on the subject to be treated, capacity of the subject's immune system to synthesize antibodies, and the degree of protection desired. Precise amounts of active ingredient required to be administered may depend
20 on the judgment of the practitioner and may be peculiar to each subject.

The vaccine may be given in a single dose schedule, or preferably in a multiple dose schedule. A multiple dose schedule is one in which a primary course of vaccination may be with 1-10 separate doses, followed by
25 other doses given at subsequent time intervals required to maintain and or reenforce the immune response, for example, at 1-4 months for a second dose, and if needed, a subsequent dose(s) after several months. The dosage regimen will also, at least in part, be determined by the
30 need of the individual and be dependent upon the judgment of the practitioner.

In addition, the vaccine containing the immunogenic HCV antigen(s) may be administered in conjunction with other immunoregulatory agents, for example, im-
35 mune globulins.

Preparation of Antibodies Against HCV Epitopes

The immunogenic polypeptides prepared as described above are used to produce antibodies, both polyclonal and monoclonal. If polyclonal antibodies are desired, a selected mammal (e.g., mouse, rabbit, goat, horse, etc.) is immunized with an immunogenic polypeptide bearing an HCV epitope(s). Serum from the immunized animal is collected and treated according to known procedures. If serum containing polyclonal antibodies to an HCV epitope contains antibodies to other antigens, the polyclonal antibodies can be purified by immunoaffinity chromatography. Techniques for producing and processing polyclonal antisera are known in the art, see for example, Mayer and Walker (1987).

Alternatively, polyclonal antibodies may be isolated from a mammal which has been previously infected with HCV. An example of a method for purifying antibodies to HCV epitopes from serum from an infected individual, based upon affinity chromatography and utilizing a fusion polypeptide of SOD and a polypeptide encoded within cDNA clone 5-1-1, is presented in EPO Pub. No. 318,216.

Monoclonal antibodies directed against HCV epitopes can also be readily produced by one skilled in the art. The general methodology for making monoclonal antibodies by hybridomas is well known. Immortal antibody-producing cell lines can be created by cell fusion, and also by other techniques such as direct transfection of B lymphocytes with oncogenic DNA, or transfection with Epstein-Barr virus. See, e.g., M. Schreier et al. (1980); Hammerling et al. (1981); Kennett et al. (1980); see also, U.S. Patent Nos. 4,341,761; 4,399,121; 4,427,783; 4,444,887; 4,466,917; 4,472,500; 4,491,632; and 4,493,890. Panels of monoclonal antibodies produced against HCV epitopes can be screened for various properties; i.e., for isotype, epitope affinity, etc.

Antibodies, both monoclonal and polyclonal, which are directed against HCV epitopes are particularly useful in diagnosis, and those which are neutralizing are useful in passive immunotherapy. Monoclonal antibodies, in particular, may be used to raise anti-idiotypic antibodies.

Anti-idiotypic antibodies are immunoglobulins which carry an "internal image" of the antigen of the infectious agent against which protection is desired. See, for example, Nisonoff, A., et al. (1981) and Dreesman et al. (1985).

Techniques for raising anti-idiotypic antibodies are known in the art. See, for example, Grzych (1985), MacNamara et al. (1984), and Uytdehaag et al. (1985). These anti-idiotypic antibodies may also be useful for treatment and/or diagnosis of NANBH, as well as for an elucidation of the immunogenic regions of HCV antigens.

It would also be recognized by one of ordinary skill in the art that a variety of types of antibodies directed against HCV epitopes may be produced. As used herein, the term "antibody" refers to a polypeptide or group of polypeptides which are comprised of at least one antibody combining site. An "antibody combining site" or "binding domain" is formed from the folding of variable domains of an antibody molecule(s) to form three-dimensional binding spaces with an internal surface shape and charge distribution complementary to the features of an epitope of an antigen, which allows an immunological reaction with the antigen. An antibody combining site may be formed from a heavy and/or a light chain domain (VH and VL, respectively), which form hypervariable loops which contribute to antigen binding. The term "antibody" includes, for example, vertebrate antibodies, hybrid antibodies, chimeric antibodies, altered antibodies, univalent

antibodies, the Fab proteins, and single domain antibodies.

5 A "single domain antibody" (dAb) is an antibody which is comprised of an VH domain, which reacts immunologically with a designated antigen. A dAb does not contain a VL domain, but may contain other antigen binding domains known to exist in antibodies, for example, the kappa and lambda domains. Methods for preparing dAbs are
10 known in the art. See, for example, Ward et al. (1989).

Antibodies may also be comprised of VH and VL domains, as well as other known antigen binding domains. Examples of these types of antibodies and methods for
15 their preparation are known in the art (see, e.g., U.S. Patent No. 4,816,467, which is incorporated herein by reference), and include the following. For example, "vertebrate antibodies" refers to antibodies which are tetramers or aggregates thereof, comprising light and
20 heavy chains which are usually aggregated in a "Y" configuration and which may or may not have covalent linkages between the chains. In vertebrate antibodies, the amino acid sequences of all the chains of a particular antibody are homologous with the chains found in one anti-
25 body produced by the lymphocyte which produces that antibody in situ, or in vitro (for example, in hybridomas). Vertebrate antibodies typically include native antibodies, for example, purified polyclonal antibodies and monoclonal antibodies. Examples of the methods for the
30 preparation of these antibodies are described infra.

"Hybrid antibodies" are antibodies wherein one pair of heavy and light chains is homologous to those in a first antibody, while the other pair of heavy and light chains is homologous to those in a different second antibody. Typically, each of these two pairs will bind different epitopes, particularly on different antigens. This
35 results in the property of "divalence", i.e., the ability

to bind two antigens simultaneously. Such hybrids may also be formed using chimeric chains, as set forth below.

5 "Chimeric antibodies", are antibodies in which the heavy and/or light chains are fusion proteins. Typically the constant domain of the chains is from one particular species and/or class, and the variable domains are from a different species and/or class. Also included is any antibody in which either or both of the heavy or
10 light chains are composed of combinations of sequences mimicking the sequences in antibodies of different sources, whether these sources be differing classes, or different species of origin, and whether or not the fusion point is at the variable/constant boundary. Thus, it is
15 possible to produce antibodies in which neither the constant nor the variable region mimic known antibody sequences. It then becomes possible, for example, to construct antibodies whose variable region has a higher specific affinity for a particular antigen, or whose
20 constant region can elicit enhanced complement fixation, or to make other improvements in properties possessed by a particular constant region.

Another example is "altered antibodies", which refers to antibodies in which the naturally occurring
25 amino acid sequence in a vertebrate antibody has been varied. Utilizing recombinant DNA techniques, antibodies can be redesigned to obtain desired characteristics. The possible variations are many, and range from the changing of one or more amino acids to the complete redesign of a
30 region, for example, the constant region. Changes in the constant region, in general, to attain desired cellular process characteristics, e.g., changes in complement fixation, interaction with membranes, and other effector functions. Changes in the variable region may be made to
35 alter antigen binding characteristics. The antibody may also be engineered to aid the specific delivery of a

molecule or substance to a specific cell or tissue site. The desired alterations may be made by known techniques in molecular biology, e.g., recombinant techniques, site
5 directed mutagenesis, etc.

Yet another example are "univalent antibodies", which are aggregates comprised of a heavy chain/light chain dimer bound to the Fc (i.e., constant) region of a second heavy chain. This type of antibody escapes
10 antigenic modulation. See, e.g., Glennie et al. (1982).

Included also within the definition of antibodies are "Fab" fragments of antibodies. The "Fab" region refers to those portions of the heavy and light chains which are roughly equivalent, or analogous, to the
15 sequences which comprise the branch portion of the heavy and light chains, and which have been shown to exhibit immunological binding to a specified antigen, but which lack the effector Fc portion. "Fab" includes aggregates
20 of one heavy and one light chain (commonly known as Fab'), as well as tetramers containing the 2H and 2L chains (referred to as F(ab)₂), which are capable of selectively reacting with a designated antigen or antigen family. "Fab" antibodies may be divided into subsets analogous to
25 those described above, i.e., "vertebrate Fab", "hybrid Fab", "chimeric Fab", and "altered Fab". Methods of producing "Fab" fragments of antibodies are known within the art and include, for example, proteolysis, and synthesis by recombinant techniques.

30 II.H. Diagnostic Oligonucleotide Probes and Kits

Using the disclosed portions of the isolated HCV cDNAs as a basis, oligomers of approximately 8 nucleotides or more can be prepared, either by excision or
35 synthetically, which hybridize with the HCV genome and are useful in identification of the viral agent(s), further characterization of the viral genome(s), as well as in

5 detection of the virus(es) in diseased individuals. The probes for HCV polynucleotides (natural or derived) are a length which allows the detection of unique viral sequences by hybridization. While 6-8 nucleotides may be a workable length, sequences of 10-12 nucleotides are preferred, and about 20 nucleotides appears optimal. Preferably, these sequences will derive from regions which lack heterogeneity. These probes can be prepared using routine methods, including automated oligonucleotide synthetic methods. Among useful probes, for example, are those derived from the newly isolated clones disclosed herein, as well as the various oligomers useful in probing cDNA libraries, set forth below. A complement to any unique portion of the HCV genome will be satisfactory. For use as probes, complete complementarity is desirable, though it may be unnecessary as the length of the fragment is increased.

20 For use of such probes as diagnostics, the biological sample to be analyzed, such as blood or serum, may be treated, if desired, to extract the nucleic acids contained therein. The resulting nucleic acid from the sample may be subjected to gel electrophoresis or other size separation techniques; alternatively, the nucleic acid sample may be dot blotted without size separation. The probes are then labeled. Suitable labels, and methods for labeling probes are known in the art, and include, for example, radioactive labels incorporated by nick translation or kinasing, biotin, fluorescent probes, and chemiluminescent probes. The nucleic acids extracted from the sample are then treated with the labeled probe under hybridization conditions of suitable stringencies, and polynucleotide duplexes containing the probe are detected.

35 The probes can be made completely complementary to the HCV genome. Therefore, usually high stringency conditions are desirable in order to prevent false

positives. However, conditions of high stringency should only be used if the probes are complementary to regions of the viral genome which lack heterogeneity. The stringency of hybridization is determined by a number of factors during hybridization and during the washing procedure, including temperature, ionic strength, length of time, and concentration of formamide. These factors are outlined in, for example, Maniatis, T. (1982).

Generally, it is expected that the HCV genome sequences will be present in serum of infected individuals at relatively low levels, i.e., at approximately 10^2 - 10^3 chimp infectious doses (CID) per ml. This level may require that amplification techniques be used in hybridization assays. Such techniques are known in the art. For example, the Enzo Biochemical Corporation "Bio-Bridge" system uses terminal deoxynucleotide transferase to add unmodified 3'-poly-dT-tails to a DNA probe. The poly dT-tailed probe is hybridized to the target nucleotide sequence, and then to a biotin-modified poly-A. PCT application 84/03520 and EPA124221 describe a DNA hybridization assay in which: (1) analyte is annealed to a single-stranded DNA probe that is complementary to an enzyme-labeled oligonucleotide; and (2) the resulting tailed duplex is hybridized to an enzyme-labeled oligonucleotide. EPA 204510 describes a DNA hybridization assay in which analyte DNA is contacted with a probe that has a tail, such as a poly-dT tail, an amplifier strand that has a sequence that hybridizes to the tail of the probe, such as a poly-A sequence, and which is capable of binding a plurality of labeled strands. A particularly desirable technique may first involve amplification of the target HCV sequences in sera approximately 10,000 fold, i.e., to approximately 10^6 sequences/ml. This may be accomplished, for example, by the polymerase chain reactions (PCR) technique described which is by Saiki et al. (1986),

by Mullis, U.S. Patent No. 4,683,195, and by Mullis et al. U.S. Patent No. 4,683,202. The amplified sequence(s) may then be detected using a hybridization assay which is
5 described in EP 317,077, published May 24, 1989. These hybridization assays, which should detect sequences at the level of 10^6 /ml, utilize nucleic acid multimers which bind to single-stranded analyte nucleic acid, and which also
10 bind to a multiplicity of single-stranded labeled oligonucleotides. A suitable solution phase sandwich assay which may be used with labeled polynucleotide probes, and the methods for the preparation of probes is described in EPO 225,807, published June 16, 1987.

15 The probes can be packaged into diagnostic kits. Diagnostic kits include the probe DNA, which may be labeled; alternatively, the probe DNA may be unlabeled and the ingredients for labeling may be included in the kit in separate containers. The kit may also contain other suitably packaged reagents and materials needed for the
20 particular hybridization protocol, for example, standards, as well as instructions for conducting the test.

Immunoassay and Diagnostic Kits

25 Both the polypeptides which react immunologically with serum containing HCV antibodies, for example, those detected by the antigenic screening method described infra. in the Examples, as well those derived from or encoded within the isolated clones described in
30 the Examples, and composites thereof, and the antibodies raised against the HCV specific epitopes in these polypeptides, are useful in immunoassays to detect presence of HCV antibodies, or the presence of the virus and/or viral antigens, in biological samples. Design of the
35 immunoassays is subject to a great deal of variation, and a variety of these are known in the art. For example, the immunoassay may utilize one viral epitope; alternatively,

the immunoassay may use a combination of viral epitopes derived from these sources; these epitopes may be derived from the same or from different viral polypeptides, and may be in separate recombinant or natural polypeptides, or together in the same recombinant polypeptides. It may use, for example, a monoclonal antibody directed towards a viral epitope(s), a combination of monoclonal antibodies directed towards epitopes of one viral antigen, monoclonal antibodies directed towards epitopes of different viral antigens, polyclonal antibodies directed towards the same viral antigen, or polyclonal antibodies directed towards different viral antigens. Protocols may be based, for example, upon competition, or direct reaction, or sandwich type assays. Protocols may also, for example, use solid supports, or may be by immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the probe are also known; examples of which are assays which utilize biotin and avidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

Some of the antigenic regions of the putative polyprotein have been mapped and identified by screening the antigenicity of bacterial expression products of HCV cDNAs which encode portions of the polyprotein. See the Examples. Other antigenic regions of HCV may be detected by expressing the portions of the HCV cDNAs in other expression systems, including yeast systems and cellular systems derived from insects and vertebrates. In addition, studies giving rise to an antigenicity index and hydrophobicity/hydrophilicity profile give rise to information concerning the probability of a region's antigenicity.

The studies on antigenic mapping by expression of HCV cDNAs showed that a number of clones containing

these cDNAs expressed polypeptides which were immunologically reactive with serum from individuals with NANBH. No single polypeptide was immunologically reactive with all sera. Five of these polypeptides were very immunogenic in that antibodies to the HCV epitopes in these polypeptides were detected in many different patient sera, although the overlap in detection was not complete. Thus, the results on the immunogenicity of the polypeptides encoded in the various clones suggest that efficient detection systems may include the use of panels of epitopes. The epitopes in the panel may be constructed into one or multiple polypeptides.

Kits suitable for immunodiagnosis and containing the appropriate labeled reagents are constructed by packaging the appropriate materials, including the polypeptides of the invention containing HCV epitopes or antibodies directed against HCV epitopes in suitable containers, along with the remaining reagents and materials required for the conduct of the assay, as well as a suitable set of assay instructions.

Further Characterization of the HCV Genome, Virions, and Viral Antigens Using Probes Derived From cDNA to the Viral Genome

The HCV cDNA sequence information in the newly isolated clones described in the Examples may be used to gain further information on the sequence of the HCV genome, and for identification and isolation of the HCV agent, and thus will aid in its characterization including the nature of the genome, the structure of the viral particle, and the nature of the antigens of which it is composed. This information, in turn, can lead to additional polynucleotide probes, polypeptides derived from the HCV genome, and antibodies directed against HCV

epitopes which would be useful for the diagnosis and/or treatment of HCV caused NANBH.

5 The cDNA sequence information in the above-mentioned clones is useful for the design of probes for the isolation of additional cDNA sequences which are derived from as yet undefined regions of the HCV genome(s) from which the cDNAs in clones described herein and in EP 0,316,218 are derived. For example, labeled probes
10 containing a sequence of approximately 8 or more nucleotides, and preferably 20 or more nucleotides, which are derived from regions close to the 5'-termini or 3'-termini of the composite HCV cDNA sequence shown in Fig. 17 may be used to isolate overlapping cDNA sequences from HCV cDNA
15 libraries. Alternatively, characterization of the genomic segments could be from the viral genome(s) isolated from purified HCV particles. Methods for purifying HCV particles and for detecting them during the purification procedure are described herein, *infra*. Procedures for
20 isolating polynucleotide genomes from viral particles are known in the art, and one procedure which may be used is that described in EP 0,218,316. The isolated genomic segments could then be cloned and sequenced. An example of
25 this technique, which utilizes amplification of the sequences to be cloned, is provided *infra*., and yielded clone 16jh.

 Methods for constructing cDNA libraries are known in the art, and are discussed *supra* and *infra*; a
30 method for the construction of HCV cDNA libraries in lambda-gt11 is discussed in EPO Pub. No. 318,216. However, cDNA libraries which are useful for screening with nucleic acid probes may also be constructed in other vectors known in the art, for example, lambda-gt10 (Huynh
35 et al. (1985)).

Screening for Anti-Viral Agents for HCV

5 The availability of cell culture and animal
model systems for HCV makes it possible to screen for
anti-viral agents which inhibit HCV replication, and
particularly for those agents which preferentially allow
cell growth and multiplication while inhibiting viral
replication. These screening methods are known by those
of skill in the art. Generally, the anti-viral agents are
10 tested at a variety of concentrations, for their effect on
preventing viral replication in cell culture systems which
support viral replication, and then for an inhibition of
infectivity or of viral pathogenicity (and a low level of
toxicity) in an animal model system.

15 The methods and compositions provided herein for
detecting HCV antigens and HCV polynucleotides are useful
for screening of anti-viral agents in that they provide an
alternative, and perhaps more sensitive means, for detect-
ing the agent's effect on viral replication than the cell
20 plaque assay or ID₅₀ assay. For example, the HCV-
polynucleotide probes described herein may be used to
quantitate the amount of viral nucleic acid produced in a
cell culture. This could be accomplished, for example, by
hybridization or competition hybridization of the infected
25 cell nucleic acids with a labeled HCV-polynucleotide
probe. For example, also, anti-HCV antibodies may be used
to identify and quantitate HCV antigen(s) in the cell
culture utilizing the immunoassays described herein. In
addition, since it may be desirable to quantitate HCV
30 antigens in the infected cell culture by a competition
assay, the polypeptides encoded within the HCV cDNAs
described herein are useful in these competition assays.
Generally, a recombinant HCV polypeptide derived from the
HCV cDNA would be labeled, and the inhibition of binding
35 of this labeled polypeptide to an HCV polypeptide due to
the antigen produced in the cell culture system would be

monitored. Moreover, these techniques are particularly useful in cases where the HCV may be able to replicate in a cell line without causing cell death.

5 The anti-viral agents which may be tested for efficacy by these methods are known in the art, and include, for example, those which interact with virion components and/or cellular components which are necessary for the binding and/or replication of the virus. Typical
10 anti-viral agents may include, for example, inhibitors of virion polymerase and/or protease(s) necessary for cleavage of the precursor polypeptides. Other anti-viral agents may include those which act with nucleic acids to prevent viral replication, for example, anti-sense
15 polynucleotides, etc.

Antisense polynucleotides molecules are comprised of a complementary nucleotide sequence which allows them to hybridize specifically to designated regions of genomes or RNAs. Antisense polynucleotides may
20 include, for example, molecules that will block protein translation by binding to mRNA, or may be molecules which prevent replication of viral RNA by transcriptase. They may also include molecules which carry agents (non-covalently attached or covalently bound) which cause the
25 viral RNA to be inactive by causing, for example, scissions in the viral RNA. They may also bind to cellular polynucleotides which enhance and/or are required for viral infectivity, replicative ability, or chronicity. Antisense molecules which are to hybridize to HCV derived
30 RNAs may be designed based upon the sequence information of the HCV cDNAs provided herein. The antiviral agents based upon anti-sense polynucleotides for HCV may be designed to bind with high specificity, to be of increased solubility, to be stable, and to have low toxicity.
35 Hence, they may be delivered in specialized systems, for example, liposomes, or by gene therapy. In addition, they

may include analogs, attached proteins, substituted or altered bonding between bases, etc.

5 Other types of drugs may be based upon polynucleotides which "mimic" important control regions of the HCV genome, and which may be therapeutic due to their interactions with key components of the system responsible for viral infectivity or replication.

10 General Methods

The general techniques used in extracting the genome from a virus, preparing and probing a cDNA library, sequencing clones, constructing expression vectors, trans-
15 forming cells, performing immunological assays such as radioimmunoassays and ELISA assays, for growing cells in culture, and the like are known in the art and laboratory manuals are available describing these techniques. However, as a general guide, the following sets forth some sources currently available for such procedures, and for
20 materials useful in carrying them out.

Both prokaryotic and eukaryotic host cells may be used for expression of desired coding sequences when appropriate control sequences which are compatible with the designated host are used. Among prokaryotic hosts, E. coli
25 coli is most frequently used. Expression control sequences for prokaryotes include promoters, optionally containing operator portions, and ribosome binding sites. Transfer vectors compatible with prokaryotic hosts are commonly derived from, for example, pBR322, a plasmid
30 containing operons conferring ampicillin and tetracycline resistance, and the various pUC vectors, which also contain sequences conferring antibiotic resistance markers. These markers may be used to obtain successful transformants by selection. Commonly used prokaryotic
35 control sequences include the Beta-lactamase (penicillinase) and lactose promoter systems (Chang et al.

(1977)), the tryptophan (*trp*) promoter system (Goeddel et al. (1980)) and the lambda-derived P_L promoter and N gene ribosome binding site (Shimatake et al. (1981)) and the hybrid *tac* promoter (De Boer et al. (1983)) derived from sequences of the *trp* and *lac* UV5 promoters. The foregoing systems are particularly compatible with *E. coli*; if desired, other prokaryotic hosts such as strains of *Bacillus* or *Pseudomonas* may be used, with corresponding control sequences.

Eukaryotic hosts include yeast and mammalian cells in culture systems. *Saccharomyces cerevisiae* and *Saccharomyces carlsbergensis* are the most commonly used yeast hosts, and are convenient fungal hosts. Yeast compatible vectors carry markers which permit selection of successful transformants by conferring prototrophy to auxotrophic mutants or resistance to heavy metals on wild-type strains. Yeast compatible vectors may employ the 2 micron origin of replication (Broach et al. (1983)), the combination of CEN3 and ARS1 or other means for assuring replication, such as sequences which will result in incorporation of an appropriate fragment into the host cell genome. Control sequences for yeast vectors are known in the art and include promoters for the synthesis of glycolytic enzymes (Hess et al. (1968); Holland et al. (1978)), including the promoter for 3 phosphoglycerate kinase (Hitzeman (1980)). Terminators may also be included, such as those derived from the enolase gene (Holland (1981)). Particularly useful control systems are those which comprise the glyceraldehyde-3 phosphate dehydrogenase (GAPDH) promoter or alcohol dehydrogenase (ADH) regulatable promoter, terminators also derived from GAPDH, and if secretion is desired, leader sequence from yeast alpha factor. In addition, the transcriptional regulatory region and the transcriptional initiation region which are operably linked may be such that they are

not naturally associated in the wild-type organism. These systems are described in detail in EPO 120,551, published October 3, 1984; EPO 116,201, published August 22, 1984; and EPO 164,556, published December 18, 1985, all of which are assigned to the herein assignee, and are hereby incorporated herein by reference.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), including HeLa cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells, and a number of other cell lines. Suitable promoters for mammalian cells are also known in the art and include viral promoters such as that from Simian Virus 40 (SV40) (Fiers (1978)), Rous sarcoma virus (RSV), adenovirus (ADV), and bovine papilloma virus (BPV). Mammalian cells may also require terminator sequences and poly A addition sequences; enhancer sequences which increase expression may also be included, and sequences which cause amplification of the gene may also be desirable. These sequences are known in the art. Vectors suitable for replication in mammalian cells may include viral replicons, or sequences which insure integration of the appropriate sequences encoding NANBV epitopes into the host genome.

Transformation may be by any known method for introducing polynucleotides into a host cell, including, for example packaging the polynucleotide in a virus and transducing a host cell with the virus, and by direct uptake of the polynucleotide. The transformation procedure used depends upon the host to be transformed. For example, transformation of the E. coli host cells with lambda-gt11 containing BB-NANBV sequences is discussed in the Example section, infra. Bacterial transformation by direct uptake generally employs treatment with calcium or rubidium chloride (Cohen (1972); Maniatis (1982)). Yeast

transformation by direct uptake may be carried out using the method of Hinnen et al. (1978). Mammalian transformations by direct uptake may be conducted using the calcium phosphate precipitation method of Graham and Van der Eb (1978), or the various known modifications thereof.

Vector construction employs techniques which are known in the art. Site-specific DNA cleavage is performed by treating with suitable restriction enzymes under conditions which generally are specified by the manufacturer of these commercially available enzymes. In general, about 1 microgram of plasmid or DNA sequence is cleaved by 1 unit of enzyme in about 20 microliters buffer solution by incubation of 1-2 hr at 37° C. After incubation with the restriction enzyme, protein is removed by phenol/chloroform extraction and the DNA recovered by precipitation with ethanol. The cleaved fragments may be separated using polyacrylamide or agarose gel electrophoresis techniques, according to the general procedures found in Methods in Enzymology (1980) 65:499-560.

Sticky ended cleavage fragments may be blunt ended using *E. coli* DNA polymerase I (Klenow) in the presence of the appropriate deoxynucleotide triphosphates (dNTPs) present in the mixture. Treatment with S1 nuclease may also be used, resulting in the hydrolysis of any single stranded DNA portions.

Ligations are carried out using standard buffer and temperature conditions using T4 DNA ligase and ATP; sticky end ligations require less ATP and less ligase than blunt end ligations. When vector fragments are used as part of a ligation mixture, the vector fragment is often treated with bacterial alkaline phosphatase (BAP) or calf intestinal alkaline phosphatase to remove the 5'-phosphate and thus prevent religation of the vector; alternatively, restriction enzyme digestion of unwanted fragments can be used to prevent ligation.

Ligation mixtures are transformed into suitable cloning hosts, such as E. coli, and successful transformants selected by, for example, antibiotic resistance, and screened for the correct construction.

Synthetic oligonucleotides may be prepared using an automated oligonucleotide synthesizer as described by Warner (1984). If desired the synthetic strands may be labeled with ^{32}P by treatment with polynucleotide kinase in the presence of ^{32}P -ATP, using standard conditions for the reaction.

DNA sequences, including those isolated from cDNA libraries, may be modified by known techniques, including, for example site directed mutagenesis, as described by Zoller (1982). Briefly, the DNA to be modified is packaged into phage as a single stranded sequence, and converted to a double stranded DNA with DNA polymerase using, as a primer, a synthetic oligonucleotide complementary to the portion of the DNA to be modified, and having the desired modification included in its own sequence. The resulting double stranded DNA is transformed into a phage supporting host bacterium. Cultures of the transformed bacteria, which contain replications of each strand of the phage, are plated in agar to obtain plaques. Theoretically, 50% of the new plaques contain phage having the mutated sequence, and the remaining 50% have the original sequence. Replicates of the plaques are hybridized to labeled synthetic probe at temperatures and conditions which permit hybridization with the correct strand, but not with the unmodified sequence. The sequences which have been identified by hybridization are recovered and cloned.

DNA libraries may be probed using the procedure of Grunstein and Hogness (1975). Briefly, in this procedure, the DNA to be probed is immobilized on nitro-cellulose filters, denatured, and prehybridized with a

buffer containing 0-50% formamide, 0.75 M NaCl, 75 mM Na citrate, 0.02% (wt/v) each of bovine serum albumin, polyvinyl pyrrolidone, and Ficoll, 50 mM Na Phosphate (pH 6.5), 0.1% SDS, and 100 micrograms/ml carrier denatured DNA. The percentage of formamide in the buffer, as well as the time and temperature conditions of the prehybridization and subsequent hybridization steps depends on the stringency required. Oligomeric probes which require lower stringency conditions are generally used with low percentages of formamide, lower temperatures, and longer hybridization times. Probes containing more than 30 or 40 nucleotides such as those derived from cDNA or genomic sequences generally employ higher temperatures, e.g., about 40-42°C, and a high percentage, e.g., 50%, formamide. Following prehybridization, 5'-³²P-labeled oligonucleotide probe is added to the buffer, and the filters are incubated in this mixture under hybridization conditions. After washing, the treated filters are subjected to autoradiography to show the location of the hybridized probe; DNA in corresponding locations on the original agar plates is used as the source of the desired DNA.

For routine vector constructions, ligation mixtures are transformed into *E. coli* strain HB101 or other suitable host, and successful transformants selected by antibiotic resistance or other markers. Plasmids from the transformants are then prepared according to the method of Clewell et al. (1969), usually following chloramphenicol amplification (Clewell (1972)). The DNA is isolated and analyzed, usually by restriction enzyme analysis and/or sequencing. Sequencing may be by the dideoxy method of Sanger et al. (1977) as further described by Messing et al. (1981), or by the method of Maxam et al. (1980). Problems with band compression, which are sometimes observed in GC rich regions, were

overcome by use of T-deazoguanosine according to Barr et al. (1986).

5 The enzyme-linked immunosorbent assay (ELISA)
can be used to measure either antigen or antibody
concentrations. This method depends upon conjugation of
an enzyme to either an antigen or an antibody, and uses
the bound enzyme activity as a quantitative label. To
10 measure antibody, the known antigen is fixed to a solid
phase (e.g., a microplate or plastic cup), incubated with
test serum dilutions, washed, incubated with anti-
immunoglobulin labeled with an enzyme, and washed again.
Enzymes suitable for labeling are known in the art, and
15 include, for example, horseradish peroxidase. Enzyme
activity bound to the solid phase is measured by adding
the specific substrate, and determining product formation
or substrate utilization colorimetrically. The enzyme
activity bound is a direct function of the amount of anti-
body bound.

20 To measure antigen, a known specific antibody is
fixed to the solid phase, the test material containing
antigen is added, after an incubation the solid phase is
washed, and a second enzyme-labeled antibody is added.
25 After washing, substrate is added, and enzyme activity is
estimated colorimetrically, and related to antigen
concentration.

Examples

30 Described below are examples of the present
invention which are provided only for illustrative
purposes, and not to limit the scope of the present inven-
tion. In light of the present disclosure, numerous
embodiments within the scope of the claims will be appar-
35 ent to those of ordinary skill in the art.

Isolation and Sequence of Overlapping

HCV cDNA Clones 13i, 26j, CA59a, CA84a, CA156e and CA167b

5 The clones 13i, 26j, CA59a, CA84a, CA156e and CA167b were isolated from the lambda-gt11 library which contains HCV cDNA (ATCC No. 40394), the preparation of which is described in EPO Pub. No. 318,216 (published 31 May 1989), and WO 89/04669 (published 1 June 1989).

10 Screening of the library was with the probes described infra., using the method described in Huynh (1985). The frequencies with which positive clones appeared with the respective probes was about 1 in 50,000.

15 The isolation of clone 13i was accomplished using a synthetic probe derived from the sequence of clone 12f. The sequence of the probe was:

5' GAA CGT TGC GAT CTG GAA GAC AGG GAC AGG 3'.

20 The isolation of clone 26j was accomplished using a probe derived from the 5'-region of clone K9-1. The sequence of the probe was:

5' TAT CAG TTA TGC CAA CGG AAG CGG CCC CGA 3'.

25 The isolation procedures for clone 12f and for clone k9-1 (also called K9-1) are described in EPO Pub. No. 318,216, and their sequences are shown in Figs. 1 and 2, respectively. The HCV cDNA sequences of clones 13i and 26j, are shown in Figs. 4 and 5, respectively. Also shown
30 are the amino acids encoded therein, as well as the overlap of clone 13i with clone 12f, and the overlap of clone 26j with clone 13i. The sequences for these clones confirmed the sequence of clone K9-1. Clone K9-1 had been
35 isolated from a different HCV cDNA library (See EP 0,218,316).

Clone CA59a was isolated utilizing a probe based upon the sequence of the 5'-region of clone 26j. The sequence of this probe was:

5' CTG GTT AGC AGG GCT TTT CTA TCA CCA CAA 3'.

A probe derived from the sequence of clone CA59a was used to isolate clone CA84a. The sequence of the probe used for this isolation was:

5' AAG GTC CTG GTA GTG CTG CTG CTA TTT GCC 3'.

Clone CA156e was isolated using a probe derived from the sequence of clone CA84a. The sequence of the probe was:

5' ACT GGA CGA CGC AAG GTT GCA ATT GCT CTA 3'.

Clone CA167b was isolated using a probe derived from the sequence of clone CA 156e. The sequence of the probe was:

5' TTC GAC GTC ACA TCG ATC TGC TTG TCG GGA 3'.

The nucleotide sequences of the HCV cDNAs in clones CA59a, CA84a, CA156e, and CA167b, are shown Figs. 6, 7, 8, and 9, respectively. The amino acids encoded therein, as well as the overlap with the sequences of relevant clones, are also shown in the Figs.

Creation of "pi" HCV cDNA Library

A library of HCV cDNA, the "pi" library, was constructed from the same batch of infectious chimpanzee plasma used to construct the lambda-gt11 HCV cDNA library (ATCC No. 40394) described in EPO Pub. No. 318,216, and

utilizing essentially the same techniques. However, construction of the pi library utilized a primer-extension method, in which the primer for reverse transcriptase was based on the sequence of clone CA59A. The sequence of the primer was:

5' GGT GAC GTG GGT TTC 3'.

10

Isolation and Sequence of Clone pi14a

Screening of the "pi" HCV cDNA library described supra., with the probe used to isolate clone CA167b (See supra.) yielded clone pi14a. The clone contains about 800 base pairs of cDNA which overlaps clones CA167b, CA156e, CA84a and CA59a, which were isolated from the lambda gt-11 HCV cDNA library (ATCC No. 40394). In addition, pi14a also contains about 250 base pairs of DNA which are upstream of the HCV cDNA in clone CA167b.

20

Isolation and Sequence of Clones CA216a, CA290a and aq30a

Based on the sequence of clone CA167b a synthetic probe was made having the following sequence:

25

5' GGC TTT ACC ACG TCA CCA ATG ATT GCC CTA 3'

The above probe was used to screen the , which yielded clone CA216a, whose HCV sequences are shown in Fig. 10.

Another probe was made based on the sequence of clone CA216a having the following sequence:

30

5' TTT GGG TAA GGT CAT CGA TAC CCT TAC GTG 3'

Screening the lambda-gt11 library (ATCC No. 40394) with this probe yielded clone CA290a, the HCV sequences therein being shown in Fig. 11.

35

In a parallel approach, a primer-extension cDNA library was made using nucleic acid extracted from the same infectious plasma used in the original lambda-gt11 cDNA library described above. The primer used was based on the sequence of clones CA216a and CA290a:

5' GAA GCC GCA CGT AAG 3'

The cDNA library was made using methods similar to those described previously for libraries used in the isolation of clones pil4a and k9-1. The probe used to screen this library was based on the sequence of clone CA290a:

5' CCG GCG TAG GTC GCG CAA TTT GGG TAA 3'

Clone ag30a was isolated from the new library with the above probe, and contained about 670 basepairs of HCV sequence. See Fig. 12. Part of this sequence overlaps the HCV sequence of clones CA216a and CA290a. About 300 base-pairs of the ag30a sequence, however, is upstream of the sequence from clone CA290a. The non-overlapping sequence shows a start codon (*) and stop codons that may indicate the start of the HCV ORF. Also indicated in Fig. 12 are putative small encoded peptides (#) which may play a role in regulating translation, as well as the putative first amino acid of the putative polypeptide (/), and downstream amino acids encoded therein.

Isolation and Sequence of Clone CA205a

Clone CA205a was isolated from the original lambda gt-11 library (ATCC No. 40394), using a synthetic probe derived from the HCV sequence in clone CA290a (Fig. 11). The sequence of the probe was:

5' TCA GAT CGT TGG TGG AGT TTA CTT GTT GCC 3'.

5 The sequence of the HCV cDNA in CA205a, shown in Fig. 13, overlaps with the cDNA sequences in both clones ag30a and CA290a. The overlap of the sequence with that of CA290a is shown by the dotted line above the sequence (the figure also shows the putative amino acids encoded in this fragment).

10 As observed from the HCV cDNA sequences in clones CA205a and ag30a, the putative HCV polyprotein appears to begin at the ATG start codon; the HCV sequences in both clones contain an in-frame, contiguous double stop codon (TGATAG) forty two nucleotides upstream from this ATG. The HCV ORF appears to begin after these stop
15 codons, and to extend for at least 8907 nucleotides (See the composite HCV cDNA shown in Fig. 17).

Isolation and Sequence of Clone 18g

20 Based on the sequence of clone ag30a (See Fig. 12) and of an overlapping clone from the original lambda gt-11 library (ATCC No. 40394), CA230a, a synthetic probe was made having the following sequence:

5' CCA TAG TGG TCT GCG GAA CCG GTG AGT ACA 3'.

25 Screening of the original lambda-gt11 HCV cDNA library with the probe yielded clone 18g, the HCV cDNA sequence of which is shown in Fig. 14. Also shown in the figure are the overlap with clone ag30a, and putative polypeptides encoded within the HCV cDNA.

30 The cDNA in clone 18g (C18g or 18g) overlaps that in clones ag30a and CA205a, described supra. The sequence of C18g also contains the double stop codon region observed in clone ag30a. The polynucleotide region upstream of these stop codons presumably represents part
35 of the 5'-region of the HCV genome, which may contain short ORFs, and which can be confirmed by direct sequenc-

ing of the purified HCV genome. These putative small encoded peptides may play a regulatory role in translation. The region of the HCV genome upstream of that
5 represented by C18g can be isolated for sequence analysis using essentially the technique described in EPO Pub. No. 318,216 for isolating cDNA sequences upstream of the HCV cDNA sequence in clone 12f. Essentially, small synthetic oligonucleotide primers of reverse transcriptase, which
10 are based upon the sequence of C18g, are synthesized and used to bind to the corresponding sequence in HCV genomic RNA. The primer sequences are proximal to the known 5'-terminal of C18g, but sufficiently downstream to allow the design of probe sequences upstream of the primer
15 sequences. Known standard methods of priming and cloning are used. The resulting cDNA libraries are screened with sequences upstream of the priming sites (as deduced from the elucidated sequence of C18g). The HCV genomic RNA is obtained from either plasma or liver samples from
20 individuals with NANBH. Since HCV appears to be a Flavivirus-like virus, the 5'-terminus of the genome may be modified with a "cap" structure. It is known that Flavivirus genomes contain 5'-terminal "cap" structures. (Yellow
25 Fever virus, Rice et al. (1988); Dengue virus, Hahn et al (1988); Japanese Encephalitis Virus (1987)).

Isolation and Sequence of Clones from
the beta-HCV cDNA library

30 Clones containing cDNA representative of the 3'-terminal region of the HCV genome were isolated from a cDNA library constructed from the original infectious chimpanzee plasma pool which was used for the creation of the HCV cDNA lambda-gt11 library (ATCC No. 40394),
35 described in EPO Pub. No. 318,216. In order to create the DNA library, RNA extracted from the plasma was "tailed" with poly rA using poly (rA) polymerase, and cDNA was

synthesized using oligo(dT)₁₂₋₁₈ as a primer for reverse transcriptase. The resulting RNA:cDNA hybrid was digested with RNAase H, and converted to double stranded HCV cDNA. The resulting HCV cDNA was cloned into lambda-gt10, using essentially the technique described in Huynh (1985), yielding the beta (or b) HCV cDNA library. The procedures used were as follows.

An aliquot (12ml) of the plasma was treated with proteinase K, and extracted with an equal volume of phenol saturated with 0.05M Tris-Cl, pH 7.5, 0.05% (v/v) beta-mercaptoethanol, 0.1% (w/v) hydroxyquinolone, 1 mM EDTA. The resulting aqueous phase was re-extracted with the phenol mixture, followed by 3 extractions with a 1:1 mixture containing phenol and chloroform:isoamyl alcohol (24:1), followed by 2 extractions with a mixture of chloroform and isoamyl alcohol (1:1). Subsequent to adjustment of the aqueous phase to 200 mM with respect to NaCl, nucleic acids in the aqueous phase were precipitated overnight at -20°C, with 2.5 volumes of cold absolute ethanol. The precipitates were collected by centrifugation at 10,000 RPM for 40 min., washed with 70% ethanol containing 20 mM NaCl, and with 100% cold ethanol, dried for 5 min. in a dessicator, and dissolved in water.

The isolated nucleic acids from the infectious chimpanzee plasma pool were tailed with poly rA utilizing poly-A polymerase in the presence of human placenta ribonuclease inhibitor (HPRI) (purchased from Amersham Corp.), utilizing MS2 RNA as carrier. Isolated nucleic acids equivalent to that in 2 ml of plasma were incubated in a solution containing TMN (50 mM Tris HCl, pH 7.9, 10 mM MgCl₂, 250 mM NaCl, 2.5 mM MnCl₂, 2 mM dithiothreitol (DTT)), 40 micromolar alpha-[³²P] ATP, 20 units HPRI (Amersham Corp.), and about 9 to 10 units of RNase free poly-A polymerase (BRL). Incubation was for 10 min. at 37°C, and the reactions were stopped with EDTA (final

concentration about 250 mM). The solution was extracted with an equal volume of phenol-chloroform, and with an equal volume of chloroform, and nucleic acids were precipitated overnight at -20°C with 2.5 volumes of ethanol in the presence of 200 mM NaCl.

Isolation of Clone b5a

The beta HCV cDNA library was screened by hybridization using a synthetic probe, which had a sequence based upon the HCV cDNA sequence in clone 15e. The isolation of clone 15e is described in EPO Pub. No. 318,216, and its sequence is shown in Fig. 3. The sequence of the synthetic probe was:

5' ATT GCG AGA TCT ACG GGG CCT GCT ACT CCA 3'.

Screening of the library yielded clone beta-5a (b5a), which contains an HCV cDNA region of approximately 1000 base pairs. The 5'-region of this cDNA overlaps clones 35f, 19g, 26g, and 15e (these clones are described supra). The region between the 3'-terminal poly-A sequence and the 3'-sequence which overlaps clone 15e, contains approximately 200 base pairs. This clone allows the identification of a region of the 3'-terminal sequence the HCV genome.

The sequence of b5a is contained within the sequence of the HCV cDNA in clone 16jh (described infra). Moreover, the sequence is also present in CC34a, isolated from the original lambda-gt11 library (ATCC No. 40394). (The original lambda-gt11 library is referred to herein as the "C" library).

Isolation and Sequence of Clones Generated by PCR
Amplification of the 3'-Region of the HCV Genome

5 Multiple cDNA clones have been generated which
 contain nucleotide sequences derived from the 3'-region of
 the HCV genome. This was accomplished by amplifying a
 targeted region of the genome by a polymerase chain re-
 action technique described in Saiki et al. (1986), and in
 10 Saiki et al. (1988), which was modified as described
 below. The HCV RNA which was amplified was obtained from
 the original infectious chimpanzee plasma pool which was
 used for the creation of the HCV cDNA lambda-gt11 library
 (ATCC No. 40394) described in EPO Pub. No. 318,216.
 15 Isolation of the HCV RNA was as described supra. The
 isolated RNA was tailed at the 3'-end with ATP by *E. coli*
 poly-A polymerase as described in Sippel (1973), except
 that the nucleic acids isolated from chimp serum were
 substituted for the nucleic acid substrate. The tailed
 20 RNA was then reverse transcribed into cDNA by reverse
 transcriptase, using an oligo dT-primer adapter, es-
 sentially as described by Han (1987), except that the
 components and sequence of the primer-adapter were:

25	<u>Stuffer</u>	<u>NotI</u>	<u>SP6 Promoter</u>	<u>Primer</u>
	AATTC	GCGGCCGC	CATACGATTTAGGTGACACTATAGAA	T ₁₅

The resultant cDNA was subjected to amplification by PCR
 using two primers:

30	<u>Primer</u>	<u>Sequence</u>
	JH32 (30mer)	ATAGCGGCCGCCCTCGATTGCGAGATCTAC
	JH11 (20mer)	AATTCGGGCGGCCGCCATACGA

35 The JH32 primer contained 20 nucleotide sequences
 hybridizable to the 5'-end of the target region in the
 cDNA, with an estimated T_m of 66°C. The JH11 was derived

from a portion of the oligo dT-primer adapter; thus, it is specific to the 3'-end of the cDNA with a T_m of 64°C.

Both primers were designed to have a recognition site for the restriction enzyme, NotI, at the 5'-end, for use in subsequent cloning of the amplified HCV cDNA.

The PCR reaction was carried out by suspending the cDNA and the primers in 100 microliters of reaction mixture containing the four deoxynucleoside triphosphates, buffer salts and metal ions, and a thermostable DNA polymerase isolated from Thermus aquaticus (Taq polymerase), which are in a Perkin Elmer Cetus PCR kit (N801-0043 or N801-0055). The PCR reaction was performed for 35 cycles in a Perkin Elmer Cetus DNA thermal cycler. Each cycle consisted of a 1.5 min denaturation step at 94°C, an annealing step at 60°C for 2 min, and a primer extension step at 72°C for 3 min. The PCR products were subjected to Southern blot analysis using a 30 nucleotide probe, JH34, the sequence of which was based upon that of the 3'-terminal region of clone 15e. The sequence of JH34 is:

5' CTT GAT CTA CCT CCA ATC ATT CAA AGA CTC 3'.

The PCR products detected by the HCV cDNA probe ranged in size from about 50 to about 400 base pairs.

In order to clone the amplified HCV cDNA, the PCR products were cleaved with NotI and size selected by polyacrylamide gel electrophoresis. DNA larger than 300 base pairs was cloned into the NotI site of pUC18S. The vector pUC18S is constructed by including a NotI polylinker cloned between the EcoRI and Sall sites of pUC18. The clones were screened for HCV cDNA using the JH34 probe. A number of positive clones were obtained and sequenced. The nucleotide sequence of the HCV cDNA insert in one of these clones, 16jh, and the amino acids encoded

therein, are shown in Fig. 15. A nucleotide heterogeneity, detected in the sequence of the HCV cDNA in clone 16jh as compared to another clone of this region, is indicated in the figure.

Compiled HCV cDNA Sequences

An HCV cDNA sequence has been compiled from a series of overlapping clones derived from the various HCV cDNA libraries described supra.. In this sequence, the compiled HCV cDNA sequence obtained from clones b114a, 18g, ag30a, CA205a, CA290a, CA216a, pi14a, CA167b, CA156e, CA84a, and CA59a is upstream of the compiled HCV cDNA sequence published in EPO Pub. No. 318,216, which is shown in Fig. 16. The compiled HCV cDNA sequence obtained from clones b5a and 16jh downstream of the compiled HCV cDNA sequence published in EPO Pub. No. 318,216.

Fig. 17 shows the compiled HCV cDNA sequence derived from the above-described clones and the compiled HCV cDNA sequence published in EPO Pub. No. 318,216. The clones from which the sequence was derived are b114a, 18g, ag30a, CA205a, CA290a, CA216a, pi14a, CA167b, CA156e, CA84a, CA59a, K9-1 (also called k9-1), 26j, 13i, 12f, 14i, 11b, 7f, 7e, 8h, 33c, 40b, 37b, 35, 36, 81, 32, 33b, 25c, 14c, 8f, 33f, 33g, 39c, 35f, 19g, 26g, 15e, b5a, and 16jh. In the figure the three dashes above the sequence indicate the position of the putative initiator methionine codon.

Clone b114a was obtained using the cloning procedure described for clone b5a, supra., except that the probe was the synthetic probe used to detect clone 18g, supra. Clone b114a overlaps with clones 18g, ag30a, and CA205a, except that clone b114a contains an extra two nucleotides upstream of the sequence in clone 18g (i.e., 5'-CA). These extra two nucleotides have been included in the HCV genomic sequence shown in Fig. 17.

It should be noted that although several of the clones described supra. have been obtained from libraries other than the original HCV cDNA lambda-gt11 C library (ATCC No. 40394), these clones contain HCV cDNA sequences which overlap HCV cDNA sequences in the original library. Thus, essentially all of the HCV sequence is derivable from the original lambda-gt11 C library (ATCC No. 40394) which was used to isolate the first HCV cDNA clone (5-1-1). The isolation of clone 5-1-1 is described in EPO Pub. No. 318,216.

Purification of Fusion Polypeptide C100-3
(Alternate method)

The fusion polypeptide, C100-3 (also called HCV c100-3 and alternatively, c100-3), is comprised of superoxide dismutase (SOD) at the N-terminus an in-frame C100 HCV polypeptide at the C-terminus. A method for preparing the polypeptide by expression in yeast, and differential extraction of the insoluble fraction of the extracted host yeast cells, is described in EPO Pub. No. 318,216. An alternative method for the preparation of this fusion polypeptide is described below. In this method the antigen is precipitated from the crude cell lysate with acetone; the acetone precipitated antigen is then subjected to ion-exchange chromatography, and further purified by gel filtration.

The fusion polypeptide, C100-3 (HCV c100-3), is expressed in yeast strain JSC 308 (ATCC No. 20879) transformed with pAB24C100-3 (ATCC No. 67976); the transformed yeast are grown under conditions which allow expression (i.e., by growth in YEP containing 1% glucose). (See EPO Pub. No. 318,216). A cell lysate is prepared by suspending the cells in Buffer A (20 mM Tris HCl, pH 8.0, 1 mM EDTA, 1 mM PMSF. The cells are broken by grinding with glass beads in a Dynomill type homogenizer or its

equivalent. The extent of cell breakage is monitored by counting cells under a microscope with phase optics.

5 Broken cells appear dark, while viable cells are light-colored. The percentage of broken cells is determined.

When the percentage of broken cells is approximately 90% or greater, the broken cell debris is separated from the glass beads by centrifugation, and the glass beads are washed with Buffer A. After combining the washes and homogenate, the insoluble material in the lysate is obtained by centrifugation. The material in the pellet is washed to remove soluble proteins by suspension in Buffer B (50 mM glycine, pH 12.0, 1 mM DTT, 500 mM NaCl), followed by Buffer C (50 mM glycine, pH 10.0, 1 mM DTT). The insoluble material is recovered by centrifugation, and solubilized by suspension in Buffer C containing SDS. The extract solution may be heated in the presence of beta-mercaptoethanol and concentrated by ultrafiltration. The HCV c100-3 in the extract is precipitated with cold acetone. If desired, the precipitate may be stored at temperatures at about or below -15°C.

Prior to ion exchange chromatography, the acetone precipitated material is recovered by centrifugation, and may be dried under nitrogen. The precipitate is suspended in Buffer D (50 mM glycine, pH 10.0, 1 mM DTT, 7 M urea), and centrifuged to pellet insoluble material. The supernatant material is applied to an anion exchange column previously equilibrated with Buffer D. Fractions are collected and analyzed by ultraviolet absorbance or gel electrophoresis on SDS polyacrylamide gels. Those fractions containing the HCV c100-3 polypeptide are pooled.

In order to purify the HCV c100-3 polypeptide by gel filtration, the pooled fractions from the ion-exchange column are heated in the presence of beta-mercaptoethanol

and SDS, and the eluate is concentrated by ultrafiltration. The concentrate is applied to a gel filtration column previously equilibrated with Buffer E (20 mM Tris HCl, pH 7.0, 1 mM DTT, 0.1% SDS). The presence of HCV c100-3 in the eluted fractions, as well as the presence of impurities, are determined by gel electrophoresis on polyacrylamide gels in the presence of SDS and visualization of the polypeptides. Those fractions containing purified HCV c100-3 are pooled. Fractions high in HCV c100-3 may be further purified by repeating the gel filtration process. If the removal of particulate material is desired, the HCV c100-3 containing material may be filtered through a 0.22 micron filter.

Expression and Antigenicity of Polypeptides
Encoded in HCV cDNA

Polypeptides Expressed in E. coli

The polypeptides encoded in a number of HCV cDNAs which span the HCV genomic ORF were expressed in E. coli, and tested for their antigenicity using serum obtained from a variety of individuals with NANBH. The expression vectors containing the cloned HCV cDNAs were constructed from pSODcfl (Steimer et al. (1986)). In order to be certain that a correct reading frame would be achieved, three separate expression vectors, pcflAB, pcflCD, and pcflEF were created by ligating either of three linkers, AB, CD, and EF to a BamHI-EcoRI fragment derived by digesting to completion the vector pSODcfl with EcoRI and BamHI, followed by treatment with alkaline phosphatase. The linkers were created from six oligomers, A, B, C, D, E, and F. Each oligomer was phosphorylated by treatment with kinase in the presence of ATP prior to annealing to its complementary oligomer. The sequences of the synthetic linkers were the following.

	<u>Name</u>	<u>DNA Sequence (5' to 3')</u>
5	A	GATC CTG AAT TCC TGA TAA
	B	GAC TTA AGG ACT ATT TTA A
	C	GATC CGA ATT CTG TGA TAA
10	D	GCT TAA GAC ACT ATT TTA A
	E	GATC CTG GAA TTC TGA TAA
	F	GAC CTT AAG ACT ATT TTA A

Each of the three linkers destroys the original EcoRI site, and creates a new EcoRI site within the linker, but within a different reading frame. Hence, the HCV cDNA EcoRI fragments isolated from the clones when inserted into the expression vector, were in three different reading frames.

The HCV cDNA fragments in the designated lambda-gt11 clones were excised by digestion with EcoRI; each fragment was inserted into pcflAB, pcflCD, and pcflEF. These expression constructs were then transformed into D1210 E. coli cells, the transformants were cloned, and recombinant bacteria from each clone were induced to express the fusion polypeptides by growing the bacteria in the presence of IPTG.

Expression products of the indicated HCV cDNAs were tested for antigenicity by direct immunological screening of the colonies, using a modification of the method described in Helfman et al. (1983). Briefly, as shown in Fig. 18, the bacteria were plated onto nitrocellulose filters overlaid on ampicillin plates to give approximately 1,000 colonies per filter. Colonies were replica plated onto nitrocellulose filters, and the replicas were regrown overnight in the presence of 2 mM

5 IPTG and ampicillin. The bacterial colonies were lysed by
suspending the nitrocellulose filters for about 15 to 20
min in an atmosphere saturated with CHCl_3 vapor. Each
10 filter then was placed in an individual 100 mm Petri dish
containing 10 ml of 50 mM Tris HCl, pH 7.5, 150 mM NaCl, 5
mM MgCl_2 , 3% (w/v) BSA, 40 micrograms/ml lysozyme, and 0.1
microgram/ml DNase. The plates were agitated gently for
at least 8 hours at room temperature. The filters were
15 rinsed in TBST (50 mM Tris HCl, pH 8.0, 150 mM NaCl, 0.005%
Tween 20). After incubation, the cell residues were
rinsed and incubated in TBS (TBST without Tween) contain-
ing 10% sheep serum; incubation was for 1 hour. The
filters were then incubated with pretreated sera in TBS
20 from individuals with NANBH, which included: 3
chimpanzees; 8 patients with chronic NANBH whose sera were
positive with respect to antibodies to HCV C100-3
polypeptide (described in EPO Pub. No. 318,216, and
supra.) (also called C100); 8 patients with chronic NANBH
25 whose sera were negative for anti-C100 antibodies; a
convalescent patient whose serum was negative for anti-
C100 antibodies; and 6 patients with community acquired
NANBH, including one whose sera was strongly positive with
respect to anti-C100 antibodies, and one whose sera was
30 marginally positive with respect to anti-C100 antibodies.
The sera, diluted in TBS, was pretreated by preabsorption
with hSOD. Incubation of the filters with the sera was
for at least two hours. After incubation, the filters
were washed two times for 30 min with TBST. Labeling of
35 expressed proteins to which antibodies in the sera bound
was accomplished by incubation for 2 hours with ^{125}I -
labeled sheep anti-human antibody. After washing, the
filters were washed twice for 30 min with TBST, dried, and
autoradiographed.

A number of clones (see infra.) expressed
polypeptides containing HCV epitopes which were im-

munologically reactive with serum from individuals with NANBH. Five of these polypeptides were very immunogenic in that antibodies to HCV epitopes in these polypeptides were detected in many different patient sera. The clones encoding these polypeptides, and the location of the polypeptide in the putative HCV polyprotein (wherein the amino acid numbers begin with the putative initiator codon) are the following: clone 5-1-1, amino acids 1694-1735; clone C100, amino acids 1569-1931; clone 33c, amino acids 1192-1457; clone CA279a, amino acids 1-84; and clone CA290a amino acids 9-177. The location of the immunogenic polypeptides within the putative HCV polyprotein are shown immediately below.

15

20

25

30

35

5 Clones encoding polypeptides of proven reactivity
 with sera from NANBH patients.

	<u>Clone</u>	<u>Location within the HCV polyprotein</u> (amino acid no. beginning with puta- tive initiator methionine)
10		
	CA279a	1-84
	CA74a	437-582
	13i	511-690
	CA290a	9-177
15	33c	1192-1457
	40b	1266-1428
	5-1-1	1694-1735
	81	1689-1805
	33b	1916-2021
20	25c	1949-2124
	14c	2054-2223
	8f	2200-3325
	33f	2287-2385
	33g	2348-2464
25	39c	2371-2502
	15e	2796-2886
	C100	1569-1931

30 The results on the immunogenicity of the
 polypeptides encoded in the various clones examined sug-
 gest efficient detection and immunization systems may
 include panels of HCV polypeptides/epitopes.

35 Expression of HCV Epitopes in Yeast
 Three different yeast expression vectors which
 allow the insertion of HCV cDNA into three different read-

ing frames are constructed. The construction of one of the vectors, pAB24C100-3 is described in EPO Pub. No. 318,216. In the studies below, the HCV cDNA from the clones listed in supra. in the antigenicity mapping study using the E. coli expressed products are substituted for the C100 HCV cDNA. The construction of the other vectors replaces the adaptor described in the above E. coli studies with one of the following adaptors:

Adaptor 1

ATT TTG AAT TCC TAA TGA G
AC TTA AGG ATT ACT CAG CT

Adaptor 2

AAT TTG GAA TTC TAA TGA G
AC CTT AAG ATT ACT CAG CT.

The inserted HCV cDNA is expressed in yeast transformed with the vectors, using the expression conditions described supra. for the expression of the fusion polypeptide, C100-3. The resulting polypeptides are screened using the sera from individuals with NANBH, described supra. for the screening of immunogenic polypeptides encoded in HCV cDNAs expressed in E. coli.

Comparison of the Hydrophobic Profiles of HCV Polyproteins with West Nile Virus Polyprotein and with Dengue Virus NS1

The hydrophobicity profile of an HCV polyprotein segment was compared with that of a typical Flavivirus, West Nile virus. The polypeptide sequence of the West Nile virus polyprotein was deduced from the known polynucleotide sequences encoding the non-structural

proteins of that virus. The HCV polyprotein sequence was deduced from the sequence of overlapping cDNA clones. The profiles were determined using an antigen program which
5 uses a window of 7 amino acid width (the amino acid in question, and 3 residues on each side) to report the average hydrophobicity about a given amino acid residue. The parameters giving the reactive hydrophobicity for each amino acid residue are from Kyte and Doolittle (1982).
10 Fig. 19 shows the hydrophobic profiles of the two polyproteins; the areas corresponding to the non-structural proteins of West Nile virus, ns1 through ns5, are indicated in the figure. As seen in the figure, there is a general similarity in the profiles of the HCV
15 polyprotein and the West Nile virus polyprotein.

The sequence of the amino acids encoded in the 5'-region of HCV cDNA shown in Fig. 16 has been compared with the corresponding region of one of the strains of Dengue virus, described supra., with respect to the
20 profile of regions of hydrophobicity and hydrophilicity (data not shown). This comparison indicated that the polypeptides from HCV and Dengue encoded in this region, which corresponds to the region encoding NS1 (or a portion thereof), have a similar hydrophobic/hydrophilic profile.
25

The similarity in hydrophobicity profiles, in combination with the previously identified homologies in the amino acid sequences of HCV and Dengue Flavivirus in EP 0,218,316 suggests that HCV is related to these members of the Flavivirus family.
30

Characterization of the Putative Polypeptides Encoded Within the HCV ORF

The sequence of the HCV cDNA sense strand, shown in Fig. 17, was deduced from the overlapping HCV cDNAs in
35 the various clones described in EPO Pub. No. 318,216 and those described supra. It may be deduced from the

sequence that the HCV genome contains primarily one long continuous ORF, which encodes a polyprotein. In the sequence, nucleotide number 1 corresponds to the first nucleotide of the initiator MET codon; minus numbers indicate that the nucleotides are that distance away in the 5'-direction (upstream), while positive numbers indicate that the nucleotides are that distance away in the 3'-direction (downstream). The composite sequence shows the "sense" strand of the HCV cDNA.

The amino acid sequence of the putative HCV polyprotein deduced from the HCV cDNA sense strand sequence is also shown in Fig. 17, where position 1 begins with the putative initiator methionine.

Possible protein domains of the encoded HCV polyprotein, as well as the approximate boundaries, are the following (the polypeptides identified within the parentheses are those which are encoded in the Flavivirus domain):

	<u>Putative Domain</u>	<u>Approximate Boundary</u> <u>(amino acid nos.)</u>
25	"C" (nucleocapsid protein)	1-120
	"E" (Virion envelope protein(s) and possibly matrix (M) proteins	120-400
30	"NS1" (complement fixation antigen?)	400-660
	"NS2" (unknown function)	660-1050
35	"NS3" (protease?)	1050-1640

"NS4" (unknown function) 1640-2000

"NS5" (polymerase) 2000-? end

5

It should be noted, however, that hydrophobicity profiles (described infra), indicate that HCV diverges from the Flavivirus model, particularly with respect to the region upstream of NS2. Moreover, the boundaries indicated are not intended to show firm demarcations between the putative polypeptides.

15

The Hydrophilic and Antigenic Profile of the Polypeptide

Profiles of the hydrophilicity/hydrophobicity and the antigenic index of the putative polyprotein encoded in the HCV cDNA sequence shown in Fig. 16 were determined by computer analysis. The program for hydrophilicity/hydrophobicity was as described supra. The antigenic index results from a computer program which relies on the following criteria: 1) surface probability, 2) prediction of alpha-helicity by two different methods; 3) prediction of beta-sheet regions by two different methods; 4) prediction of U-turns by two different methods; 5) hydrophilicity/hydrophobicity; and flexibility. The traces of the profiles generated by the computer analyses are shown in Fig. 20. In the hydrophilicity profile, deflection above the abscissa indicates hydrophilicity, and below the abscissa indicates hydrophobicity. The probability that a polypeptide region is antigenic is usually considered to increase when there

is a deflection upward from the abscissa in the hydrophilic and/or antigenic profile. It should be noted, however, that these profiles are not necessarily indicators of the strength of the immunogenicity of a polypeptide.

Identification of Co-linear Peptides in HCV and Flaviviruses

The amino acid sequence of the putative polyprotein encoded in the HCV cDNA sense strand was compared with the known amino acid sequences of several members of Flaviviruses. The comparison shows that homology is slight, but due to the regions in which it is found, it is probably significant. The conserved co-linear regions are shown in Fig. 21. The amino acid numbers listed below the sequences represent the number in the putative HCV polyprotein (See Fig. 17.)

The spacing of these conserved motifs is similar between the Flaviviruses and HCV, and implies that there is some similarity between HCV and these flaviviral agents.

The following listed materials are on deposit under the terms of the Budapest Treaty with the American Type Culture Collection (ATCC), 12301 Parklawn Dr., Rockville, Maryland 20852, and have been assigned the following Accession Numbers.

35

	<u>lambda-gt11</u>	<u>ATCC No.</u>	<u>Deposit Date</u>
	HCV cDNA library	40394	1 Dec. 1987
5	clone 81	40388	17 Nov. 1987
	clone 91	40389	17 Nov. 1987
	clone 1-2	40390	17 Nov. 1987
	clone 5-1-1	40391	18 Nov. 1987
	clone 12f	40514	10 Nov. 1988
10	clone 35f	40511	10 Nov. 1988
	clone 15e	40513	10 Nov. 1988
	clone K9-1	40512	10 Nov. 1988
	JSC 308	20879	5 May 1988
	pS356	67683	29 April 1988

- 15 In addition, the following deposits were made on 11 May 1989.

	<u>Strain</u>	<u>Linkers</u>	<u>ATCC No.</u>
20	D1210 (Cf1/5-1-1)	EF	67967
	D1210 (Cf1/81)	EF	67968
	D1210 (Cf1/CA74a)	EF	67969
	D1210 (Cf1/35f)	AB	67970
	D1210 (Cf1/279a)	EF	67971
25	D1210 (Cf1/C36)	CD	67972
	D1210 (Cf1/13i)	AB	67973
	D1210 (Cf1/C33b)	EF	67974
	D1210 (Cf1/CA290a)	AB	67975
	HB101 (AB24/C100 #3R)		67976

- 30 The following derivatives of strain D1210 were deposited on 3 May 1989.

	<u>Strain Derivative</u>	<u>ATCC No.</u>
	pCF1CS/C8f	67956
	pCF1AB/C12f	67952
5	pCF1EF/14c	67949
	pCF1EF/15e	67954
	pCF1AB/C25c	67958
	pCF1EF/C33c	67953
	pCF1EF/C33f	67050
10	pCF1CD/33g	67951
	pCF1CD/C39c	67955
	pCF1EF/C40b	67957
	pCF1EF/CA167b	67959

15 The following strains were deposited on May 12, 1989.

	<u>Strain</u>	<u>ATCC No.</u>
	Lambda gt11(C35)	40603
	Lambda gt10(beta-5a)	40602
20	D1210 (C40b)	67980
	D1210 (M16)	67981

25 Upon allowance and issuance of this application as a
United States Patent, all restriction on availability of
these deposits will be irrevocably removed; and access to
the designated deposits will be available during pendency
of the above-named application to one determined by the
Commissioner to be entitled thereto under 37 CFR 1.14 and
35 USC 1.22. Moreover, the designated deposits will be
30 maintained for a period of thirty (30) years from the date
of deposit, or for five (5) years after the last request
for the deposit; or for the enforceable life of the U.S.
patent, whichever is longer. The deposited materials
mentioned herein are intended for convenience only, and
35 are not required to practice the present invention in view

of the descriptions herein, and in addition these materials are incorporated herein by reference.

5

Industrial Applicability

The invention, in the various manifestations disclosed herein, has many industrial uses, some of which are the following. The HCV cDNAs may be used for the design of probes for the detection of HCV nucleic acids in
10 samples. The probes derived from the cDNAs may be used to detect HCV nucleic acids in, for example, chemical synthetic reactions. They may also be used in screening programs for anti-viral agents, to determine the effect of the agents in inhibiting viral replication in cell culture
15 systems, and animal model systems. The HCV polynucleotide probes are also useful in detecting viral nucleic acids in humans, and thus, may serve as a basis for diagnosis of HCV infections in humans.

In addition to the above, the cDNAs provided
20 herein provide information and a means for synthesizing polypeptides containing epitopes of HCV. These polypeptides are useful in detecting antibodies to HCV antigens. A series of immunoassays for HCV infection, based on recombinant polypeptides containing HCV epitopes
25 are described herein, and will find commercial use in diagnosing HCV induced NANBH, in screening blood bank donors for HCV-caused infectious hepatitis, and also for detecting contaminated blood from infectious blood donors. The viral antigens will also have utility in monitoring
30 the efficacy of anti-viral agents in animal model systems. In addition, the polypeptides derived from the HCV cDNAs disclosed herein will have utility as vaccines for treatment of HCV infections.

The polypeptides derived from the HCV cDNAs,
35 besides the above stated uses, are also useful for raising anti-HCV antibodies. Thus, they may be used in anti-HCV

vaccines. However, the antibodies produced as a result of immunization with the HCV polypeptides are also useful in detecting the presence of viral antigens in samples. Thus, they may be used to assay the production of HCV polypeptides in chemical systems. The anti-HCV antibodies may also be used to monitor the efficacy of anti-viral agents in screening programs where these agents are tested in tissue culture systems. They may also be used for passive immunotherapy, and to diagnose HCV caused NANBH by allowing the detection of viral antigen(s) in both blood donors and recipients. Another important use for anti-HCV antibodies is in affinity chromatography for the purification of virus and viral polypeptides. The purified virus and viral polypeptide preparations may be used in vaccines. However, the purified virus may also be useful for the development of cell culture systems in which HCV replicates.

Antisense polynucleotides may be used as inhibitors of viral replication.

For convenience, the anti-HCV antibodies and HCV polypeptides, whether natural or recombinant, may be packaged into kits.

CLAIMS

- 5 1. A recombinant polynucleotide comprising a
sequence derived from HCV cDNA, wherein the HCV cDNA is in
clone 13i, or clone 26j, or clone 59a, or clone 84a, or
clone CA156e, or clone 167b, or clone pil4a, or clone
CA216a, or clone CA290a, or clone ag30a, or clone 205a, or
10 clone 18g, or clone 16jh, or wherein the HCV cDNA is of a
sequence indicated by nucleotide numbers -319 to 1348 or
8659 to 8866 in Fig. 17.
- 15 2. A recombinant polynucleotide according to
claim 1, encoding an epitope of HCV.
3. A recombinant vector comprising the
polynucleotide of claim 1 or claim 2.
- 20 4. A host cell transformed with the vector of
claim 3.
5. A recombinant expression system comprising
an open reading frame (ORF) of DNA derived from the re-
combinant polynucleotide of claim 1 or claim 2, wherein
25 the ORF is operably linked to a control sequence compat-
ible with a desired host.
6. A cell transformed with the recombinant
expression system of claim 5.
30
7. A polypeptide produced by the cell of claim
6.
- 35 8. A purified polypeptide comprising an epitope
encoded within HCV cDNA wherein the HCV cDNA is of a

sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17.

5 9. An immunogenic polypeptide produced by a
cell transformed with a recombinant expression vector
comprising an ORF of DNA derived from HCV cDNA, wherein
the HCV cDNA is comprised of a sequence derived from the
10 HCV cDNA sequence in clone CA279a, or clone CA74a, or
clone 13i, or clone CA290a, or clone 33C or clone 40b, or
clone 33b, or clone 25c, or clone 14c, or clone 8f, or
clone 33f, or clone 33g, or clone 39c, or clone 15e, and
wherein the ORF is operably linked to a control sequence
15 compatible with a desired host.

10. A peptide comprising an HCV epitope,
wherein the peptide is of the formula

20 AA_x-AA_y ,

wherein x and y designate amino acid numbers shown in Fig.
17, and wherein the peptide is selected from the group
consisting of AA1-AA25, AA1-AA50, AA1-AA84, AA9-AA177,
25 AA1-AA10, AA5-AA20, AA20-AA25, AA35-AA45, AA50-AA100,
AA40-AA90, AA45-AA65, AA65-AA75, AA80-90, AA99-AA120,
AA95-AA110, AA105-AA120, AA100-AA150, AA150-AA200,
AA155-AA170, AA190-AA210, AA200-AA250, AA220-AA240,
AA245-AA265, AA250-AA300, AA290-AA330, AA290-305,
AA300-AA350, AA310-AA330, AA350-AA400, AA380-AA395,
30 AA405-AA495, AA400-AA450, AA405-AA415, AA415-AA425,
AA425-AA435, AA437-AA582, AA450-AA500, AA440-AA460,
AA460-AA470, AA475-AA495, AA500-AA550, AA511-AA690, AA515-
AA550, AA550-AA600, AA550-AA625, AA575-AA605, AA585-AA600,
AA600-AA650, AA600-AA625, AA635-AA665, AA650-AA700,
35 AA645-AA680, AA700-AA750, AA700-AA725, AA700-AA750,
AA725-AA775, AA770-AA790, AA750-AA800, AA800-AA815,

AA825-AA850, AA850-AA875, AA800-AA850, AA920-AA990,
AA850-AA900, AA920-AA945, AA940-AA965, AA970-AA990,
5 AA950-AA1000, AA1000-AA1060, AA1000-AA1025, AA1000-AA1050,
AA1025-AA1040, AA1040-AA1055, AA1075-AA1175,
AA1050-AA1200, AA1070-AA1100, AA1100-AA1130,
AA1140-AA1165, AA1192-AA1457, AA1195-AA1250,
AA1200-AA1225, AA1225-AA1250, AA1250-AA1300,
10 AA1260-AA1310, AA1260-AA1280, AA1266-AA1428,
AA1300-AA1350, AA1290-AA1310, AA1310-AA1340, AA1345-
AA1405, AA1345-AA1365, AA1350-AA1400, AA1365-AA1380,
AA1380-AA1405, AA1400-AA1450, AA1450-AA1500,
AA1460-AA1475, AA1475-AA1515, AA1475-AA1500,
AA1500-AA1550, AA1500-AA1515, AA1515-AA1550,
15 AA1550-AA1600, AA1545-AA1560, AA1569-AA1931,
AA1570-AA1590, AA1595-AA1610, AA1590-AA1650,
AA1610-AA1645, AA1650-AA1690, AA1685-AA1770,
AA1689-AA1805, AA1690-AA1720, AA1694-AA1735,
AA1720-AA1745, AA1745-AA1770, AA1750-AA1800,
20 AA1775-AA1810, AA1795-AA1850, AA1850-AA1900,
AA1900-AA1950, AA1900-AA1920, AA1916-AA2021,
AA1920-AA1940, AA1949-AA2124, AA1950-AA2000,
AA1950-AA1985, AA1980-AA2000, AA2000-AA2050,
AA2005-AA2025, AA2020-AA2045, AA2045-AA2100,
25 AA2045-AA2070, AA2054-AA2223, AA2070-AA2100,
AA2100-AA2150, AA2150-AA2200, AA2200-AA2250,
AA2200-AA2325, AA2250-AA2330, AA2255-AA2270,
AA2265-AA2280, AA2280-AA2290, AA2287-AA2385,
AA2300-AA2350, AA2290-AA2310, AA2310-AA2330,
30 AA2330-AA2350, AA2350-AA2400, AA2348-AA2464,
AA2345-AA2415, AA2345-AA2375, AA2370-AA2410,
AA2371-AA2502, AA2400-AA2450, AA2400-AA2425,
AA2415-AA2450, AA2445-AA2500, AA2445-AA2475,
AA2470-AA2490, AA2500-AA2550, AA2505-AA2540,
35 AA2535-AA2560, AA2550-AA2600, AA2560-AA2580,
AA2600-AA2650, AA2605-AA2620, AA2620-AA2650,

AA2640-AA2660, AA2650-AA2700, AA2655-AA2670,
AA2670-AA2700, AA2700-AA2750, AA2740-AA2760,
AA2750-AA2800, AA2755-AA2780,
5 AA2780-AA2830, AA2785-AA2810, AA2796-AA2886,
AA2810-AA2825, AA2800-AA2850, AA2850-AA2900,
AA2850-AA2865, AA2885-AA2905, AA2900-AA2950,
AA2910-AA2930, AA2925-AA2950, AA2945-end(C' terminal).

10 11. A polypeptide comprised of the peptide of
claim 10.

12. An immunogenic polypeptide attached to a
solid substrate, wherein the polypeptide is according to
15 claim 7, or claim 8, or claim 9, or claim 10, or claim 11,
or wherein the polypeptide is comprised of an epitope
encoded within HCV cDNA wherein the HCV cDNA is of a
sequence indicated by nucleotide numbers -319 to 1348 or
8659 to 8866 in Fig. 17.
20

13. A monoclonal antibody directed against an
epitope encoded in HCV cDNA, wherein the HCV cDNA is of a
sequence indicated by nucleotide numbers -319 to 1348 or
8659 to 8866 in Fig. 17, or is the sequence present in
25 clone 13i, or clone 26j, or clone 59a, or clone 84a, or
clone CA156e, or clone 167b, or clone pil4a, or clone
CA216a, or clone CA290a, or clone ag30a, or clone 205a, or
clone 18g, or clone 16jh.

30 14. A preparation of purified polyclonal anti-
bodies directed against a polypeptide comprised of an
epitope encoded within HCV cDNA, wherein the HCV cDNA is
of a sequence indicated by nucleotide numbers -319 to 1348
or 8659 to 8866 in Fig. 17, or is the sequence present in
35 clone 13i, or clone 26j, or clone 59a, or clone 84a, or
clone CA156e, or clone 167b, or clone pil4a, or clone

CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh.

5 15. A polynucleotide probe for HCV, wherein the probe is comprised of an HCV sequence derived from an HCV cDNA sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or from the complement of the HCV cDNA sequence.

10

16. A kit for analyzing samples for the presence of polynucleotides from HCV comprising a polynucleotide probe containing a nucleotide sequence of about 8 or more nucleotides, wherein the nucleotide
15 sequence is derived from HCV cDNA which is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, wherein the polynucleotide probe is in a suitable container.

20

17. A kit for analyzing samples for the presence of an HCV antigen comprising an antibody which reacts immunologically with an HCV antigen, wherein the antigen contains an epitope encoded within HCV cDNA which is of a sequence indicated by nucleotide numbers -319 to 1348 or
25 8659 to 8866 in Fig. 17, or wherein the HCV cDNA is in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone p114a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh.

30

18. A kit for analyzing samples for the presence of an HCV antibody comprising an antigenic polypeptide containing an HCV epitope encoded within HCV cDNA which is of a sequence indicated by nucleotide
35 numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is in clone 13i, or clone 26j, or clone 59a, or clone 84a, or

5 clone CA156e, or clone 167b, or clone pil4a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh.

10 19. A kit for analyzing samples for the presence of an HCV antibody comprising an antigenic polypeptide expressed from HCV cDNA in clone CA279a, or clone CA74a, or clone 13i, or clone CA290a, or clone 33C or clone 40b, or clone 33b, or clone 25c, or clone 14c, or clone 8f, or clone 33f, or clone 33g, or clone 39c, or clone 15e, wherein the antigenic polypeptide is present in a suitable container.

15 20. A method for detecting HCV nucleic acids in a sample comprising:

20 (a) reacting nucleic acids of the sample with a polynucleotide probe for HCV, wherein the probe is comprised of an HCV sequence derived from an HCV cDNA sequence is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, and wherein the reacting is under conditions which allow the formation of a polynucleotide duplex between the probe and the HCV nucleic acid from the sample,

25 (b) detecting a polynucleotide duplex which contains the probe, formed in step (a).

30 21. An immunoassay for detecting an HCV antigen comprising:

35 (a) incubating a sample suspected of containing an HCV antigen with an antibody directed against an HCV epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is the sequence present in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone pil4a, or clone

5 CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, and wherein the incubating is under conditions which allow formation of an antigen-antibody complex; and (b) detecting an antibody-antigen complex formed in step (a) which contains the antibody.

10 22. An immunoassay for detecting antibodies directed against an HCV antigen comprising:

15 (a) incubating a sample suspected of containing anti-HCV antibodies with an antigen polypeptide containing an epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is the sequence present in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone pil4a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, and wherein the incubating is under conditions which allow formation of an antigen-antibody complex; and

20 (b) detecting an antibody-antigen complex formed in step (a) which contains the antigen polypeptide.

25 23. An immunoassay for detecting antibodies directed against an HCV antigen comprising:

30 (a) incubating a sample suspected of containing anti-HCV antibodies with the polypeptide of claim 9, under conditions which allow formation of an antigen-antibody complex; and

(b) detecting an antibody-antigen complex formed in step (a) which contains the antigen polypeptide.

35 24. A vaccine for treatment of HCV infection comprising an immunogenic polypeptide containing an HCV epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or

8659 to 8866 in Fig. 17 or is the sequence present in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone p114a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, and wherein the immunogenic polypeptide is present in a pharmacologically effective dose in a pharmaceutically acceptable excipient.

10

25. A method for producing antibodies to HCV comprising administering to an individual an isolated immunogenic polypeptide containing an HCV epitope encoded in HCV cDNA, wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17, or is of the sequence present in clone CA279a, or clone CA74a, or clone 13i, or clone CA290a, or clone 33C or clone 40b, or clone 33b, or clone 25c, or clone 14c, or clone 8f, or clone 33f, or clone 33g, or clone 39c, or clone 15e, and wherein the immunogenic polypeptide is present in a pharmacologically effective dose in a pharmaceutically acceptable excipient.

15

20

26. An antisense polynucleotide derived from HCV cDNA, wherein the HCV cDNA is that shown in Fig. 17.

25

27. A method for preparing purified fusion polypeptide C100-3 comprising:

30

(a) providing a crude cell lysate containing polypeptide C100-3,

(b) treating the crude cell lysate with an amount of acetone which causes the polypeptide to precipitate,

(c) isolating and solubilizing the precipitated material,

35

(d) isolating the C100-3 polypeptide by anion exchange chromatography, and

(e) further isolating the C100-3 polypeptide of step (d) by gel filtration.

5

28. A method for preparing an HCV polypeptide comprising:

10 (a) providing a host cell transformed with a recombinant expression system comprising an open reading frame (ORF) of DNA derived from HCV cDNA, wherein the HCV cDNA is in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone p114a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, or wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 15 1348 or 8659 to 8866 in Fig. 17, wherein the ORF is operably linked to a control sequence compatible with a desired host; and

20 (b) incubating the host cell under conditions with allow expression of the HCV polypeptide.

29. A method for preparing an immunogenic HCV polypeptide comprising:

25 (a) providing a host cell transformed with a recombinant expression vector comprising an ORF of DNA derived from HCV cDNA, wherein the HCV cDNA is comprised of a sequence derived from the HCV cDNA sequence in clone CA279a, or clone CA74a, or clone 13i, or clone CA290a, or clone 33c, or clone 40b, or clone 33b, or clone 25c, or clone 14c, or clone 8f, or clone 33f, or clone 33g, or 30 clone 39c, or clone 15e, wherein the OEF is operably linked to a control sequence compatible with the desired host; and

35 (b) incubating the host cell under conditions which allow expression of the HCV polypeptide.

30. A method for preparing a host cell

transformed with a recombinant polynucleotide comprising a sequence of HCV cDNA derived from the HCV cDNA in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone 5
CA156e, or clone 167b, or clone pil4a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, or wherein the HCV cDNA is of a sequence indicated by nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17 comprising:

- 10 (a) providing a host cell capable of transformation;
- (b) providing the recombinant polynucleotide;
- and
- (c) incubating (a) with (b) under conditions
- 15 which allow transformation of the host cell with the polynucleotide.

31. A method for preparing a recombinant polynucleotide comprised of a sequence of HCV cDNA derived from the HCV cDNA in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e, or clone 167b, or clone pil4a, or clone CA216a, or clone CA290a, or clone ag30a, or clone 205a, or clone 18g, or clone 16jh, or wherein the HCV cDNA is of a sequence indicated by
20 nucleotide numbers -319 to 1348 or 8659 to 8866 in Fig. 17 comprising:

- 25 (a) providing a host cell transformed with the recombinant polynucleotide; and
- (b) isolating said polynucleotide from said
- 30 host cell.

32. A method for preparing blood free of HCV comprising:

- 35 (a) providing a sample of blood suspected of containing HCV and anti-HCV antibodies;

- (b) providing an immunogenic polypeptide prepared according to claim 28 or 29;
- 5 (c) incubating the sample of (a) with the immunogenic polypeptide of (b) under conditions which allow the formation of antibody-HCV polypeptide complexes;
- (d) detecting the complexes formed in step (c); and
- 10 (e) saving the blood from which complexes were not detected in (d).

33. A method for preparing blood free of HCV comprising:

- 15 (a) providing nucleic acids from a sample of blood suspected of containing HCV polynucleotides;
- (b) providing a probe for HCV, wherein the probe is comprised of an HCV sequence derived from an HCV cDNA which is of a sequence indicated by nucleotide
- 20 numbers -319 to 1348 or 8659 to 8866 in Fig. 17,
- (c) reacting (a) with (b) under conditions which allow the formation of a polynucleotide duplex between the probe and the HCV nucleic acid from the sample;
- 25 (d) detecting a polynucleotide which contains the probe, formed in step (c); and
- (e) saving the blood from which complexes were not detected in (d).

34. A method for producing a hybridoma which produces anti-HCV monoclonal antibodies comprising:

- 30 (a) immunizing an individual with an immunogenic polypeptide containing an epitope encoded in HCV cDNA, wherein the HCV cDNA is HCV cDNA in clone 13i, or clone 26j, or clone 59a, or clone 84a, or clone CA156e,
- 35 or clone 167b, or clone pil4a, or clone CA216a, or clone

CA290a, or clone ag30a, or clone 205a, or clone 18g, or
clone 16jh, or wherein the HCV cDNA is of a sequence
indicated by nucleotide numbers -319 to 1348 or 8659 to
5 8866 in Fig. 17; or

(b) immunizing an individual with an
immunogenic polypeptide prepared according to claim 29;

(c) immortalizing antibody producing cells from
the immunized individual;

10 (d) selecting an immortal cell which produces
antibodies which react with an HCV epitope in the
immunogenic polypeptide of (a) or (b); and

(e) growing said immortal cell.

15

20

25

30

35

FIG. 1 Translation of DNA 12f

IlePheLysIleArgMetTyrValGlyGlyValGluHisArgLeuGluAlaAlaCysAsn
1 CCATATTTAAAATCAGGATGTACGTGGGAGGGGTGGAACACAGGCTGGAAGCTGCCTGCA
GGTATAAATTTTAGTCCTACATGCACCCCTCCCAGCTTGTGTCCGACCTTCGACGGACGT

TrpThrArgGlyGluArgCysAspLeuGluAspArgAspArgSerGluLeuSerProLeu
61 ACTGGACGCGGGGCGAACGTTGCGATCTGGAAGACAGGGACAGGTCGAGCTCAGCCCGT
TGACCTGCGCCCGCTTGCAACGCTAGACCTTCTGTCCCTGTCCAGGCTCGAGTCGGGCA

LeuLeuThrThrThrGlnTrpGlnValLeuProCysSerPheThrThrLeuProAlaLeu
121 TACTGCTGACCACTACACAGTGGCAGGTCTCCCGTGTTCCTTCACAACCCTACCAGCCT
ATGACGACTGGTGATGTGTCACCGTCCAGGAGGGCACAAGGAAGTGTGGGATGGTCGGA

SerThrGlyLeuIleHisLeuHisGlnAsnIleValAspValGlnTyrLeuTyrGlyVal
181 TGTCCACCGGCTCATCCACCTCCACCAGAACATTGTGGACGTGCAGTACTTGTACGGGG
ACAGGTGGCCGGAGTAGGTGGAGGTGGTCTTGTAACACCTGCACGTCATGAACATGCCCC

GlySerSerIleAlaSerTrpAlaIleLysTrpGluTyrValValLeuLeuPheLeuLeu
241 TGGGGTCAAGCATCGCGTCCTGGGCCATTAAGTGGGAGTACGTCGTTCTCCTGTTCTTC
ACCCAGTTCGTAGCGCAGGACCGGTAATTACCCCTCATGCAGCAAGAGGACAAGGAAG

LeuAlaAspAlaArgValCysSerCysLeuTrpMetMetLeuLeuIleSerGlnAlaGlu
301 TGCTTGACAGACGCGCGCTCTGCTCCTGCTTGTGGATGATGCTACTCATATCCCAAGCGG
ACGAACGTCTGCGCGCGCAGACGAGGACGAACACCTACTACGATGAGTATAGGGTTCGCC

AlaAlaLeuGluAsnLeuValIleLeuAsnAlaAlaSerLeuAlaGlyThrHisGlyLeu
361 AGGCGGCTTTGGAGAACCTCGTAATACTTAATGCAGCATCCCTGGCCGGGACGCACGGTC
TCCGCCGAAACCTCTTGAGCATTATGAATTACGTCGTAGGGACCGGCCCTGCGTGCCAG

Val
421 TTGTATC
AACATAG

SUBSTITUTE SHEET

3 / 40

ValCysSerCysLeuTrpMetMetLeuLeuIleSerGlnAlaGluAlaAlaLeuGluAsn
841 GCGTCTGCTCCTGCTTGTGGATGATGCTACTCATATCCCAAGCGGAAGCGGCTTTGGAGA
CGCAGACGAGGACGAACACCTACTACGATGAGTATAGGGTTCCGCTTCGCCGAAACCTCT

LeuValIleLeuAsnAlaAlaSerLeuAlaGlyThrHisGlyLeuValSerPheLeuVal
901 ACCTCGTAATACTTAATGCAGCATCCCTGGCCGGGACGCACGGTCTTGTATCCTTCCTCG
TGGAGCATTATGAATTACGTCGTAGGGACCGGCCCTGCGTGCCAGAACATAGGAAGGAGC

PhePheCysPheAlaTrpTyrLeuLysGlyLysTrpValProGlyAlaValTyrThrPhe
961 TGTTCCTTCTGCTTTGCATGGTATCTGAAGGGTAAGTGGGTGCCGGGAGCGGTCTACACCT
ACAAGAAGACGAAACGTACCATAGACTTCCCATTCACCCACGGGGCTCGCCAGATGTGGA

TyrGlyMetTrpProLeuLeuLeuLeuLeuAlaLeuProGlnArgAlaTyrAlaLeu
1021 TCTACGGGATGTGGCCTCTCCTCCTGCTCCTGTTGGCGTTGCCCGAGCGGGCGTACGCCG
AGATGCCCTACACCGGAGAGGAGGACGAGGACAACCGCAACGGGGTTCGCCCGCATGCCGC

AspThrGluValAlaAlaSerCysGlyGlyValValLeuValGlyLeuMetAlaLeuThr
1081 TGGACACGGAGGTGGCCGCGTCGTGTGGCGGTGTGTTCTCGTCGGGTGATGGCGCTAA
ACCTGTGCCTCCACCGCGCAGCACACCGCCACAACAAGAGCAGCCCACTACCGCGATT

LeuSerProTyrTyrLysArgTyrIleSerTrpCysLeuTrpTrpLeuGlnTyrPheLeu
1141 CTCTGTCAACATATTACAAGCGCTATATCAGCTGGTGCTTGTGGTGGCTTCAGTATTTTC
GAGACAGTGGTATAATGTTCCGATATAGTCGACCACGAACACCACCGAAGTCATAAAG

ThrArgValGluAlaGlnLeuHisValTrpIleProProLeuAsnValArgGlyGlyArg
1201 TGACCAGAGTGAAGCGCAACTGCACGTGTGGATTCCCCCCTCAACGTCCGAGGGGGGC
ACTGGTCTCACCTTCGCGTTGACGTGCACACCTAAGGGGGGAGTTGCAGGCTCCCCCG

AspAlaValIleLeuLeuMetCysAlaValHisProThrLeuValPheAspIleThrLys
1261 GCGACGCTGTCATCTTACTCATGTGTGCTGTACACCGACTCTGGTATTTGACATCACCA
CGCTGCGACAGTAGAATGAGTACACACGACATGTGGGCTGAGACCATAAACTGTAGTGGT

LeuLeuLeuAlaValPheGlyProLeuTrpIleLeuGlnAla
1321 AATTGCTGCTGGCGTCTTCGGACCCCTTTGGATTCTTCAAGCCAG
TTAACGACGACCGGCAGAAGCCTGGGGAAACCTAAGAAGTTCGGTC

FIG. 2-2

SUBSTITUTE SHEET

4 / 40

FIG. 3

```

1  GlyAlaGlyLysArgValTyrTyrLeuThrArgAspProThrThrProLeuAlaArgAla
   CGGCGCTGGAAAGAGGGTCTACTACCTCACCCGTGACCCTACAACCCCCCTCGCGAGAGC
   GCCGCGACCTTTCTCCAGATGATGGAGTGGGCACTGGGATGTTGGGGGAGCGCTCTCG

61  AlaTrpGluThrAlaArgHisThrProValAsnSerTrpLeuGlyAsnIleIleMetPhe
   TCGTGAGGAGACAGCAAGACACACTCCAGTCAATTCCTGGCTAGGCAACATAATCATGTT
   ACGCACCTCTGTGCTTCTGTGTGAGGTCAAGTAAGGACCGATCCGTTGTATTAGTACAA

121  AlaProThrLeuTrpAlaArgMetIleLeuMetThrHisPhePheSerValLeuIleAla
   TGCCCCCACACTGTGGGCGAGGATGATACTGATGACCCATTTCTTTAGCGTCCTTATAGC
   ACGGGGGTGTGACACCCGCTCCTACTATGACTACTGGGTAAAGAAATCGCAGGAATATCG

181  ArgAspGlnLeuGluGlnAlaLeuAspCysGluIleTyrGlyAlaCysTyrSerIleGlu
   CAGGGACCAGCTTGAACAGGCCCTCGATTGCGAGATCTACGGGGCTGCTACTCCATAGA
   GTCCCTGGTCGAACCTGTCCGGGAGCTAACGCTCTAGATGCCCCGACGATGAGGTATCT

241  ProLeuAspLeuProProIleIleGlnArgLeu
   ACCACTTGATCTACCTCCAATCATTCAAAGACTC
   TGGTGAAC TAGATGGAGGTTAGTAAGTTTCTGAG

```

FIG. 5

Translation of DNA 26j

```

1  LeuPheTyrHisHisLysPheAsnSerSerGlyCysProGluArgLeuAlaSerCysArg
   GCTTTTCTATCACCACAAGTTCAACTCTTCAGGCTGTCCTGAGAGGCTAGCCAGCTGCCG
   CGAAAAGATAGTGGTGTTCAAGTTGAGAAGTCCGACAGGACTCTCCGATCGGTCGACGGC

61  ProLeuThrAspPheAspGlnGlyTrpGlyProIleSerTyrAlaAsnGlySerGlyPro
   ACCCCTTACCGATTTTGACCAGGGCTGGGGCCCTATCAGTTATGCCAACGGAAGCGGCCC
   TGGGGAATGGCTAAACTGGTCCCGACCCCGGATAGTCAATACGGTTGCCTTCGCCGGG

121  AspGlnArgProTyrCysTrpHisTyrProProLysProCysGlyIleValProAlaLys
   CGACCAGCGCCCCTACTGCTGGCACTACCCCCAAACCTTGCGGTATTGTGCCCCGAA
   GCTGGTCGCGGGGATGACGACCGTGATGGGGGTTTGAACGCCATAACACGGGCGCTT

181  SerValCysGlyProValTyrCysPheThrProSerProValValVal
   GAGTGTGTGTGGTCCGGTATATTGCTTCACTCCAGCCCCGTGGTGGTGGG
   CTCACACACACCAGGCCATATAACGAAGTGAGGGTCGGGGCACCACCACC

```

---Overlap with 131---

SUBSTITUTE SHEET

FIG. 4

Translation of DNA 131

1 ProSerProValValValGlyThrThrAspArgSerGlyAlaProThrTyrSerTrpGly
 CTCCCAGCCCCGTGGTGGTGGGAACGACCGACAGGTCGGGCGCGCTACCTACAGCTGGG
 GAGGGTCGGGCGACCAACCACTTGTCTGGCTGTCCAGCCCCGCGCGGATGGATGTGACCC

61 GluAsnAspThrAspValPheValLeuAsnAsnThrArgProProLeuGlyAsnTrpPhe
 GTGAAAATGATACGGACGTCTTCGTCCTTAACAATACCAGGCCACCGCTGGGCAATTGGT
 CACTTTTACTATGCCTGCAGAAGCAGGAATTGTTATGGTCCGGTGGCGACCCGTTAACCA

121 GlyCysThrTrpMetAsnSerThrGlyPheThrLysValCysGlyAlaProProCysVal
 TCGGTTGTACCTGGATGAACCTCAACTGGATTACCAAAGTGTGCGGAGCGCTCCTTGTG
 AGCCAACATGGACCTACTTGAGTTGACCTAAGTGGTTTCACACGCCTCGCGGAGGAACAC

181 IleGlyGlyAlaGlyAsnAsnThrLeuHisCysProThrAspCysPheArgLysHisPro
 TCATCGGAGGGGCGGGCAACAACACCTGCACTGCCCCACTGATTGCTTCGCAAGCATC
 AGTAGCCTCCCCGCCCCGTGTTGTGGGACGTGACGGGGTGACTAACGAAGCGTTCTGTAG

241 AspAlaThrTyrSerArgCysGlySerGlyProTrpLeuThrProArgCysLeuValAsp
 CGGACGCCACATACTCTCGGTGCGGCTCCGGTCCCTGGCTCACACCCAGGTGCTTGGTCTG
 GCCTGCGGTGTATGAGAGCCACGCCGAGGCCAGGGACCGAGTGTGGGTCCACGGACCAGC

301 TyrProTyrArgLeuTrpHisTyrProCysThrIleAsnTyrThrIlePheLysIleArg
 ACTACCCGTATAGGCTTTGGCATTATCCTTGTACCATCAACTACACCATATTTAAAATCA
 TGATGGGCATATCCGAAACCGTAATAGGAACATGGTAGTTGATGTGGTATAAATTTTAGT

361 MetTyrValGlyGlyValGluHisArgLeuGluAlaAlaCysAsnTrpThrArgGlyGlu
 GGATGTACGTGGGAGGGGTCGAGCACAGGCTGGAAGCTGCCTGCAACTGGACGCGGGGCG
 CCTACATGCACCTCCCCAGCTCGTGTCCGACCTTCGACGGACGTTGACCTGCGCCCCGC

421 -----Overlap with 12f-----
 ArgCysAspLeuGluAspArgAspArgSerGluLeuSerProLeuLeuLeuThrThrThr
 AACGTTGCGATCTGGAAGACAGGGACAGGTCCGAGCTCAGCCCGTTACTGCTGACCACTA
 TTGCAACGCTAGACCTTCTGTCCCTGTCCAGGCTCGAGTCGGGCAATGACGACTGGTGAT

481 GlnTrpGlnValLeuProCysSerPheThrThrLeuProAlaLeuSerThrGlyLeu
 CACAGTGGCAGGTCTCCCGTGTTCCTTCACAACCCTGCCAGCCTTGTCCACCGGCCTCA
 GTGTCACCGTCCAGGAGGGCACAAGGAAGTGTGGGACGGTCGGAACAGGTGGCCGGAGT

SUBSTITUTE SHEET

FIG. 6

Translation of DNA CA59a

LeuValMetAlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIleAlaGlyAla
1 TTGGTAATGGCTCAGCTGCTCCGGATCCCACAAGCCATCTTGGACATGATCGCTGGTGCT
AACCATTACCGAGTCGACGAGGCCTAGGGTGTTTCGGTAGAACCTGTACTAGCGACCACGA

HisTrpGlyValLeuAlaGlyIleAlaTyrPheSerMetValGlyAsnTrpAlaLysVal
61 CACTGGGGAGTCCTGGCGGGCATAGCGTATTTCTCCATGGTGGGGAACTGGGCGAAGGTC
GTGACCCCTCAGGACCGCCCGTATCGCATAAAGAGGTACCACCCCTTGACCCGCTTCCAG

LeuValValLeuLeuLeuPheAlaGlyValAspAlaGluThrHisValThrGlyGlySer
121 CTGGTAGTGCTGCTGCTATTGCGCGCGTCGACGCGGAAACCCACGTCACCGGGGGAAGT
GACCATCACGACGACGATAAACGGCCGACGCTGCGCCTTTGGGTGCAGTGGCCCCCTTCA

AlaGlyHisThrValSerGlyPheValSerLeuLeuAlaProGlyAlaLysGlnAsnVal
181 GCCGGCCACAC TGTGTCTGGATTTGTTAGCCTCCTCGCACCAGGCGCCAAGCAGAACGTC
CGGCCGGTGTGACACAGACCTAAACAATCGGAGGAGCGTGGTCCGCGGTTCTGCTTGACG

GlnLeuIleAsnThrAsnGlySerTrpHisLeuAsnSerThrAlaLeuAsnCysAsnAsp
241 CAGCTGATCAACACCAACGGCAGTTGGCACCTCAATAGCACGGCCCTGAACTGCAATGAT
GTCGACTAGTTGTGGTTGCCGTCAACCGTGGAGTTATCGTGCCGGGACTTGACGTTACTA

301 SerLeuAsnThrGlyTrpLeuAlaGlyLeuPheTyrHisHisLysPheAsnSerSerGly
AGCCTCAACACCGGCTGGTTGGCAGGGCTTTTCTATCACCACAAGTTCAACTCTTCAGGC
TCGGAGTTGTGGCCGACCAACCGTCCCGAAAAGATAGTGGTGTTCAGTTGAGAAGTCCG
-----Overlap with 26j-----

-----Overlap with K9-1-----
361 CysProGluArgLeuAlaSerCysArgPro
TGTCCTGAGAGGCTAGCCAGCTGCCGACCCC
ACAGGACTCTCCGATCGGTGACGGCTGGGG

SUBSTITUTE SHEET

7 / 40

Translation of DNA CA84a

FIG. 7

1 GlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArgMetAlaTrpAsp
 CGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATATAACGGGTCACCGCATGGCATGGG
 GCGTTC AACGTTAACGAGATAGATAGGGCCGGTATATTGCCAGTGGCGTACCGTACCC

61 MetMetMetAsnTrpSerProThrThrAlaLeuValMetAlaGlnLeuLeuArgIlePro
 ATATGATGATGAACTGGTCCCCTACGACGGCGTTGGTAATGGCTCAGCTGCTCCGGATCC
 TATACTACTACTTGACCAGGGGATGCTGCCGCAACCATTACCGAGTCGACGAGGCC TAGG

121 GlnAlaIleLeuAspMetIleAlaGlyAlaHisTrpGlyValLeuAlaGlyIleAlaTyr
 CACAAGCCATCTTGGACATGATCGCTGGTGCTCACTGGGGAGTCTTGGCGGGCATAGCGT
 GTGTTCCGGTAGAACCTGTACTAGCGACCACGAGTGACCCCTCAGGACCGCCCGTATCGCA

-----Overlap with CA59a-----
 181 PheSerMetValGlyAsnTrpAlaLysValLeuValValLeuLeuPheAlaGlyVal
 ATTTCTCCATGGTGGGGAAC TGGGCGAAGGTCCTGGTAGTGCTGCTGCTATTTGCCGGCG
 TAAAGAGGTACCACCCCTTGACCCGCTTCCAGGACCATCAGACGACGATAAACGGCCGC

241 AspAlaGluThrHisValThrGly
 TCGACGCGGAAACCCACGTCACCGGGG
 AGCTGCGCCTTTGGGTGCAGTGGCCCC

Translation of DNA CA156e

FIG. 8

1 CysTrpValAlaMetThrProThrValAlaThrArgAspGlyLysLeuProAlaThrGln
 GTGTTGGGTGGCGATGACCCCTACG GTGGCCACCAGGGATGGCAAAC TCCCGCGACGCA
 CACAACCCACCGCTACTGGGGATGCCACCGGTGGTCCCTACCGTTTGAGGGGCGCTGCGT

61 LeuArgArgHisIleAspLeuLeuValGlySerAlaThrLeuCysSerAlaLeuTyrVal
 GCTTCGACGTCACATCGATCTGCTTGTGCGGAGCGCCACCCTCTGTTCCGGCCCTCTACGT
 CGAAGCTGCAGTGTAGCTAGACGAACAGCCCTCGCGGTGGGAGACAAGCCGGGAGATGCA

121 GlyAspLeuCysGlySerValPheLeuValGlyGlnLeuPheThrPheSerProArgArg
 GGGGGACCTATGCGGTCTGTCTTTCTTGTGCGGCCAACTGTTACCTTCTCTCCAGGCG
 CCCCCTGGATACGCCAGACAGAAAGAACAGCQGGTTGACAAGTGGAAGAGAGGGTCCGC

181 HisTrpThrThrGlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArg
 CCACTGGACGACGCAAGGTTGCAATTGCTCTATCTATCCCGGCCATATAACGGGTCACCG
 GGTGACCTGCTGCGTTCCAACGTTAACGAGATAGATAGGGCCGGTATATTGCCAGTGGC

-----Overlap with CA84a-----
 241 MetAlaTrpAspMetMetMetAsnTrpSerProThrThrAlaLeuValValAlaGlnLeu
 CATGGCATGGGATATGATGATGAACTGGTCCCCTACGACGGCGTTGGTAGTGGCTCAGCT
 GTACCGTACCCTATACTACTACTTGACCAGGGGATGCTGCCGCAACCATCACCGAGTCGA

301 LeuArgIleProGlnAla
 GCTCCGGATCCCACAAGCC
 CGAGGCCTAGGGTGTTCGG

SUBSTITUTE SHEET

FIG. 9

Translation of DNA CA167b

SerThrGlyLeuTyrHisValThrAsnAspCysProAsnSerSerIleValTyrGluAla
1 CTCCACGGGGCTTTACCACGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGGC
GAGGTGCCCCGAAATGGTGTCAGTGGTTACTAACGGGATTGAGCTCATAACACATGCTCCG

AlaAspAlaIleLeuHisThrProGlyCysValProCysValArgGluGlyAsnAlaSer
61 GGCCGATGCCATCCTGCACACTCCGGGGTGCGTCCCTTGCCTTCGTGAGGGCAACGCCTC
CCGGCTACGGTAGGACGTGTGAGGCCCCACGCAGGGAACGCAAGCACTCCCGTTGCGGAG

ArgCysTrpValAlaMetThrProThrValAlaThrArgAspGlyLysLeuProAlaThr
121 GAGGTGTTGGGTGGCGATGACCCCTACGGTGGCCACCAGGGATGGCAAACCCCCGCGAC
CTCCACAACCCACCGCTACTGGGGATGCCACCGGTGGTCCCTACCGTTTGAGGGGCGCTG

-----Overlap with CA156e-----
GlnLeuArgArgHisIleAspLeuLeuValGlySerAlaThrLeuCysSerAlaLeuTyr
181 GCAGCTTCGACGTCACATCGATCTGCTTGTCTGGGAGCGCTACCTCTGTTCGGCCCTCTA
CGTCGAAGCTGCAGTGTAGCTAGACGAACAGCCCTCGCGATGGGAGACAAGCCGGGAGAT

ValGlyAspLeuCysGlySerValPheLeu
241 CGTGGGGGACTTGTGCGGGTCTGTCTTTCTTG
GCACCCCTGAACACGCCAGACAGAAAGAAC

SUBSTITUTE SHEET

FIG. 10

Translation of DNA ssCA216a

ArgArgArgSerArgAsnLeuGlyLysValIleAspThrLeuThrCysGlyPheAlaAsp
1 CCCGGCGTAGGTGCGCAATTTGGGTAAGGTCATCGATACCCTTACGTGCGGCTTCGCCG
GGGCCGCATCCAGCGCGTTAAACCCATTCCAGTAGCTATGGGAATGCACGCCGAAGCGGC

LeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyAlaAlaArgAlaLeuAla
61 ACCTCATGGGGTACATACCGCTCGTCGGCGCCCTCTTGGAGGCGCTGCCAGGGCCCTGG
TGGAGTACCCCATGTATGGCGAGCAGCCGCGGGGAGAACCTCCGCGACGGTCCCGGGACC

HisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsnLeuProGlyCys
121 CGCATGGCGTCCGGGTCTTGAAGACGGCGTGAACATGCAACAGGGAACCTTCCTGGTT
CGGTACCGCAGGCCCAAGACCTTCTGCCGCACTTGATACGTTGTCCCTTGAAGGACCAA

SerPheSerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValProAlaSerAlaTyr
181 GCTCTTTCTCTATCTTCTTCTGGCCCTGCTCTCTTGCTTGACTGTGCCCGCTTCGGCCT
CGAGAAAGAGATAGAAGGAAGACCGGGACGAGAGAACGAACCTGACACGGGCGAAGCCGGA

GlnValArgAsnSerThrGlyLeuTyrHisValThrAsnAspCysProAsnSerSerIle
241 ACCAAGTGCGCAACTCCACGGGGCTTTACCACGTCACCAATGATTGCCCTAACTCGAGTA
TGGTTCACGCGTTGAGGTGCCCCGAAATGGTGAGTGTTACTAACGGGATTGAGCTCAT

-----overlap with CA167b-----
ValTyrGluAlaAlaAspAlaIleLeuHisThrProGlyCysValProCysValArgGlu
301 TTGTGTACGAAGCGGGCCGATGCCATCCTGCACACTCCGGGGTGCGTCCCTTGCGTTCTGTG
AACACATGCTTCGCCGGCTACGGTAGGACGTGTGAGGCCCCACGCAGGGAACGCAAGCAC

GlyAsnAlaSerArgCysTrpValAlaMetThrProThrValAla
361 AGGGCAACGCCCTCGAGGTGTTGGGTGGCGATGACCCCTACGGTGGCC
TCCCGTTGCGGAGCTCCACAACCCACCGCTACTGGGGATGCCACCGG

SUBSTITUTE SHEET

FIG. 11

Translation of DNA ssCA290a

LysLysAsnLysArgAsnThrAsnArgArgProGlnAspValLysPheProGlyGlyGly
 1 AAAAAAAAAAACAACGTAACACCAACCGTCGCCCACAGGACGTCAAGTTCCCGGTGGCG
 TTTTTTTTTTGTTCATTGTGGTTGGCAGCGGTGTCCTGCAGTTCAAGGGCCACCGC

GlnIleValGlyGlyValTyrLeuLeuProArgArgGlyProArgLeuGlyValArgAla
 61 GTCAGATCGTTGGTGGAGTTTACTTGTTCGCCGCGAGGGGCCCTAGATTGGGTGTGCGCG
 CAGTCTAGCAACCACCTCAATGAACAACGGCGCTCCCCGGGATCTAACCACACGCGC

ThrArgLysThrSerGluArgSerGlnProArgGlyArgArgGlnProIleProLysAla
 121 CGACGAGAAAGACTTCCGAGCGGTGCGAACCTCGAGGTAGACGCCAGCCTATCCCCAAGG
 GCTGCTCTTTCTGAAGGCTCGCCAGCGTTGGAGCTCCATCTGCGGTGCGATAGGGGTTC

ArgArgProGluGlyArgThrTrpAlaGlnProGlyTyrProTrpProLeuTyrGlyAsn
 181 CTCGTCGGCCCGAGGGCAGGACCTGGGCTCAGCCCGGTACCTTGGCCCCCTCTATGGCA
 GAGCAGCCGGGCTCCCGTCTGGACCCGAGTCGGGCCCATGGGAACCGGGGAGATACCGT

GluGlyCysGlyTrpAlaGlyTrpLeuLeuSerProArgGlySerArgProSerTrpGly
 241 ATGAGGGCTGCGGGTGGGCGGGATGGCTCCTGTCTCCCGTGGCTCTCGGCCTAGCTGGG
 TACTCCCGAAGCCACCCGCCCTACCGAGGACAGAGGGGCACCGAGAGCCGGATCGACCC

ProThrAspProArgArgArgSerArgAsnLeuGlyLysValIleAspThrLeuThrCys
 301 GCCCCACAGACCCCGGCGTAGGTGCGCGCAATTTGGGTAAGGTCATCGATACCCCTACGT
 CGGGGTGTCTGGGGGCCGCATCCAGCGCGTTAAACCATTCAGTAGCTATGGGAATGCA

GlyPheAlaAspLeuMetGlyTyrIleProLeuValGlyAlaProLeuGlyGlyAlaAla
 361 GCGGCTTCGCCGACCTCATGGGGTACATACCGCTCGTCGGCGCCCCCTCTGGAGGCGCTG
 CGCCGAAGCGGCTGGAGTACCCCATGTATGGCGAGCAGCCGCGGGGAGAACCTCCGCGAC

ArgAlaLeuAlaHisGlyValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsn
 421 CCAGGGCCCTGGCGCATGGCGTCCGGGTTCTGGAAGACGGCGTGAACCTATGCAACAGGGA
 GGTCCCGGGACCGGTACCGCAGGCCCAAGACCTTCTGCCGCACTTGATACGTTGTCCCT

LeuProGlyCysSerPheSerThrPhe
 481 ACCTTCCTGGTTGCTCTTCTCTACCTTC
 TGAAGGACCAACGAGAAAGAGATGGAAG

SUBSTITUTE SHEET

11 / 40

Translation of DNA ag30a

FIG. 12-1

#MetSerValValGlnProProGlyProProLeu

#MetAlaLeuValOP

- 1 CGCAGAAAGCGTCTAGCCATGGCGTTAGTATGAGTGTCTGTCAGCCTCCAGGACCCCCC
GCGTCTTTTCGAGATCGGTACCGCAATCATACTACAGCACGTCTGGAGGTCTGGGGGGG
ProGlyGluProAM
- 61 TCCCGGGAGAGCCATAGTGGTCTGCGGAACCGGTGAGTACACCGGAATTGCCAGGACGAC
AGGGCCCTCTCGGTATCACCAGACGCTTGGCCACTCATGTGGCCTTAACGGTCTGCTG
- #MetProGlyAspLeuGlyValProProGlnAsp
- 121 CGGGTCCTTTCTTGGATCAACCCGCTCAATGCCTGGAGATTGGGGCGTCCCCCGCAAGA
GCCCAGGAAAGAACCTAGTTGGGCGAGTTACGGACCTCTAAACCCGCACGGGGGCGTTCT
CysAM OP AM GlyAlaCys
*
- 181 CTGCTAGCCGAGTAGTGTGGGTGCGGAAAGGCCTTGTGGTACTGCCTGATAGGGTGCTT
GACGATCGGCTCATCACACCAGCGCTTTCGGAACACCATGACGGACTATCCACGAA
GluCysProGlyArgSerArgArgProCysThrMetSerThrAsnProLysProGlnLys
- 241 GCGAGTGCCCCGGGAGGTCTCGTAGACCGTGCACCATGAGCACGAATCCTAAACCTCAAA
CGCTCACGGGGCCCTCCAGAGCATCTGGCACGTGGTACTCGTGCTTAGGATTTGGAGTTT
LysAsnLysArgAsnThrAsnArgArgProGlnAspValLysPheProGlyGlyGlyGln
- 301 -----
AAAAAAACAAACGTAACACCAACCGTGGCCACAGGACGTCAAGTTCCCGGTGGCGGTC
TTTTTTTGTGTTGATTGTGGTTGGCAGCGGGTGTCTTGCAGTTCAAGGGCCACCGCCAG
IleValGlyGlyValTyrLeuLeuProArgArgGlyProArgLeuGlyValArgAlaThr
- 361 -----
AGATCGTTGGTGGAGTTTACTTGTGCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCGA
TCTAGCAACCACCTCAAATGAACAACGGCGCGTCCCCGGGATCTAACCACACGCGCGCT
ArgLysThrSerGluArgSerGlnProArgGlyArgArgGlnProIleProLysAlaArg
- 421 -----
CGAGAAAGACTTCCGAGCGGTGCAACCTCGAGGTAGACGTACGCTATCCCCAAGGCTC
GCTCTTTCTGAAGGCTCGCCAGCGTTGGAGCTCCATCTGCAGTCGGATAGGGGTCCGAG
ArgProGluGlyArgThrTrpAlaGlnProGlyTyrProTrpProLeuTyrGlyAsnGlu
- 481 -----overlap with CA290a-----
GTCGGCCCCGAGGGCAGGACCTGGGCTCAGCCCGGGTACCCTTGGCCCTCTATGGCAATG
CAGCCGGGCTCCCGTCTGGACCCGAGTCGGGCCCCATGGGAACGGGGAGATACCGTTAC
GlyCysGlyTrpAlaGlyTrpLeuLeuSerProArgGlySerArgProSerTrpGlyPro
- 541 -----
AGGGCTGCGGGTGGGCGGGATGGCTCCTGTCTCCCGTGGCTCTCGGCCTAGCTGGGGCC
TCCCGACGCCACCCGCCCTACCGAGGACAGAGGGGCACCGAGAGCCGGATCGACCCCGG
ThrAspProArgArgArgSerArgAsnLeuGlyLysValIleAspThrLeuThrCysGly

SUBSTITUTE SHEET

12 / 40

601 CCACAGACCCCGGCGTAGGTCGCGCAATTTGGGTAAGGTCATCGATACCCTTACGTGCG
GGTGTCTGGGGGCCGCATCCAGCGCGTTAAACCCATTCCAGTAGCTATGGGAATGCACGC

Phe

661 GCTTC
CGAAG

* = Start of long HCV ORF
| = Putative first amino acid of large HCV polyprotein
= Putative small encoded peptides (that may play a
translational regulatory role)

FIG. 12-2

SUBSTITUTE SHEET

13/40

FIG. 13

Translation of DNA CA205a

1 ValLeuGlyArgGluArgProCysGlyThrAlaOP AM GlyAlaCysGluCysProGly
 GTCTTGGGTCGCGAAAGGCCTTGTGGTACTGCCTGATAGGGTGCTTGCAGAGTGCCCCGGG
 CAGAACCCAGCGCTTCCGGAACACCATGACGGACTATCCCA CGAACGCTCACGGGGCCC

*

61 ArgSerArgArgProCysThrMetSerThrAsnProLysProGlnArgLysThrLysArg
 AGGTCTCGTAGACCGTGCACCATGAGCACGAATCCTAAACCTCAAAGAAAAACCAAACGT
 TCCAGAGCATCTGGCAGCTGGTACTCGTGCTTAGGATTGGAGTTTCTTTTGGTTTGCA

121 AsnThrAsnArgArgProGlnAspValLysPheProGlyGlyGlyGlnIleValGlyGly
 AACACCAACCGTCGCCCCACAGGACGTCAAGTTCCCGGGTGGCGGTTCAGATCGTTGGTGGA
 TTGTGGTTGGCAGCGGGTGTCCTGCAGTTCAAGGGCCCACCGCCAGTCTAGCAACCACCT

181 ValTyrLeuLeuProArgArgGlyProArgLeuGlyValArgAlaThrArgLysThrSer
 GTTTACTTGTGTCGCGCAGGGGCCCTAGATTGGGTGTGCGCGCGACGAGAAAGACTTCC
 CAAATGAACAACGGCGCGTCCCGGGATCTAACCACACGCGCGCTGCTCTTCTGAAGG

-----overlap with CA290a-----

241 GluArgSerGlnProArgGlyArgArgGlnProIleProLysAlaArgArgProGluGly
 GAGCGGTGCGAACCTCGAGGTAGACGTCAGCCTATCCCCAAGGCTCGTCGGCCCCGAGGGC
 CTCGCCAGCGTTGGAGCTCCATCTGCAGTCGGATAGGGGTTCGAGCAGCCGGGCTCCCG

301 ArgThrTrpAlaGlnProGlyTyrProTrpProLeuTyrGlyAsnGluGlyCys
 AGGACCTGGGCTCAGCCCGGTACCTTGGCCCCCTCTATGGCAATGAGGGCTGCG
 TCCTGGACCCGAGTCGGGCCCATGGGAACCGGGAGATACCGTTACTCCCGACGC

* = putative initiator methionine codon

SUBSTITUTE SHEET.

14 / 40

FIG. 14

Translation of DNA 18g

```

#ProProOP
#SerThrMetAsnHisSerProValArgAsnTyrCysLeuHisAlaGluSerValAM Pro
#LeuHisHisGluSerLeuProCysGluGluLeuLeuSerSerArgArgLysArgLeuAla
1 CTCCACCATGAATCACTCCCCTGTGAGGAAGTACTGTCTTCACGCAGAAAGCGTCTAGCC
GAGGTGGTACTTAGTGAGGGGACACTCCTTGATGACAGAAGTGCCTCTTTCGCAGATCGG

-----
#MetSerValValGlnProProGlyProProLeuProGlyGluProAM
MetAlaLeuValOP
61 ATGGCGTTAGTATGAGTGTCTGTCAGCCTC CAGGACCCCCCTCCCGGGAGAGCCATAGT
TACCGCAATCATACTCACAGCACGTGGGAGGTCTTGGGGGGAGGGCCCTCTCGGTATCA

-----
121 GGTCTGCGGAACCGGTGAGTACACCGGAATTGCCAGGACGACCGGGTCCTTTCTTGGATC
CCAGACGCCTTGCCCACTCATGTGGCCTTAACGGTCCTGCTGGCCCAGGAAAGAACCTAG

-----overlap with ag30a-----
#MetProGlyAspLeuGlyValProProGlnAspCysAM
181 AACCCGCTCAATGCCTGGAGATTTGGGCGTGGCCCCGCAAGACTGCTAGCCGAGTAGTGT
TTGGGCGAGTTACGGACCTCTAAACCCGCACGGGGGCGTTCTGACGATCGGCTCATCACA

-----
OP AM GlyAlaCysGluCysProGlyArgSer
*
241 TGGGTCGCGAAAGGCCTTGTGGTACTGCCTGATAGGGTGCTTGCGAGTGCCCCGGGAGGT
ACCCAGCGCTTTCCGGAACACCATGACGGACTATCCACGAACGCTCACGGGGCCCTCCA

-----
ArgArg
301 CTCGTAGA
GAGCATCT

```

* = Start of long HCV ORF
 # = Putative small encoded peptides (that may play a translational regulatory role)

SUBSTITUTE SHEET

FIG. 15

Translation of DNA 16jh

-----Overlap with 15e -----
1 GlyAlaCysTyrSerIleGluProLeuAspLeuProProIleIleGlnArgLeuHisGly
GGGGCCTGCTACTCCATAGAACCACTGGATCTACCTCCAATCATTCAAAGACTCCATGGC
CCCCGGACGATGAGGTATCTTGGTGACCTAGATGGAGGTTAGTAAGTTTCTGAGGTACCG
61 LeuSerAlaPheSerLeuHisSerTyrSerProGlyGluIleAsnArgValAlaAlaCys
CTCAGCGCATTTTCACTCCACAGTTACTCTCCAGGTGAAATTAATAGGGTGGCCGCATGC
GAGTCGCGTAAAAGTGAGGTGTCAATGAGAGGTCCACTTTAATTATCCCACCGGCGTACG
Gly*
G
121 LeuArgLysLeuGlyValProProLeuArgAlaTrpArgHisArgAlaArgSerValArg
CTCAGAAAACCTTGGGGTACCGCCCTTGCGAGCTTGGAGACACCGGGCCCGGAGCGTCCGC
GAGTCTTTTGAACCCCATGGGCGGAACGCTCGAACCTCTGTGGCCCGGGCCTCGCAGGCG
181 AlaArgLeuLeuAlaArgGlyGlyArgAlaAlaIleCysGlyLysTyrLeuPheAsnTrp
GCTAGGCTTCTGGCCAGAGGAGGCAGGGCTGCCATATGTGGCAAGTACCTCTTCAACTGG
CGATCCGAAGACCGGTCTCCTCCGTCCCACGGTATACACCGTTCATGGAGAAGTTGACC
241 AlaValArgThrLysLeuLys
GCAGTAAGAACAAGCTCAAAC
CGTCATTCTTGTTCGAGTTTG

* = nucleotide heterogeneity

SUBSTITUTE SHEET

16 / 40

COMBINED ORF OF DNAs p114a THROUGH 15e

FIG. 16-1

(p114a/CA167b/CA156e/CA84a/CA59a/K9-1/12f/14i/11b/7f/7e/
8h/33c/40b/37b/35/36/81/32/33b/25c/14c/8f/33f/33g/39c/
35f/19g/26g & 15e)

1 ArgSerArgAsnLeuGlyLysValIleAspThrLeuThrCysGlyPheAlaAspLeuMet
AGGTCGCGCAATTTGGGTAAGGTCAATCGATACCCCTACGTGCGGCTTCGCGGACCTCATG
TCCAGCGCGTTAAACCCATTCCAGTAGCTATGGGAATGCACGCCGAAGCGGCTGGAGTAC

61 GlyTyrIleProLeuValGlyAlaProLeuGlyGlyAlaAlaArgAlaLeuAlaHisGly
GGGTACATACCGCTCGTGGGCGCCCTCTTGGAGGCGCTGCCAGGCCCTGGCGCATGGC
CCCATGTATGGCGAGCAGCCGCGGGGAGAACCTCCGCGACGGTCCCGGGACCGGTACCG

121 ValArgValLeuGluAspGlyValAsnTyrAlaThrGlyAsnLeuProGlyCysSerPhe
GTCCGGGTTCTGGAAGACGGCTGAACATATGCAACAGGGAACCTTCCTGGTTGCTCTTTC
CAGGCCCAAGACCTTCTGCGCACTTGATACGTTGTCCCTTGAAGGACCAACGAGAAAG

181 SerIlePheLeuLeuAlaLeuLeuSerCysLeuThrValProAlaSerAlaTyrGlnVal
TCTATCTTCTTCTGGCCCTGCTCTCTTGTGACTGTGCGCGCTTCGGCCTACCAAGTG
AGATAGAAGGAAGACCGGACGAGAGAAACGAACGTACACGGGCGAAGCCGGATGGTTCAC

241 ArgAsnSerThrGlyLeuTyrHisValThrAsnAspCysProAsnSerSerIleValTyr
CGCAACTCCACGGGGCTTTACCACGTACCAATGATGCCCCTAACTCGAGTATTGTGTAC
GCGTTGAGGTGCCCCGAAATGGTGAGTGGTTACTAACGGGATTGAGCTCATAACACATG

301 GluAlaAlaAspAlaIleLeuHisThrProGlyCysValProCysValArgGluGlyAsn
GAGGCGCCGATGCCATCCTGCACACTCCGGGGTGGTCCCTTGGCTGCTGAGGGCAAC
CTCCGCGCGCTACGGTAGGACGTGTGAGGCCCCACGCAGGGAACGCAAGCACTCCCGTTG

361 AlaSerArgCysTrpValAlaMetThrProThrValAlaThrArgAspGlyLysLeuPro
GCCTCGAGGTGTTGGGTGGCGATGACCCCTACGGTGGCCACCAGGGATGGCAAACTCCCC
CGGAGCTCCACAACCCACCGCTACTGGGATGCCACCGGTGGTCCCTACCGTTTGGGGG

421 AlaThrGlnLeuArgArgHisIleAspLeuLeuValGlySerAlaThrLeuCysSerAla
GCGACGCAGCTTCGACGTACATCGATCTGCTGTGCGGGAGCGCCACCTCTGTTTCGGCC
CGCTGCGTCGAAGCTGCAGTGTAGCTAGACGAACAGCCCTCGCGGTGGGAGACAAGCCGG

481 LeuTyrValGlyAspLeuCysGlySerValPheLeuValGlyGlnLeuPheThrPheSer
CTCTACGTGGGGGACCTATGCGGGTCTGTCTTCTTGTGCGGCCAACTGTTCACCTTCTCT
GAGATGCACCCCTGGATACGCCAGACAGAAAGAACAGCCGTTGACAAGTGAAGAGA

541 ProArgArgHisTrpThrThrGlnGlyCysAsnCysSerIleTyrProGlyHisIleThr
CCCAGGCCCACTGGACGACGCAAGTTGCAATTGCTCTATCTATCCCGGCCATATAACG
GGGTCCGCGTGACCTGCTGCGTTCCAACGTTAACGAGATAGTAGGGCCGGTATATTGC

601 GlyHisArgMetAlaTrpAspMetMetMetAsnTrpSerProThrThrAlaLeuValMet
GGTCACCGCATGGCATGGGATATGATGATGAACGTGCCCTACGACGGCGTTGGTAATG
CCAGTGGCGTACCGTACCTATACTACTTGTACAGGGGATGCTGCCGCAACCATTAC

661 AlaGlnLeuLeuArgIleProGlnAlaIleLeuAspMetIleAlaGlyAlaHisTrpGly
GCTCAGCTGCTCCGGATCCCAAGCCATCTTGGACATGATCGCTGGTGTCACTGGGGA
CGAGTGCAGAGGCCTAGGGTGTTCGGTAGAACCTGTACTAGCGACCACGAGTGACCCCT

721 ValLeuAlaGlyIleAlaTyrPheSerMetValGlyAsnTrpAlaLysValLeuValVal
GTCTGGCGGGCATAGCGTATTTCTCCATGGTGGGGAAGTGGGGAAGTCTGGTAGTG
CAGGACCGCCCGTATCGCATAAAGAGGTACCACCCCTTGACCGCTTCCAGGACCATCAC

781 LeuLeuLeuPheAlaGlyValAspAlaGluThrHisValThrGlyGlySerAlaGlyHis
CTGCTGCTATTTGCCGGCGTGCAGCGGAAACCCACGTACCGGGGGAAGTGCCGGCCAC
GACGACGATAAACGGCCGACGTGCGCCTTTGGGTGCAGTGGCCCCCTTACCGCCGGTG

SUBSTITUTE SHEET

17 / 40

FIG. 16-2

841 ThrValSerGlyPheValSerLeuLeuAlaProGlyAlaLysGlnAsnValGlnLeuIle
 ACTGTGTCTGGATTGTGTAGCCTCCTCGCACCAGGCGCAAGCAGAACGTCCAGCTGATC
 TGACACAGACCTAAACAATCGGAGGAGCGTGGTCCGCGGTTCTGCTTGCAGGTGCGACTAG

901 AsnThrAsnGlySerTrpHisLeuAsnSerThrAlaLeuAsnCysAsnAspSerLeuAsn
 AACACCAACGGCAGTTGGCACCTCAATAGCACGGCCCTGAAGTGAATGATAGCCTCAAC
 TTGTGGTTGCCGTCAACCGTGGAGTTATCGTGCCGGGACTTGACGTTACTATCGGAGTTG

961 ThrGlyTrpLeuAlaGlyLeuPheTyrHisHisLysPheAsnSerSerGlyCysProGlu
 ACCGGCTGGTTGGCAGGGCTTTTCTATCACCACAAGTTCAACTCTTCAGGCTGTCTGAG
 TGGCCGACCAACCGTCCCGAAAAGATAGTGGTGTCAAGTTGAGAAGTCCGACAGGACTC

1021 ArgLeuAlaSerCysArgProLeuThrAspPheAspGlnGlyTrpGlyProIleSerTyr
 AGGCTAGCCAGCTGCCGACCCCTTACCGATTTTGACCAGGGCTGGGGCCCTATCAGTTAT
 TCCGATCGGTTCGACGGCTGGGGAATGGCTAAAACTGGTCCCGACCCCGGGATAGTCAATA

1081 AlaAsnGlySerGlyProAspGlnArgProTyrCysTrpHisTyrProProLysProCys
 GCCAACGGAAGCGGCCCCGACAGCGCCCTACTGCTGGCACTACCCCCAAAACCTTGC
 CGGTGCTTCGCCGGGGCTGGTCCGGGGATGACACCGTGATGGGGGGTTTTGGAACG

1141 GlyIleValProAlaLysSerValCysGlyProValTyrCysPheThrProSerProVal
 GGTATTGTGCCCCGAAGAGTGTGTGGTCCGGTATATTGCTTCACTCCAGCCCCGTG
 CCATAACACGGGCGTTCTCACACACACAGGCCATATAACGAAGTGAGGGTGGGGCAC

1201 ValValGlyThrThrAspArgSerGlyAlaProThrTyrSerTrpGlyGluAsnAspThr
 GTGGTGGGAACGACGACAGGTGCGGGCGCGCCACCTACAGTGGGGTGAAAATGATACG
 CACCACCCTTGCTGGCTGTCCAGCCCGCGGGGTGGATGTCGACCCCACTTTTACTATGC

1261 AspValPheValLeuAsnAsnThrArgProProLeuGlyAsnTrpPheGlyCysThrTrp
 GACGTCTTCGTCTTAACAATACAGGCCACCGCTGGGCAATTGGTTTCGGTTGTACCTGG
 CTGCAGAAGCAGGAATTGTTATGGTCCGGTGGCGACCCGTTAACAAGCCAACATGGACC

1321 MetAsnSerThrGlyPheThrLysValCysGlyAlaProProCysValIleGlyGlyAla
 ATGAACCAACTGGATTACCAAAGTGTGCGGAGCGCCTCCTTGTGTATCGGAGGGGGCG
 TACTTGAGTTGACCTAAGTGGTTTCACACGCCTCGCGGAGGAACACAGTAGCCTCCCCGC

1381 GlyAsnAsnThrLeuHisCysProThrAspCysPheArgLysHisProAspAlaThrTyr
 GGCAACAACACCCTGCCTGCCCCACTGATTGCTTCCGCAAGCATCCGGACGCCACATAC
 CCGTTGTTGTGGGACGTGACGGGGTACTAACGAAGGCGTTCTGAGGCTCGCGGTGTATG

1441 SerArgCysGlySerGlyProTrpIleThrProArgCysLeuValAspTyrProTyrArg
 TCTCGGTGCGGCTCCGGTCCCTGGATCACACCCAGGTGCTGGTTCGACTACCGGTATAGG
 AGAGCCACGCCGAGGCCAGGACCTAGTGTGGGTCCACGGACAGCTGATGGGCATATCC

1501 LeuTrpHisTyrProCysThrIleAsnTyrThrIlePheLysIleArgMetTyrValGly
 CTTTGGCATTATCCTTGTTACCATCAACTACCATATTTAAATCAGGATGTACGTGGGA
 GAAACCGTAATAGGAACATGGTAGTTGATGTGGTATAAATTTAGTCTACATGCACCCCT

1561 GlyValGluHisArgLeuGluAlaAlaCysAsnTrpThrArgGlyGluArgCysAspLeu
 GGGGTGGAACACAGGCTGGAAGCTGCCTGCAACTGGACGCGGGGCGAACGTTGCGATCTG
 CCCAGCTTGTGTCCGACCTTCGACGGACGTGACCTGCGCCCCGCTTGCAACGCTAGAC

1621 GluAspArgAspArgSerGluLeuSerProLeuLeuLeuThrThrThrGlnTrpGlnVal
 GAAGACAGGGACAGGTCCGAGCTCAGCCCGTTACTGCTGACCACTACACAGTGGCAGGTC
 CTCTGTCCCTGTCCAGGCTCGAGTCCGGCAATGACGACTGGTGATGTGTACCGTCCAG

1681 LeuProCysSerPheThrThrLeuProAlaLeuSerThrGlyLeuIleHisLeuHisGln
 CTCCCGTGTTCCTTCAACAACCTACCAGCCTTGTCACCGGCCCTCATCCACCTCCACCAG
 GAGGGCACAAAGGAAGTGTGGGATGGTCGGAACAGGTGGCCGAGTAGGTGGAGGTGGTC

1741 AsnIleValAspValGlnTyrLeuTyrGlyValGlySerSerIleAlaSerTrpAlaIle
 AACATTGTGGACGTGCAGTACTGTACGGGGTGGGGTCAAGCATCGCGTCTGGGCCATT
 TTGTAACACCTGCACGTATGAACATGCCCAACCCAGTTCGTAGCGCAGGACCCGGTAA

1801 LysTrpGluTyrValValLeuLeuPheLeuLeuAlaAspAlaArgValCysSerCys
 AAGTGGGAGTACGTGTTCTCTCTCTGCTTGCAGACGCGCGCTCTGCTCCTGC

SUBSTITUTE SHEET

18/40

FIG. 16-3

TTCACCCCTCATGCAGCAAGAGGACAAGGAAGACGAACGTCTGCGCGCGCAGACGAGGACG
1861 LeuTrpMetMetLeuLeuIleSerGlnAlaGluAlaAlaLeuGluAsnLeuValIleLeu
TTGTGGATGATGCTACTCATATCCCAAGCGGAGGCGGCTTTGGAGAACCTCGTAATACTT
AACACCTACTACGATGAGTATAGGGTTTCGCTCCGCCGAAACCTCTTGAGCATTATGAA
1921 AsnAlaAlaSerLeuAlaGlyThrHisGlyLeuValSerPheLeuValPhePheCysPhe
AATGCAGCATCCCTGGCCGGGACGCACGGTCTGTATCCTTCCTCGTGTCTTCTGCTTT
TTACGTCGTAGGGACCGGCCCTGCGTGCCAGAACATAGGAAGGAGCACAAGAAGACGAAA
1981 AlaTrpTyrLeuLysGlyLysTrpValProGlyAlaValTyrThrPheTyrGlyMetTrp
GCATGGTATTGAAGGGTAAGTGGGTGCCCGGAGCGGTCTACACCTTCTACGGGATGTGG
CGTACCATAAACTTCCCATTCACCCACGGGCTCGCCAGATGTGAAGATGCCCTACACC
2041 ProLeuLeuLeuLeuLeuAlaLeuProGlnArgAlaTyrAlaLeuAspThrGluVal
CCTCTCCTCCTGCTCCTGTTGGCGTTGCCCGAGCGGGCGTACGCGCTGGACACGGAGGTG
GGAGAGGAGGACGAGGACAACCGCAACGGGGTCCGCCGATGCGCGACCTGTGCCTCCAC
2101 AlaAlaSerCysGlyGlyValValLeuValGlyLeuMetAlaLeuThrLeuSerProTyr
GCCGCGTCGTGTGGCGGTGTGTCTCGTGGGTGATGGCGCTGACTCTGTCACCATAT
CGGCGCAGCACACCGCCACAACAAGAGCAGCCCACTACCGGACTGAGACAGTGGTATA
2161 TyrLysArgTyrIleSerTrpCysLeuTrpTrpLeuGlnTyrPheLeuThrArgValGlu
TACAAGCGCTATATCAGCTGGTGTGTCTCGTGGGTTCAGTATTTTCTGACCAGATGGAA
ATGTTCCGATATAGTCGACCACGAACACCACCGAAGTCATAAAAGACTGGTCTCACCTT
2221 AlaGlnLeuHisValTrpIleProProLeuAsnValArgGlyGlyArgAspAlaValIle
GCGCAACTGCACGTGTGGATTCCCCCCTCAACGTCCGAGGGGGGCGGACGCCGTATC
CGCGTTGACGTGCACACCTAAGGGGGGAGTTGCAGGCTCCCCCGCGCTGCGGCAGTAG
2281 LeuLeuMetCysAlaValHisProThrLeuValPheAspIleThrLysLeuLeuAla
TTACTCATGTGTGTGTACACCCGACTCTGGTATTTGACATCACCAAATGTCTGTGGCC
AATGAGTACACACGACATGTGGGCTGAGACCATAAACTGTAGTGGTTTACGACGACCGG
2341 ValPheGlyProLeuTrpIleLeuGlnAlaSerLeuLeuLysValProTyrPheValArg
GTCTTCGGACCCCTTTGATTCTTCAAGCCAGTTTGCTTAAAGTACCCTACTTTGTGCGC
CAGAAGCCTGGGGAAACCTAAGAAGTTCGGTCAAACGAATTTTCATGGGATGAAACACGCG
2401 ValGlnGlyLeuLeuArgPheCysAlaLeuAlaArgLysMetIleGlyGlyHisTyrVal
GTCCAAGGCCTTCTCCGTTCTGCGGTTAGCGCGGAAGATGATCGGAGGCCATTACGTG
CAGGTTCCGGAAGAGGCCAAGACGCGCAATCGCGCCTTCTACTAGCCTCCGGTAATGCAC
2461 GlnMetValIleIleLysLeuGlyAlaLeuThrGlyThrTyrValTyrAsnHisLeuThr
CAAATGGTCATCATTAAGTTAGGGGCGCTTACTGGCACCTATGTTTATAACCATCTCACT
GTTTACCAGTAGTAATTCAATCCCCGGAATGACCGTGGATACAAATATTGGTAGAGTGA
2521 ProLeuArgAspTrpAlaHisAsnGlyLeuArgAspLeuAlaValAlaValGluProVal
CCTCTTCGGGACTGGGCGCACACGGCTTGCGAGATCTGGCGGTGGCTGTAGAGCCAGTC
GGAGAAGCCCTGACCCGCGTGTGCGGAACGCTCTAGACCGGACCGACATCTCGGTGAG
2581 ValPheSerGlnMetGluThrLysLeuIleThrTrpGlyAlaAspThrAlaAlaCysGly
GTCTTCTCCCAAATGGAGACCAAGCTCATCAGTGGGGGGCAGATACCGCCGCGTGGGT
CAGAAGAGGGTTTACCTCTGTTTCGAGTAGTGCACCCCGCTATGGCGGCGCACGCCA
2641 AspIleIleAsnGlyLeuProValSerAlaArgArgGlyArgGluIleLeuLeuGlyPro
GACATCATCAACGGCTTGCTGTTTCCGCCCCGAGGGGCGGGAGATACTGCTCGGGCCA
CTGTAGTAGTTGCCGAACGGACAAGGGCGGCGTCCCCGGCCCTCTATGACGAGCCCGGT
2701 AlaAspGlyMetValSerLysGlyTrpArgLeuLeuAlaProIleThrAlaTyrAlaGln
GCCGATGGAATGGTCTCCAAGGGTGGAGGTTGCTGGCGCCCATCACGGCGTACGCCAG
CGGCTACCTTACCAGAGGTTCCCCACCTCCAACGACCGGGGTAGTGCAGCATGCGGGTC
2761 GlnThrArgGlyLeuLeuGlyCysIleIleThrSerLeuThrGlyArgAspLysAsnGln
CAGACAAGGGGCTCCTAGGCTGCATAATCACCAGCCTAACTGGCCGGGACAAAAACCAA
GTCTGTTCCCGGAGGATCCACGTATTAGTGGTTCGATTGACCGGCCCTGTTTTGGTT

SUBSTITUTE SHEET

19/40
FIG. 16-4

2821 ValGluGlyGluValGlnIleValSerThrAlaAlaGlnThrPheLeuAlaThrCysIle
GTGGAGGGTGAGGTCCAGATTGTGTCAACTGCTGCCCAAACCTTCCTGGCAACGTGCATC
CACCTCCCACTCCAGGTCTAACACAGTTGACGACGGGTTTGAAGGACCGTTGCACGTAG

2881 AsnGlyValCysTrpThrValTyrHisGlyAlaGlyThrArgThrIleAlaSerProLys
AATGGGGTGTGCTGACTGTCTACACGGGGCCGGAACGAGGACCATCGCGTCACCCAAG
TTACCCACACGACCTGACAGATGGTGCCCCGGCCTTGCTCCTGGTAGCGCAGTGGGTTT

2941 GlyProValIleGlnMetTyrThrAsnValAspGlnAspLeuValGlyTrpProAlaPro
GGTCCTGTTCATCCAGATGTATACCAATGTAGACCAAGACCTTGTGGGCTGGCCCGCTCCG
CCAGGACAGTAGGTCTACATATGGTTACATCTGGTTCTGGAACACCCGACCGGGCGAGGC

3001 GlnGlySerArgSerLeuThrProCysThrCysGlySerSerAspLeuTyrLeuValThr
CAAGGTAGCCGCTCATTGACACCTGCACTTGGCGCTCCTCGGACCTTTACCTGGTCAAG
GTTCCATCGGCGAGTAACTGTGGGACGTGAACGCCGAGGAGCCTGGAAATGGACCAGTGC

3061 ArgHisAlaAspValIleProValArgArgArgGlyAspSerArgGlySerLeuLeuSer
AGGCACGCCGATGTTCATCCCGTGGCGCGGGGTGATAGCAGGGGCGAGCTGCTGTGCG
TCCGTGGCGCTACAGTAAGGGCACGCGGCCGCCCACTATCGTCCCCGTGGACGACAGC

3121 ProArgProIleSerTyrLeuLysGlySerSerGlyGlyProLeuLeuCysProAlaGly
CCCCGGCCCATTTCTACTGTAAAGGCTCCTCGGGGGTCCGCTGTTGTGCCCGCGGGG
GGGGCCGGGTAAAGGATGAACCTTCCGAGGAGCCCCCAGGCGACAACACGGGGCGCCCC

3181 HisAlaValGlyIlePheArgAlaAlaValCysThrArgGlyValAlaLysAlaValAsp
CAGCCGTGGGCATATTTAGGGCCGCGGTGTGCACCCGTGGAGTGGCTAAGCGGTGGAC
GTGCGGCACCCGTATAAATCCCGGCGCCACAGTGGGCACCTCACCGATTCCGCCACCTG

3241 PheIleProValGluAsnLeuGluThrThrMetArgSerProValPheThrAspAsnSer
TTTATCCCTGTGGAGAACCTAGAGACAACCATGAGGTCCCGGTGTTACGGATAACTCC
AAATAGGGACACCTCTGGATCTCTGTGGTACTCCAGGGGGCCACAAGTCCCTATTGAGG

3301 SerProProValValProGlnSerPheGlnValAlaHisLeuHisAlaProThrGlySer
TCTCCACCACTAGTGTCCCGAGAGCTTCCAGGTGGCTCACCTCCATGCTCCACAGGCAGC
AGAGGTGGTTCATCAGGGGTCTCGAAGGTCCACCGAGTGGAGGTACGAGGGTGTCCGTGC

3361 GlyLysSerThrLysValProAlaAlaTyrAlaAlaGlnGlyTyrLysValLeuValLeu
GGCAAAGCACCAAGGTCCCGGCTGCATATGCAGCTCAGGGCTATAAGGTGCTAGTACTC
CCGTTTTCTGTGTTCCAGGGCCGACGTATACGTGAGTCCCGATATCCACGATCATGAG

3421 AsnProSerValAlaAlaThrLeuGlyPheGlyAlaTyrMetSerLysAlaHisGlyIle
AACCCCTCTGTTGCTGCAACACTGGGCTTGGTGCTTACATGTCCAAGGCTCATGGGATC
TTGGGGAGACAAAGACGTTGTGACCCGAAACACGAATGTACAGGTCCCGAGTACCCTAG

3481 AspProAsnIleArgThrGlyValArgThrIleThrThrGlySerProIleThrTyrSer
GATCCTAACATCAGGACCGGGTGAGAACAATTACCACTGGCAGCCCCATCACGTACTCC
CTAGGATTGTAGTCTGCGCCCACTCTGTGTTAATGGTGACCGTGGGGTAGTGCATGAGG

3541 ThrTyrGlyLysPheLeuAlaAspGlyGlyCysSerGlyGlyAlaTyrAspIleIleIle
ACCTACGGCAAGTTCTTGGCGACGGCGGGTGTCTGGGGGGCGCTTATGACATAATAATT
TGGATGCCGTTCAAGGAACGGCTGCCGCCACGAGCCCCCGGAATACTGTATTATTAA

3601 CysAspGluCysHisSerThrAspAlaThrSerIleLeuGlyIleGlyThrValLeuAsp
TGTGACGAGTGCCACTCCACGGATGCCACATCCATCTTGGGCATCGGCCTGTCTTGGAC
ACACTGCTCAGGTGAGGTGCTACGGTGTAGGTAGAACCGTAGCCGTGACAGGAAGTGC

3661 GlnAlaGluThrAlaGlyAlaArgLeuValValLeuAlaThrAlaThrProProGlySer
CAAGCAGAGACTGCGGGGGCGAGACTGGTTGTGCTCGCCACCGCCACCCCTCGGGGCTCC
GTTCTGTTCTGACGCCCCGCTCTGACCAACACGAGCGGTGGCGGTGGGGAGGCCCGAGG

3721 ValThrValProHisProAsnIleGluGluValAlaLeuSerThrThrGlyGluIlePro
GTCACGTGCCCCATCCCAACATCGAGGAGGTGCTCTGTCCACCACCGGAGAGATCCCT
CAGTGACACGGGGTAGGGTTGTAGCTCTCCAAACGAGACAGGTGGTGGCCTCTCTAGGGA

3781 PheTyrGlyLysAlaIleProLeuGluValIleLysGlyGlyArgHisLeuIlePheCys
TTTTACGGCAAGGCTATCCCCCTCGAAGTAATCAAGGGGGGAGACATCTCATCTTCTGT

SUBSTITUTE SHEET

20/40

FIG. 16-5

AAAATGCCGTTCCGATAGGGGAGCTTCATTAGTTCCTCCCCCTCTGTAGAGTAGAAGACA

3841 HisSerLysLysLysCysAspGluLeuAlaAlaLysLeuValAlaLeuGlyIleAsnAla
CATTCAAAGAAGAAGTGCAGCAACTCGCCGCAAAGCTGGTTCGCATTGGGCATCAATGCC
GTAAGTTTCTTCTTCACGCTGCTTGAGCGCGTTTCGACCAGCGTAACCGTAGTTACGG

3901 ValAlaTyrTyrArgGlyLeuAspValSerValIleProThrSerGlyAspValValVal
GTGGCCTACTACCGCGGTCTTGACGTGTCCGTCATCCCGACCAGCGCGATGTTGTCGTC
CACCGGATGATGGCGCCAGAACTGCACAGGCAGTAGGGCTGGTTCGCGCTACAACAGCAG

3961 ValAlaThrAspAlaLeuMetThrGlyTyrThrGlyAspPheAspSerValIleAspCys
GTGGCAACCGATGCCCTCATGACCGGCTATACCGCGGACTTCGACTCGGTGATAGACTGC
CACCGTTGGCTACGGGAGTACTGGCCGATATGCCCGCTGAAGCTGAGCCACTATCTGACG

4021 AsnThrCysValThrGlnThrValAspPheSerLeuAspProThrPheThrIleGluThr
AATACGTGTGTACCCAGACAGTCGATTTACGCCTTGACCCTACCTTACCATTGAGACA
TTATGCACACAGTGGGTCTGTACGCTAAAGTCGGAAGTGGGATGGAAGTGGTAACCTCTGT

4081 IleThrLeuProGlnAspAlaValSerArgThrGlnArgArgGlyArgThrGlyArgGly
ATCACGCTCCCCAGGATGCTGTCTCCCGCACTCAACGTCGGGCGAGGACTGGCAGGGGG
TAGTGCAGAGGGGGTCTACGACAGAGGGCGTGAGTTGCAGCCCCCTCTGACCGTCCCC

4141 LysProGlyIleTyrArgPheValAlaProGlyGluArgProSerGlyMetPheAspSer
AAGCCAGGCATCTACAGATTTGTGGCACCGGGGAGCGCCCTCCGGCATGTTCCGACTCG
TTCGGTCCGTAGATGTCTAACACCGTGGCCCCCTCGCGGGGAGGCCGTACAAGCTGAGC

4201 SerValLeuCysGluCysTyrAspAlaGlyCysAlaTrpTyrGluLeuThrProAlaGlu
TCCGTCTCTGTGAGTGCTATGACGAGGCTGTGCTTGGTATGAGCTCAAGCCCGCCGAG
AGGCAGGAGACACTCACGATACTGCGTCCGACACGAACCATACTCGAGTGGGGCGGCTC

4261 ThrThrValArgLeuArgAlaTyrMetAsnThrProGlyLeuProValCysGlnAspHis
ACTACAGTTAGGCTACGAGCGTACATGAACACCCCGGGGCTTCCCGTGTGCCAGGACCAT
TGATGTCAATCCGATGCTCGCATGTACTGTGGGGCCCCGAAGGGCACACGGTCTTGTA

4321 LeuGluPheTrpGluGlyValPheThrGlyLeuThrHisIleAspAlaHisPheLeuSer
CTTGAATTTTGGGAGGGCGTCTTACAGGCCTCACTCATATAGATGCCACTTCTATACC
GAACCTAAAACCCCTCCCGCAGAAATGTCCGGAGTGAGTATCTACGGGTGAAAGATAGG

4381 GlnThrLysGlnSerGlyGluAsnLeuProTyrLeuValAlaTyrGlnAlaThrValCys
CAGACAAAGCAGAGTGGGGAGAACCTTCCCTTACCTGGTAGCGTACCAAGCCACCGTGTGC
GTCTGTTTCTGCTCACCCCTCTTGAAGGAATGGACCATCGCATGGTTCTGGTGGCACACG

4441 AlaArgAlaGlnAlaProProProSerTrpAspGlnMetTrpLysCysLeuIleArgLeu
GCTAGGGCTCAAGCCCTCCCCATCGTGGGACCAGATGTGGAAGTGTGATTCGCCTC
CGATCCCGAGTTCGGGGAGGGGGTAGCACCTGGTCTACACCTTCAAACTAAGCGGAG

4501 LysProThrLeuHisGlyProThrProLeuLeuTyrArgLeuGlyAlaValGlnAsnGlu
AAGCCACCCCTCCATGGGCCAACCCCTGCTATACAGACTGGGCGCTGTTTCAAGATGAA
TTCGGGTGGGAGGTACCGGTTGTGGGGACGATATGTCTGACCCGACAAAGTCTTACTT

4561 IleThrLeuThrHisProValThrLysTyrIleMetThrCysMetSerAlaAspLeuGlu
ATCACCTGACGCACCCAGTCACCAAATACATCATGACATGTCATGTCGGCCGACCTGGAG
TAGTGGGACTGCGTGGGTCACTGGTTTATGTAGTACTGTACGTACAGCCGGCTGGACCTC

4621 ValValThrSerThrTrpValLeuValGlyGlyValLeuAlaAlaLeuAlaAlaTyrCys
GTCGTACAGAGCACCTGGGTGCTCGTTGGCGGCTCCTGGCTGCTTGGCCGCGTATTGC
CAGCAGTGCTCGTGGACCCAGAGCAACCGCCGAGGACCGACGAAACCGCGCATAACG

4681 LeuSerThrGlyCysValValIleValGlyArgValValLeuSerGlyLysProAlaIle
CTGTCAACAGGCTGCGTGGTCATAGTGGGCAGGGTCTGCTGTCCGGGAAGCCGGCAATC
GACAGTTGTCCGACGCACCACTATCACCCGTCCAGCAGAACAGGCCCTTCGGCCGTTAG

4741 IleProAspArgGluValLeuTyrArgGluPheAspGluMetGluGluCysSerGlnHis
ATACCTGACAGGGAAGTCTCTACCGAGAGTTTCGATGAGATGGAAGAGTGCTCTCAGCAC
TATGGACTGTCCCTTCAAGAGATGGCTCTCAAGCTACTCTACCTTCTCAGAGAGTCGTG

SUBSTITUTE SHEET

21 / 40

FIG. 16-6

4801 LeuProTyrIleGluGlnGlyMetMetLeuAlaGluGlnPheLysGlnLysAlaLeuGly
TTACCGTACATCGAGCAAGGGATGATGCTCGCCGAGCAGTTCAAGCAGAAGGCCCTCGGC
AATGGCATGTAGCTCGTTCCCTACTACGAGCGGCTCGTCAAGTTCGTCTTCCGGGAGCCG

4861 LeuLeuGlnThrAlaSerArgGlnAlaGluValIleAlaProAlaValGlnThrAsnTrp
CTCCTGCAGACCGCGTCCCGTCAGGCAGAGGTTATCGCCCCGTGTGCCAGACCAACTGG
GAGGACGTCTGGCGCAGGGCAGTCCGTCTCCAATAGCGGGACGACAGGTCTGGTTGACC

4921 GlnLysLeuGluThrPheTrpAlaLysHisMetTrpAsnPheIleSerGlyIleGlnTyr
CAAAAACCTCGAGACCTTCTGGGCGAAGCATATGTGGAACCTCATCAGTGGGATACAATAC
GTTTTGAGCTCTGGAAGACCCGCTTCGTATACACCTTGAAGTAGTCACCCTATGTTATG

4981 LeuAlaGlyLeuSerThrLeuProGlyAsnProAlaIleAlaSerLeuMetAlaPheThr
TTGGCGGGCTTGTCACGCTGCCTGGTAACCCGCCATTGCTTCATGTATGGCTTTTACA
AACCGCCCGAACAGTTGCGACGGACCATTTGGGCGGTAACGAAGTAACTACCGAAAATGT

5041 AlaAlaValThrSerProLeuThrThrSerGlnThrLeuLeuPheAsnIleLeuGlyGly
GCTGCTGTACCAGCCCACTAACCCTAGCCAAACCCCTCCTCTTCAACATATTGGGGGGG
CGACGACAGTGGTCGGGTGATTGGTGATCGGTTTGGGAGGAGAAGTTGTATAACCCCCC

5101 TrpValAlaAlaGlnLeuAlaAlaProGlyAlaAlaThrAlaPheValGlyAlaGlyLeu
TGGGTGGCTGCCAGCTCGCCGCCCCCGGTGCCGCTACTGCCTTTGTGGGCGCTGGCTTA
ACCCACCGACGGGTGAGCGGGCGGGGGCCACGGCGATGACGAAACACCCGCGACCGAAT

5161 AlaGlyAlaAlaIleGlySerValGlyLeuGlyLysValLeuIleAspIleLeuAlaGly
GCTGGCGCCGCCATCGGCAGTGTGGACTGGGGAAGGTCTCTATAGACATCCTTGACGGG
CGACCGCGGGGTAGCCGTCACAACCTGACCCCTTCCAGGAGTATCTGTAGGAACGTCCC

5221 TyrGlyAlaGlyValAlaGlyAlaLeuValAlaPheLysIleMetSerGlyGluValPro
TATGGCGCGGGCGTGGCGGGAGCTCTTGTGGCATTCAAGATCATGAGCGGTGAGGTCCCC
ATACCGCGCCCGCACCGCCCTCGAGAACACCGTAAGTTCTAGTACTCGCCACTCCAGGGG

5281 SerThrGluAspLeuValAsnLeuLeuProAlaIleLeuSerProGlyAlaLeuValVal
TCCACGGAGGACCTGGTCAATCTACTGCCCCCATCCTCTCGCCCGAGCCCTCGTAGTC
AGGTGCTCCTGGACCACTTAGATGACGGCGGTAGGAGAGCGGGCTCGGGAGCATCAG

5341 GlyValValCysAlaAlaIleLeuArgArgHisValGlyProGlyGluGlyAlaValGln
GGCGTGGTCTGTGCAGCAATACTGCGCCGGCACGTTGGCCCGGGCGAGGGGGCAGTGCAG
CCGCACCAGACACGTCGTTATGACGCGGGCGTGCAACCGGGCCCGCTCCCCCGTCACGTC

5401 TrpMetAsnArgLeuIleAlaPheAlaSerArgGlyAsnHisValSerProThrHisTyr
TGGATGAACCGGCTGATAGCCTTCGCCTCCCGGGGAACCATGTTTCCCCACGCACTAC
ACCTACTTGGCCGACTATCGGAAGCGGAGGGCCCCCTTGGTACAAAGGGGTGCGTGATG

5461 ValProGluSerAspAlaAlaAlaArgValThrAlaIleLeuSerSerLeuThrValThr
GTGCCGAGAGCGATGCAGCTGCCCCGCTCACTGCCATACTCAGCAGCTCACTGTAACC
CACGGCCTCTCGCTACGTCGACGGGCGCAGTGACGGTATGAGTCGTGAGTGACATTGG

5521 GlnLeuLeuArgArgLeuHisGlnTrpIleSerSerGluCysThrThrProCysSerGly
CAGCTCCTGAGGCGACTGCACAGTGGATAAGCTCGGAGTGTAACCTCCATGCTCCGGT
GTCGAGGACTCCGCTGACGTGTCACCTATTGAGCCTCACATGGTGAGGTACGAGGCCA

5581 SerTrpLeuArgAspIleTrpAspTrpIleCysGluValLeuSerAspPheLysThrTrp
TCCTGGCTAAGGGACATCTGGGACTGGATATGCGAGGTGTTGAGCGACTTTAAGACCTGG
AGGACCGATTCCCTGTAGACCTGACCTATACGCTCCACAACCTCGCTGAAATTCGGACC

5641 LeuLysAlaLysLeuMetProGlnLeuProGlyIleProPheValSerCysGlnArgGly
CTAAAGCTAAGCTCATGCCACAGCTGCCTGGGATCCCCCTTTGTGTCTGCCAGCGCGGG
GATTTTCGATTTCGAGTACGGTGTGACGGACCCTAGGGGAAACACAGGACGGTCGCGCCC

5701 TyrLysGlyValTrpArgValAspGlyIleMethHisThrArgCysHisCysGlyAlaGlu
TATAAGGGGTCTGGCGAGTGGACGGCATCATGCACACTCGCTGCCACTGTGGAGCTGAG
ATATCCCCCAGACCGCTCACCTGCCGTAGTACGTGTGAGCGACGGTGACACCTCGACTC

5761 IleThrGlyHisValLysAsnGlyThrMetArgIleValGlyProArgThrCysArgAsn
ATCACTGGACATGTCAAAAACGGGACGATGAGGATCGTGGTCTAGGACCTGCAGGAAC

SUBSTITUTE SHEET

22/40

FIG. 16-7

TAGTGACCTGTACAGTTTTTGCCCTGCTACTCCTAGCAGCCAGGATCCTGGACGTCTCTG

5821 MetTrpSerGlyThrPheProIleAsnAlaTyrThrThrGlyProCysThrProLeuPro
ATGTGGAGTGGGACCTTCCCCATTAAATGCCTACACCACGGGCCCCCTGTACCCCCCTTCTT
TACACCTCACCCCTGGAAGGGTAATTACGGATGTGGTGCCCGGGGACATGGGGGAAGGA

5881 AlaProAsnTyrThrPheAlaLeuTrpArgValSerAlaGluGluTyrValGluIleArg
GCGCCGAACCTACACGTTGCGCTATGGAGGGTGCTGTCAGAGGAATATCTGGAGATAAGG
CGCGGCTTGATGTGCAAGCGGATACCTCCACAGACGTCTCCTTATACACCTCTATTCC

5941 GlnValGlyAspPheHisTyrValThrGlyMetThrThrAspAsnLeuLysCysProCys
CAGGTGGGGGACTTCCACTACGTGACGGGTATGACTACTGACAATCTCAAATGCCCGTGC
GTCCACCCCTGAAGGTGATGCACTGCCATACTGATGACTGTTAGAGTTTACGGGCACG

6001 GlnValProSerProGluPhePheThrGluLeuAspGlyValArgLeuHisArgPheAla
CAGGTCCCATCGCCGAATTTTTTACAGAAATGGACGGGGTGCGCTACATAGGTTTGCG
GTCCAGGGTAGCGGGCTTAAAAAGTGTCTTAACCTGCCCCACGGGATGTATCCAAACGC

6061 ProProCysLysProLeuLeuArgGluGluValSerPheArgValGlyLeuHisGluTyr
CCCCCTGCAAGCCCTTGCTGCGGGAGGAGGTATCATTCAGAGTAGGACTCCACGAATAC
GGGGGACGTTCCGGGAACGACGCCCTCTCCATAGTAAGTCTCATCTGAGGTGCTTATG

6121 ProValGlySerGlnLeuProCysGluProGluProAspValAlaValLeuThrSerMet
CCGGTAGGGTCCGAATTACCTTGCGAGCCCGAACCGGACGTGGCCGTGTTGACGTCCATG
GGCCATCCAGCGTTAATGGAACGCTCGGGCTTGGCCTGCACCGGCACAACCTGCAGGTAC

6181 LeuThrAspProSerHisIleThrAlaGluAlaAlaGlyArgArgLeuAlaArgGlySer
CTCACTGATCCCTCCCATATAACAGCAGAGGCGGGCGGGCGAAGGTGGCGAGGGGATCA
GAGTGACTAGGGAGGGTATATTGTCGTCTCCGCGGGCCCGCTTCCAACCGCTCCCTAGT

6241 ProProSerValAlaSerSerSerAlaSerGlnLeuSerAlaProSerLeuLysAlaThr
CCCCCTCTGTGGCCAGCTCCTCGGCTAGCCAGCTATCCGCTCCATCTCTCAAGGCAACT
GGGGGAGACACCGGTCGAGGAGCCGATCGGTGATAGCGAGGTAGAGAGTTCCGTTGA

6301 CysThrAlaAsnHisAspSerProAspAlaGluLeuIleGluAlaAsnLeuLeuTrpArg
TGCACCGCTAACCATGACTCCCTGATGCTGAGCTCATAGAGGCAACCTCCTATGGAGG
ACGTGGCGATTGTTACTGAGGGGACTACGACTCGAGTATCTCCGTTGGAGGATACCTCC

6361 GlnGluMetGlyGlyAsnIleThrArgValGluSerGluAsnLysValValIleLeuAsp
CAGGAGATGGGCGGCAACATCACCAGGGTTGAGTCAGAAAACAAGTGGTGGTATTCTGGAC
GTCCTCTACCCGCCGTTGTAGTGGTCCCAACTCAGTCTTTTGTTCACCACTAAGACCTG

6421 SerPheAspProLeuValAlaGluGluAspGluArgGluIleSerValProAlaGluIle
TCCTTCGATCCGCTTGTGGCGGAGGAGGACGCGGGAGATCTCCGTACCCGCAGAAATC
AGGAAGCTAGGCGAACACCGCCTCCTCTGCTCGCCCTCTAGAGGCATGGGCGTCTTTAG

6481 LeuArgLysSerArgArgPheAlaGlnAlaLeuProValTrpAlaArgProAspTyrAsn
CTGCGGAAGTCTCGGAGATTGCGCCAGGCCCTGCCGCTTGGGCGCGGCCGACTATAAC
GACGCCCTCAGAGCCTCTAAGCGGGTCCGGGACGGGCAAACCCGCGCCGCTGATATTG

6541 ProProLeuValGluThrTrpLysLysProAspTyrGluProProValValHisGlyCys
CCCCGCTAGTGGAGACGTGGAAGAGCCCGACTACGAACCACCTGTGGTCCATGGCTGT
GGGGGCGATCACCTCTGCACCTTTTTCGGGCTGATGCTTGGTGGACACCAGGTACCGACA

6601 ProLeuProProProLysSerProProValProProProArgLysLysArgThrValVal
CCGCTTCCACCTCCAAAGTCCCTCCTGTGCCTCCGCTCGGAAGAAGCGGACGGTGGTC
GGCGAAGGTGGAGGTTTCAGGGGAGGACACGGAGGCGGAGCCTTCTTCGCTGCCACCAAG

6661 LeuThrGluSerThrLeuSerThrAlaLeuAlaGluLeuAlaThrArgSerPheGlySer
CTCACTGAATCAACCCTATCTACTGCCTTGGCCGAGCTCGCCACCAAGCTTTGGCAGC
GAGTGACTTAGTGGGATAGATGACGGAACCGGCTCGAGCGGTGGTCTTCGAAACCGTCG

6721 SerSerThrSerGlyIleThrGlyAspAsnThrThrThrSerSerGluProAlaProSer
TCCTCAACTTCCGGCATTACGGGCGACAATACGACAACATCCTCTGAGCCCGCCCCCTCT
AGGAGTTGAAGGCCGTAATGCCGCTGTTATGCTGTTGTAGGAGACTCGGGCGGGGAAGA

SUBSTITUTE SHEET

23/40

FIG. 16-8

6781 GlyCysProProAspSerAspAlaGluSerTyrSerSerMetProProLeuGluGlyGlu
GGCTGCCCCCGACTCCGACGCTGAGTCCCTATTCCTCCATGCCCCCCTGGAGGGGGAG
CCGACGGGGGGGCTGAGGCTGCGACTCAGGATAAGGAGGTACGGGGGGGACCTCCCCCTC

6841 ProGlyAspProAspLeuSerAspGlySerTrpSerThrValSerSerGluAlaAsnAla
CCTGGGGATCCGGATCTTAGCGACGGGTCAATGCTCAACGGTCAGTAGTGAGGCCAACGCG
GGACCCCTAGGCCTAGAATCGCTGCCCCAGTACCAGTTGCCAGTCATCACTCCGGTTGCGC

6901 GluAspValValCysCysSerMetSerTyrSerTrpThrGlyAlaLeuValThrProCys
GAGGATGTGCTGTGCTGCTCAATGTCTTACTCTTGACAGGCGCACTCGTCACCCCGTGC
CTCCTACAGCACACGACGAGTTACAGAATGAGAACCTGTCCGCGTGAGCAGTGGGGCAGC

6961 AlaAlaGluGluGlnLysLeuProIleAsnAlaLeuSerAsnSerLeuLeuArgHisHis
GCCGCGGAAGAACAGAACTGCCCATCAATGCCTAAGCAACTCGTTGCTACGTCACCCAC
CGGCGCCTTCTGTCTTTGACGGGTAGTTACGTGATTCTGTTAGCAACGATGCAGTGGTG

7021 AsnLeuValTyrSerThrThrSerArgSerAlaCysGlnArgGlnLysLysValThrPhe
AATTGGGTGATTCCACCACCTCAGCGAGTGCTTGCCAAAGGCAGAAAGTCACATTT
TTAAACCACATAAGGTGGTGGAGTGCGTCACGAACGGTTTCCGTCTTCTTTTCAGTGTA

7081 AspArgLeuGlnValLeuAspSerHisTyrGlnAspValLeuLysGluValLysAlaAla
GACAGACTGCAAGTTCTGGACAGCCATTACCAGGACGTACTCAAGGAGGTTAAAGCAGCG
CTGTCTGACGTTCAAGACCTGTCGGTAATGGTCTGTCATGAGTTCTCTCAATTTCTGCGC

7141 AlaSerLysValLysAlaAsnLeuLeuSerValGluGluAlaCysSerLeuThrProPro
GCGTCAAAAGTGAAGGCTAACTTGCTATCCGTAGAGGAAGCTTGACGCTGACGCCCCCA
CGCAGTTTCACTTCCGATTGAACGATAGGCATCTCTTCGAACGTCCGACTGCGGGGGT

7201 HisSerAlaLysSerLysPheGlyTyrGlyAlaLysAspValArgCysHisAlaArgLys
CACTCAGCCAAATCCAAGTTTGTTATGGGGCAAAGACGTCCGTTGCCATGCCAGAAAG
GTGAGTCGGTTTAGGTTCAAACCAATACCCGTTTCTGTCAGGCAACGGTACGGTCTTTC

7261 AlaValThrHisIleAsnSerValTrpLysAspLeuLeuGluAspAsnValThrProIle
GCCGTAAACCCACATCAACTCCGTGTGGAAAGACCTTCTGGAAGACAATGTAACACCAATA
CGGCATTGGGTGTAGTTGAGGCACACCTTCTGGAAGACCTTCTGTATACATTGTGGTTAT

7321 AspThrThrIleMetAlaLysAsnGluValPheCysValGlnProGluLysGlyGlyArg
GACACTACCATCATGGCTAAGAACGAGGTTTCTGCGTTTCAGCCTGAGAAGGGGGGTCTG
CTGTGATGGTAGTACCGATTCTTGCTCCAAAGACGCAAGTCGGAATCTTCCCCCAGCA

7381 LysProAlaArgLeuIleValPheProAspLeuGlyValArgValCysGluLysMetAla
AAGCCAGCTCGTCTCATCGTGTCCCGATCTGGGCGTGCGCGTGTGCGAAAAGATGGCT
TTCGGTCGAGCAGAGTAGCACAAAGGGGCTAGACCCGCACGCGCACACGCTTTTCTACCGA

7441 LeuTyrAspValValThrLysLeuProLeuAlaValMetGlySerSerTyrGlyPheGln
TTGTACGACGTGGTTACAAAGCTCCCTTGCCCGTGATGGGAAGCTCCTACGGATTCCAA
AACATGTGTCACCAATGTTTCGAGGGGAACCGGCACTACCTTCGAGGATGCCTAAGGTT

7501 TyrSerProGlyGlnArgValGluPheLeuValGlnAlaTrpLysSerLysLysThrPro
TACTCACCAGGACAGCGGTTGAATTCCTCGTCAAGCGTGAAGTCCAAGAAAACCCCA
ATGAGTGGTCTGTGCGCCCACTTAAGGAGCACGTTGCGACCTTCAGGTTCTTTTGGGGT

7561 MetGlyPheSerTyrAspThrArgCysPheAspSerThrValThrGluSerAspIleArg
ATGGGGTCTCGTATGATACCCGTGCTTTGACTCCACAGTCACTGAGAGCGACATCCGT
TACCCCAAGAGCATACTATGGGCGACGAACTGAGGTGTCACTGACTCTCGCTGTAGGCA

7621 ThrGluGluAlaIleTyrGlnCysCysAspLeuAspProGlnAlaArgValAlaIleLys
ACGGAGGAGCAATCTACCAATGTTGTGACCTCGACCCCAAGCCCGGTGGCCATCAAG
TGCTCTCTCGTTAGATGGTTACAACACTGGAGCTGGGGGTTCGGGCGACCGGTAGTTC

7681 SerLeuThrGluArgLeuTyrValGlyGlyProLeuThrAsnSerArgGlyGluAsnCys
TCCCTCACCAGAGGCTTTATGTTGGGGGCCCTCTTACCAATTCAAGGGGGGAGAACTGC
AGGGAGTGGCTCTCCGAAATACAACCCCGGGAGAATGGTTAAGTTCCCCCTCTTGACG

7741 GlyTyrArgArgCysArgAlaSerGlyValLeuThrThrSerCysGlyAsnThrLeuThr
GGCTATCGCAGGTGCCGCGAGCGGCTACTGACAACCTAGCTGTGGTAACACCCCTCACT

SUBSTITUTE SHEET

24 / 40

FIG. 16-9

CCGATAGCGTCCACGGCGCGCTCGCCGCATGACTGTTGATCGACACCATTTGTGGGAGTGA

7801 CysTyrIleLysAlaArgAlaAlaCysArgAlaAlaGlyLeuGlnAspCysThrMetLeu
TGCTACATCAAGGCCCGGGCAGCCTGTCGAGCCGAGGGCTCCAGGACTGCACCATGCTC
ACGATGTAGTTCCGGGCGCGTCGGACAGCTCGGCGTCCCGAGGTCCTGACGTGGTACGAG

7861 ValCysGlyAspAspLeuValValIleCysGluSerAlaGlyValGlnGluAspAlaAla
GTGTGTGGCGACGACTTAGTCGTTATCTGTGAAAGCGCGGGGGTCCAGGAGGACGCGGCG
CACACACCGCTGCTGAATCAGCAATAGACACTTTCGCGCCCCCAGGTCTCTCTGCGCCCG

7921 SerLeuArgAlaPheThrGluAlaMetThrArgTyrSerAlaProProGlyAspProPro
AGCCTGAGAGCCTTCACGGAGGCTATGACCAGGTACTCCGCCCCCCTGGGGACCCCCCA
TCGGACTCTCGGAAGTGCTCCGATACTGGTCCATGAGGCGGGGGGACCCCTGGGGGGT

7981 GlnProGluTyrAspLeuGluLeuIleThrSerCysSerSerAsnValSerValAlaHis
CAACCAGAATACGACTTGGAGCTCATAACATCATGCTCTCCAACGTGTGAGTCGCCCCAC
GTTGGTCTTATGCTGAACCTCGAGTATTGTAGTACGAGGAGGTGCACAGTCAGCGGGTG

8041 AspGlyAlaGlyLysArgValTyrTyrLeuThrArgAspProThrThrProLeuAlaArg
GACGGCGCTGGAAAGAGGGTCTACTACCTCACCCGTGACCCTACAACCCCCCTCGCGAGA
CTGCCGCGACCTTTCTCCAGATGATGGAGTGGGCACTGGGATGTTGGGGGGAGCGCTCT

8101 AlaAlaTrpGluThrAlaArgHisThrProValAsnSerTrpLeuGlyAsnIleIleMet
GCTGCGTGGGAGACAGCAAGACACACTCCAGTCAATTCCTGGCTAGGCAACATAATCATG
CGACGCACCCTCTGTCGTTCTGTGTGAGGTCAGTTAAGGACCGATCCGTTGTATTAGTAC

8161 PheAlaProThrLeuTrpAlaArgMetIleLeuMetThrHisPhePheSerValLeuIle
TTTGCCCCCACACTGTGGGCGAGGATGATACTGATGACCCATTCTTTAGCGTCCCTTATA
AAACGGGGGTGTGACACCCGCTCCTACTATGACTACTGGGTAAAGAAATCGCAGGAATAT

8221 AlaArgAspGlnLeuGluGlnAlaLeuAspCysGluIleTyrGlyAlaCysTyrSerIle
GCCAGGGACCAGCTTGAACAGGCCCTCGATTGCGAGATCTACGGGGCCTGCTACTCCATA
CGGTCCCTGGTCGAACCTGTCCGGGAGCTAACGCTCTAGATGCCCCGACGATGAGGTAT

8281 GluProLeuAspLeuProProIleIleGlnArgLeu
GAACCACTTGATCTACCTCCAATCATTCAAAGACTC
CTTGGTGAAC TAGATGGAGGTTAGTAAGTTTCTGAG

SUBSTITUTE SHEET

25 / 40

FIG. 17-1

-319 CACTCCACCATGAATCACTCCCTGTGAGGAAGTACTGTCTTCACGCAGAAAGCGTCTAG
GTGAGGTGGTACTTAGTGAGGGGACACTCCTTGATGACAGAAGTCCGTCTTTTCGCAGATC

-259 CCATGGCGTTAGTATGAGTGTCTGTGCAGCCTCCAGGACCCCCCTCCCGGGAGAGCCATA
GGTACCGCAATCATACTACAGCAGTCGGAGGTCTTGGGGGGAGGGCCCTCTCGGTAT

-199 GTGGTCTGCGGAACCGGTGAGTACACCGGAATTGCCAGGACACCGGGTCTCTTTCTTGGGA
CACCAGACGCCCTTGGCCACTCATGTGGCCTTAACGGTCTGTGGCCAGGAAAGAACCT

-139 TCAACCCGCTCAATGCCTGGAGATTGTTGGGCGTGCCCCGCAAGACTGCTAGCCGAGTAGT
AGTTGGGCGAGTTACGGACCTCTAAACCCGCACGGGGGCGTTCTGACGATCGGCTCATCA

- 79 GTTGGGTGCGGAAAGGCCTTGTGGTACTGCCTGATAGGGTGCTTGCAGTGCCCCGGGAG
CAACCCAGCGCTTTCCGGAACACCATGACGGACTATCCACGAACGCTCACGGGGCCCTC

- 19 GTCTCGTAGACCGTGCACC
CAGAGCATCTGGCACGTGG

Arg Thr
1 MetSerThrAsnProLysProGlnLysLysAsnLysArgAsnThrAsnArgArgProGln
ATGAGCACGAATCCTAAACCTCAAAAAAACAACCGTAACACCAACCGTCGCCCCACAG
TACTCGTGCTTAGGATTTGGAGTTTTTTTTTTGTTTGCATTGTGGTTGGCAGCGGGTGTCT

61 AspValLysPheProGlyGlyGlyGlnIleValGlyGlyValTyrLeuLeuProArgArg
GACGTCAAGTTCCCGGGTGGCGGTGAGATCGTTGGTGGAGTTACTTGTGTCGCGCAGG
CTGCAGTTCAAGGGCCACCGCCAGTCTAGCAACCACCTCAATGAACAACGGCGCGTCC

121 GlyProArgLeuGlyValArgAlaThrArgLysThrSerGluArgSerGlnProArgGly
GGCCCTAGATTGGGTGTGCGCGCAGCAGAAAGACTTCCGAGCGGTGCGAACCTCGAGGT
CCGGGATCTAACCACACGCGCGCTGCTCTTTCTGAAGGCTCGCCAGCGTTGGAGCTCCA

181 ArgArgGlnProIleProLysAlaArgArgProGluGlyArgThrTrpAlaGlnProGly
AGACGTCAGCCTATCCCAAGGCTCGTCGGCCCGAGGGCAGGACCTGGGCTCAGCCCGGG
TCTGCAGTCGGATAGGGGTTCGAGCAGCCGGGCTCCCGTCTGGACCCGAGTCGGGCCC

241 TyrProTrpProLeuTyrGlyAsnGluGlyCysGlyTrpAlaGlyTrpLeuLeuSerPro
TACCCTTGGCCCCCTCTATGGCAATGAGGGCTGCGGGTGGCGGGATGGCTCCTGTCTCCC
ATGGGAACCGGGGAGATACCGTTACTCCCGACGCCACCCGCCCTACCGAGGACAGAGGG

301 ArgGlySerArgProSerTrpGlyProThrAspProArgArgArgSerArgAsnLeuGly
CGTGGCTCTCGGCCTAGCTGGGGCCCCACAGACCCCGGCGTAGGTGCGCAATTTGGGT
GCACCGAGAGCCGGATCGACCCCGGGGTGTCTGGGGCCGCATCCAGCGCGTTAAACCCA

361 LysValIleAspThrLeuThrCysGlyPheAlaAspLeuMetGlyTyrIleProLeuVal
AAGTTCATCGATACCTTACGTGCGGCTTCGCCGACCTCATGGGGTACATACCGCTCGTC
TTCCAGTAGCTATGGGAATGCACGCCGAAGCGGCTGGAGTACCCCATGTATGGCGAGCAG

421 GlyAlaProLeuGlyGlyAlaAlaArgAlaLeuAlaHisGlyValArgValLeuGluAsp
GGCGCCCCCTCTTGGAGGCGCTGCCAGGGCCCTGGCGCATGGCGTCCGGGTCTTGAAGAC
CCGCGGGGAGAACCTCCGCGACGGTCCCGGGACCGGTACCGCAGGCCCAAGACCTTCTG

Thr
481 GlyValAsnTyrAlaThrGlyAsnLeuProGlyCysSerPheSerIlePheLeuLeuAla
GGCGTGAAGTATGCAACAGGGAACCTTCCCTGGTTGCTCTTTCTCTATCTTCTTCTGGCC
CCGCACTTGATACGTTGTCCCTTGAAGGACCAACGAGAAAGAGATAGAAGGAAGACCGG

541 LeuLeuSerCysLeuThrValProAlaSerAlaTyrGlnValArgAsnSerThrGlyLeu
CTGCTCTCTTGGCTTACTGTGCGGCTTCGGCCTACCAAGTGGCAACTCCACGGGGCTT
GACGAGAGAACGAACGACACGGGCGAAGCCGGATGGTTCACGCGTTGAGGTGCCCGGAA

601 TyrHisValThrAsnAspCysProAsnSerSerIleValTyrGluAlaAlaAspAlaIle
TACCACGTCACCAATGATTGCCCTAACTCGAGTATTGTGTACGAGGCGGCGCATGCCATC
ATGGTGCAGTGGTTACTAACGGGATTGAGCTCATAACACATGCTCCGCCGGCTACGGTAG

661 LeuHisThrProGlyCysValProCysValArgGluGlyAsnAlaSerArgCysTrpVal
CTGCACACTCCGGGGTGGCTCCCTTGGGTTGCTGAGGGCAACGCCCTCGAGGTGTTGGGTG
GACGTGTGAGGCCCCACGACGGGAACGCAAGCACTCCCGTTGCGGAGCTCCACAACCCAC

SUBSTITUTE SHEET

26/40

FIG. 17-2

- 721 AlaMetThrProThrValAlaThrArgAspGlyLysLeuProAlaThrGlnLeuArgArg
GCGATGACCCCTACGGTGGCCACCAGGGATGGCAAACCTCCCGGACGCAGCTTCGACGT
CGCTACTGGGGATGCCACCGGTGGTCCCTACCGTTTGAGGGGCGCTGCGTGAAGCTGCA
- 781 HisIleAspLeuLeuValGlySerAlaThrLeuCysSerAlaLeuTyrValGlyAspLeu
CACATCGATCTGCTTGTGGGAGCGCCACCCTCTGTTCCGGCCCTCTACGTGGGGGACCTA
GTGTAGCTAGACGAACAGCCCTCGCGGTGGGAGACAAGCCGGGAGATGCACCCCTGGAT
- 841 CysGlySerValPheLeuValGlyGlnLeuPheThrPheSerProArgArgHisTrpThr
TGCGGGTCTGTCTTTCTTGTGGCCAACTGTTACCTTCTCTCCAGGCGCCACTGGACG
ACGCCAGACAGAAAGAACAGCCGGTTGACAAGTGAAGAGAGGGTCCGCGGTGACCTGC
- 901 ThrGlnGlyCysAsnCysSerIleTyrProGlyHisIleThrGlyHisArgMetAlaTrp
ACGCAAGGTGCAATTGCTCTATCTATCCCGCCATATAACGGGTCAACGCATGGCATGG
TGCGTTCCAACGTTAACGAGATAGATAGGGCCGGTATATTGCCAGTGGCGTACCGTACC
- 961 Val
AspMetMetMetAsnTrpSerProThrThrAlaLeuValMetAlaGlnLeuLeuArgIle
GATATGATGATGAACCTGGTCCCTACGACGGCGTTGGTAAATGGCTCAGCTGCTCCGGATC
CTATACTACTACTTGACCAGGGGATGCTGCGCAACCATTACCGAGTCGACGAGGCCCTAG
- 1021 ProGlnAlaIleLeuAspMetIleAlaGlyAlaHisTrpGlyValLeuAlaGlyIleAla
CCACAAGCCATCTTGGACATGATCGCTGGTGTCTACTGGGGAGTCTTGGCGGGCATAGCG
GGTGTTCGGTAGAACCTGTACTAGCGACCAAGAGTGACCCCTCAGGACCGCCCGTATCGC
- 1081 TyrPheSerMetValGlyAsnTrpAlaLysValLeuValValLeuLeuPheAlaGly
TATTTCTCCATGGTGGGGAACCTGGGCGAAGGTCTCTGGTAGTGTCTGCTATTGTGCCGGC
ATAAAGAGGTACCACCCCTTGACCCGCTTCCAGGACCATCAGACGACGATAAACGGCCG
- 1141 ValAspAlaGluThrHisValThrGlyGlySerAlaGlyHisThrValSerGlyPheVal
GTCGACGCGGAAACCCACGTCACCGGGGGAAGTGCCGGCCACACTGTGTCTGGATTGTGTT
CAGCTGCGCCTTTGGGTGCAGTGGCCCCCTTCACGGCCGGTGTGACACAGACCTAAACAA
- 1201 SerLeuLeuAlaProGlyAlaLysGlnAsnValGlnLeuIleAsnThrAsnGlySerTrp
AGCCTCCTCGCACCAGGCGCCAAGCAGAACGTCCAGCTGATCAACACCAACGGCAGTTGG
TCGGAGGAGCGTGGTCCGCGGTTTCGTCTTGACGGTGCAGTAGTTGTGGTTGCCGTCAACC
- 1261 HisLeuAsnSerThrAlaLeuAsnCysAsnAspSerLeuAsnThrGlyTrpLeuAlaGly
CACCTCAATAGCACGGCCCTGAATGCAATGATAGCCTCAACACCGGCTGGTTGGCAGGG
GTGGAGTTATCGTGGCGGGACTTGACGTTACTATCGGAGTTGTGGCCGACCAACCGTCCC
- 1321 LeuPheTyrHisHisLysPheAsnSerSerGlyCysProGluArgLeuAlaSerCysArg
CTTTCTATACCA CAAGTTCAACTCTTCAGGCTGTCTGAGAGGCTAGCCAGCTGCCGA
GAAAAGATAGTGGTGTCAAGTTGAGAAGTCCGACAGGACTCTCCGATCGGTGACGGCT
- 1381 ProLeuThrAspPheAspGlnGlyTrpGlyProIleSerTyrAlaAsnGlySerGlyPro
CCCCTTACCGATTTTGACCAGGGCTGGGGCCCTATCAGTTATGCCAACGGAAGCGGCCCC
GGGGAATGGCTAAACTGGTCCCGACCCCGGATAGTCAATACGGTTGCCTTCGCCGGGG
- 1441 AspGlnArgProTyrCysTrpHisTyrProProLysProCysGlyIleValProAlaLys
GACCAGCGCCCTACTGCTGGCACTACCCCCAAAACCTTGCGGTATTGTGCCCCGGAAG
CTGGTGGCGGGGATGACGACCGTATGGGGGGTTTGGAAACGCATAACACGGGCGCTTC
- 1501 SerValCysGlyProValTyrCysPheThrProSerProValValValGlyThrThrAsp
AGTGTGTGTGGTCCGGTATATTGCTTCACTCCAGCCCCGTGGTGGTGGGAACGACCGAC
TCACACACACCAGGCCATATAACGAAGTGAGGGTCGGGGCACCACCACCTTGTGCGCTG
- 1561 ArgSerGlyAlaProThrTyrSerTrpGlyGluAsnAspThrAspValPheValLeuAsn
AGGTGGGGCGCGCCACCTACAGCTGGGGTGAAAATGATACGGACGTCTTCGTCTTAAC
TCCAGCCCCGCGGGTGGATGTGACCCCACTTTTACTATGCTGCAGAAGCAGGAATTG
- 1621 AsnThrArgProProLeuGlyAsnTrpPheGlyCysThrTrpMetAsnSerThrGlyPhe
AATACCAGGCCACCGCTGGGCAATTGGTTCCGGTTGTACCTGGATGAACCTCACTGGATTTC
TTATGGTCCGGTGGCGACCCGTTAACCAAGCCAACATGGACCTACTTGAGTTGACCTAAG

SUBSTITUTE SHEET

27 / 40

FIG. 17-3

- 1681 ThrLysValCysGlyAlaProProCysValIleGlyGlyAlaGlyAsnAsnThrLeuHis
ACCAAAGTGTGCGGAGCGCTCCTTGTGTCATCGGAGGGGCGGGCAACAACCCCTGCAC
TGGTTTCACACGCCTGCGGAGGAACACAGTAGCCTCCCGCCCGTTGTTGTGGGACGTG
- 1741 CysProThrAspCysPheArgLysHisProAspAlaThrTyrSerArgCysGlySerGly
TGCCCCACTGATTGCTTCCGCAAGCATCCGACGCCACATACTCTCGGTGCGGCTCCGCT
ACGGGGTGACTAACGAAGGCGTTCGTAGGCCTGCGGTGTATGAGAGCCACGCCGAGGCCA
- Leu
- 1801 ProTrpIleThrProArgCysLeuValAspTyrProTyrArgLeuTrpHisTyrProCys
CCCTGGATCACACCCAGGTGCTGCTGACTACCCGTATAGGCTTTGGCATTATCCTTGT
GGGACCTAGTGTGGGTCCACGGACCAGCTGATGGGCATATCCGAAACCGTAATAGGAACA
- 1861 ThrIleAsnTyrThrIlePheLysIleArgMetTyrValGlyGlyValGluHisArgLeu
ACCATCAACTACACCATATTTAAATCAGGATGTACGTGGGAGGGGTCGAACACAGGCTG
TGGTAGTTGATGTGGTATAAATTTTAGTCTACATGCACCTCCCGAGCTTGTGTCGAC
- 1921 GluAlaAlaCysAsnTrpThrArgGlyGluArgCysAspLeuGluAspArgAspArgSer
GAAGCTGCCTGCAACTGGACGCGGGGCGAACGTGCGATCTGGAAGACAGGGACAGGTCC
CTTCGACGGACGTTGACCTGCGCCCGCTTGCAACGCTAGACCTTCTGTCCCTGTCCAGG
- 1981 GluLeuSerProLeuLeuLeuThrThrThrGlnTrpGlnValLeuProCysSerPheThr
GAGCTCAGCCCGTTACTGCTGACCACTACACAGTGGCAGGTCTCCCGTGTTCCTTCACA
CTCGAGTCGGGCAATGACGACTGGTGTATGTCACCGTCCAGGAGGCACAAGGAAGTGT
- 2041 ThrLeuProAlaLeuSerThrGlyLeuIleHisLeuHisGlnAsnIleValAspValGln
ACCCTACAGCCTTGTCCACCGCCTCATCCACCTCCACCAGAACATTTGGACGCTGCAG
TGGGATGGTTCGGAACAGGTGGCCGGAGTAGGTGGAGGTGGTCTTGTAAACCTGCACGTC
- 2101 TyrLeuTyrGlyValGlySerSerIleAlaSerTrpAlaIleLysTrpGluTyrValVal
TACTTGTACGGGGTGGGGTCAAGCATCGCGTCTCGGGCCATTAGTGGGAGTACGTCTGTT
ATGAACATGCCCCACCCAGTTCGTAGCGCAGGACCGGTAATTCAACCTCATGCAGCAA
- 2161 LeuLeuPheLeuLeuLeuAlaAspAlaArgValCysSerCysLeuTrpMetMetLeuLeu
CTCCTGTCTCTCTGCTTGACAGCGCGCGCTCTGCTCCTGCTTGTGGATGATGCTACTC
GAGGACAAGGAAGACGAACGCTGCGCGCGCAGACGAGGACGAACACCTACTACGATGAG
- 2221 IleSerGlnAlaGluAlaAlaLeuGluAsnLeuValIleLeuAsnAlaAlaSerLeuAla
ATATCCCAAGCGGAGCGGCTTGGAGAACCTCGTAATACCTAATGCAGCATCCCTGGCC
TATAGGGTTCGCTCCGCGAAACCTCTTGGAGCATTATGAATTACGTCGTAGGGACCGG
- 2281 GlyThrHisGlyLeuValSerPheLeuValPhePheCysPheAlaTrpTyrLeuLysGly
GGGACGCACGGTCTTGTATCCTTCTCTGTTCTTCTGCTTGTGATGGTATTTGAAGGGT
CCCTGCGTGCCAGAACATAGGAAGGAGCACAAGAAGACGAAACGTACCATAAACTTCCCA
- 2341 LysTrpValProGlyAlaValTyrThrPheTyrGlyMetTrpProLeuLeuLeuLeuLeu
AAGTGGGTGCCCCGAGCGGTCTACACCTTCTACGGGATGTGGCTCTCTCTCTGCTCTG
TTCACCCACGGGCTCGCCAGATGTGGAAGATGCCCTACACCGGAGAGGAGGACGAGGAC
- 2401 LeuAlaLeuProGlnArgAlaTyrAlaLeuAspThrGluValAlaAlaSerCysGlyGly
TTGGCGTTGCCCCAGCGGGCTACGCGCTGGACACGGAGGTGGCCGCGTCTGTGGCGGT
AACCGCAACGGGGTTCGCGCATGCGGACCTGTGCCTCCACCGGCGCAGCACCGCCCA
- 2461 ValValLeuValGlyLeuMetAlaLeuThrLeuSerProTyrTyrLysArgTyrIleSer
GTTGTTCTCTGTCGGGTGATGGCGCTGACTCTGTACCATATTACAAGCGCTATATCAGC
CAACAAGAGCAGCCCACTACCGGACTGAGACAGTGGTATAATGTTTCGGGATATAGTCG
- Asn
- 2521 TrpCysLeuTrpTrpLeuGlnTyrPheLeuThrArgValGluAlaGlnLeuHisValTrp
TGGTGCTTGTGGTGGCTTCAGTATTTTCTGACCAGAGTGGAAGCGCAACTGCACGTGTGG
ACCACGAACACCACCAAGTCATAAAGACTGGTCTCACTTCGCGTTGACGTGCACACC
- 2581 IleProProLeuAsnValArgGlyGlyArgAspAlaValIleLeuLeuMetCysAlaVal
ATTCCCCCTCAACGTCCGAGGGGGCGGACGCGCTCATCTTACTCATGTGTGCTGTA
TAAGGGGGGGAGTTGCAGGCTCCCCCGGCTGCGGCAGTAGAATGAGTACACACGACAT

SUBSTITUTE SHEET

28 / 40

FIG. 17-4

2641 HisProThrLeuValPheAspIleThrLysLeuLeuAlaValPheGlyProLeuTrp
CACCCGACTCTGGTATTGACATCACCAAATTGCTGCTGGCCGCTCTCGGACCCTTTGG
GTGGGCTGAGACCATAAACTGTAGTGGTTTAAACGACGACCGGCAGAACCTGGGGAAACC

2701 IleLeuGlnAlaSerLeuLysValProTyrPheValArgValGlnGlyLeuLeuArg
ATTCTTCAAGCCAGTTTGCTTAAAGTACCCTACTTTGTGCGCGTCCAAGGCCTTCTCCGG
TAAGAAGTTCGGTCAAACGAATTTTCATGGGATGAAACACGCGCAGGTTCGGGAAGAGGCC

2761 PheCysAlaLeuAlaArgLysMetIleGlyGlyHisTyrValGlnMetValIleIleLys
TTCTGCGCGTTAGCGCGAAGATGATCGGAGGCCATTACGTGCAATGGTCATCATTAAG
AAGACGCGCAATCGCGCCTTCTACTAGCCTCCGGTAATGCAGTTTACCAGTAGTAATTC

2821 LeuGlyAlaLeuThrGlyThrTyrValTyrAsnHisLeuThrProLeuArgAspTrpAla
TTAGGGGCGCTTACTGGCACCATTGTTTATAACCATCTCACTCCTCTTCGGGACTGGGCG
AATCCCCGGAATGACCGTGGATACAAATATTGGTAGAGTGAGGAGAAGCCCTGACCCGC

2881 HisAsnGlyLeuArgAspLeuAlaValAlaValGluProValValPheSerGlnMetGlu
CACAACGCGCTTGGGAGATCTGGCCGTGGCTGTAGAGCCAGTCGTCTTCTCCCAAATGGAG
GTGTTGCCGAACGCTCTAGACCGGCACCGACATCTCGGTCAGCAGAAGAGGGTTTACCTC

2941 ThrLysLeuIleThrTrpGlyAlaAspThrAlaAlaCysGlyAspIleIleAsnGlyLeu
ACCAAGCTCATCAGTGGGGGCGAGATACCGCCGCGTGGTGACATCATCAACGGCTTG
TGGTTCGAGTAGTGCACCCCCGCTCTATGGCGGCGCACGCCACTGTAGTAGTTGCCGAAC

3001 ProValSerAlaArgArgGlyArgGluIleLeuLeuGlyProAlaAspGlyMetValSer
CCTGTTTCCGCCCGCAGGGGCGGGAGATACTGCTCGGGCCAGCCGATGGAATGGTCTCC
GGACAAAGGCGGGCGTCCCCGGCCCTCTATGACGAGCCCGGTCGGCTACCTTACCAGAGG

3061 LysGlyTrpArgLeuLeuAlaProIleThrAlaTyrAlaGlnGlnThrArgGlyLeuLeu
AAGGGGTGGAGGTTGCTGCGCGCCATCACGGCGTAGCGCCAGCAGACAAGGGGCTCCTA
TTCCCCACCTCCAACGACCGCGGGTAGTGC CGCATGCGGGTCTGTTCCTCCCGGAGGAT

3121 GlyCysIleIleThrSerLeuThrGlyArgAspLysAsnGlnValGluGlyGluValGln
GGGTGCATAATCACCAGCCTAACTGGCCGGGACAAAAACCAAGTGGAGGTGAGGTCCAG
CCCACGTATTAGTGGTGGATTGACCGGCCCTGTTTTTGGTTACCTCCCACTCCAGGTC

3181 IleValSerThrAlaAlaGlnThrPheLeuAlaThrCysIleAsnGlyValCysTrpThr
ATTGTGTCAACTGCTGCCAAACCTTCTGGCAACGTGCATCAATGGGGTGTGCTGGACT
TAACACAGTTGACGACGGGTTTGAAGGACCGTTGCACGTAGTTACCCCAACGACCTGA

3241 ValTyrHisGlyAlaGlyThrArgThrIleAlaSerProLysGlyProValIleGlnMet
GTCTACCACGGGGCCGGAACGAGGACCATCGCGTCACCAAGGGTCTGTGTCATCCAGATG
CAGATGGTGGCCCCGGCCTTGCTCCTGGTAGCGCAGTGGGTTCCAGGACAGTAGGTCTAC

3301 TyrThrAsnValAspGlnAspLeuValGlyTrpProAlaProGlnGlySerArgSerLeu
TATACCAATGTAGACCAAGACCTTGTTGGGCTGGCCCGCTCCGCAAGGTAGCCGCTCATTG
ATATGGTTACATCTGGTTCTGGAACACCCGACCGGGCGAGGCGTTCCATCGGCGAGTAAC

3361 ThrProCysThrCysGlySerSerAspLeuTyrLeuValThrArgHisAlaAspValIle
ACACCCTGCCTTGGCGGCTCCTCGGACCTTTACCTGGTACGAGGCACGCCGATGTCATT
TGTGGGACGTGAACGCCGAGGAGCCTGGAAATGGACCAGTGTCCGTGCGGCTACAGTAA

3421 ProValArgArgArgGlyAspSerArgGlySerLeuLeuSerProArgProIleSerTyr
CCCGTGGCCCGGCGGGGTGATAGCAGGGGCGAGCCTGCTGTGCCCCGGGCCATTTCCTAC
GGGCACGCGCCCGCCCACTATCGTCCCCGTCGGACGACAGCGGGGCGGGTAAAGGATG

3481 LeuLysGlySerSerGlyGlyProLeuLeuCysProAlaGlyHisAlaValGlyIlePhe
TTGAAAGCTCCTCGGGGGTCCGCTGTTGTGCCCCGCGGGGACGCGCGTGGGCATATTT
AACTTTCCGAGGAGCCCCCAGGCGACAACACGGGGCGCCCCGTGCGGCACCCGTATAAA

3541 ArgAlaAlaValCysThrArgGlyValAlaLysAlaValAspPheIleProValGluAsn
AGGGCCGCGGTGTGCACCCGTGGAGTGGCTAAGGCGGTGGACTTTATCCCTGTGGAGAAC
TCCCGCGCCACACGTGGGCACCTACCGATTCCGCCACCTGAAATAGGGACACCTCTTG

SUBSTITUTE SHEET

29/40

FIG. 17-5

- 3601 LeuGluThrThrMetArgSerProValPheThrAspAsnSerSerProProValValPro
CTAGAGACAACCATGAGGTCCCCGGTGTTCACGGATAACTCTCTCCACCAGTAGTGCCC
GATCTCTGTTGGTACTCCAGGGGCCACAAGTGCCTATTGAGGAGAGGTGGTCATCAGGGG
- 3661 GlnSerPheGlnValAlaHisLeuHisAlaProThrGlySerGlyLysSerThrLysVal
CAGAGCTTCCAGGTGGCTCACCTCCATGCTCCACAGGCAGCGCAAAAGCACCAAGGTC
GTCTCGAAGGTCCACCGAGTGGAGGTACGAGGGTGTCCGTGCGCGTTTTCGTGGTTCCAG
- 3721 ProAlaAlaTyrAlaAlaGlnGlyTyrLysValLeuValLeuAsnProSerValAlaAla
CCGGCTGCATATGCAGCTCAGGGCTATAAGGTGCTAGTACTCAACCCCTCTGTTGCTGCA
GGCCGACGTATACGTGAGTCCCGATATTCCACGATCATGAGTTGGGGAGACAACGACGT
- 3781 ThrLeuGlyPheGlyAlaTyrMetSerLysAlaHisGlyIleAspProAsnIleArgThr
ACACTGGGCTTTGGTGCTTACATGTCCAAGGCTCATGGGATCGATCCTAACATCAGGACC
TGTGACCCGAAACCACGAATGTACAGGTTCCGAGTACCCTAGCTAGGATTGTAGTCTGG
- 3841 GlyValArgThrIleThrThrGlySerProIleThrTyrSerThrTyrGlyLysPheLeu
GGGGTGAGAACAATTACCACTGGCAGCCCCATCAGTACTCCACCTACGGCAAGTTCCCTT
CCCCACTCTTGTAAATGGTGACCGTCCGGGGTAGTGCATGAGGTGGATGCCGTTCAAGGAA
- 3901 AlaAspGlyGlyCysSerGlyGlyAlaTyrAspIleIleIleCysAspGluCysHisSer
GCCGACGGCGGGTGTCTCGGGGGCGCTTATGACATAATAATTTGTGACGAGTGCCACTCC
CGCTGCGCGCCACGAGCCCCCGGAATACTGTATTATTAAACACTGCTCACGGTGAGG
- 3961 ThrAspAlaThrSerIleLeuGlyIleGlyThrValLeuAspGlnAlaGluThrAlaGly
ACGGATGCCACATCCATCTTGGGCATCGGCACTGTCTTGACCAAGCAGAGACTGCGGGG
TGCTTACGGTGTAGGTAGAACCCGTAGCCGTGACAGGAAGTGGTTCGTCTCTGACGCCCC
- 4021 AlaArgLeuValValLeuAlaThrAlaThrProProGlySerValThrValProHisPro
GCGAGACTGGTTGTGCTCGCCACCGCCACCCCTCCGGGCTCCGTCACTGTGCCCCATCCC
CGCTCTGACCAACACGAGCGGTGGCGGTGGGGAGGCCCGAGGCAGTGACAGGGGTAGGG
- 4081 AsnIleGluGluValAlaLeuSerThrThrGlyGluIleProPheTyrGlyLysAlaIle
AACATCGAGGAGGTGCTCTGTCCACCACCGGAGAGATCCCTTTTACGGCAAGGCTATC
TTGTAGCTCCTCCAACGAGACAGGTGGTGGCTCTCTAGGGAAAATGCCGTTCCGATAG
- 4141 ProLeuGluValIleLysGlyGlyArgHisLeuIlePheCysHisSerLysLysLysCys
CCCCTCGAAGTAATCAAGGGGGGAGACATCTCATCTCTGTCAATCAAGAAGAAGTGC
GGGGAGCTTCATTAGTTCCCCCCTCTGTAGAGTAGAAGACAGTAAGTTCTTCTTCAAG
- 4201 AspGluLeuAlaAlaLysLeuValAlaLeuGlyIleAsnAlaValAlaTyrTyrArgGly
GACGAACCTCGCCGCAAGCTGGTCGCATTGGGCATCAATGCCGTGGCCTACTACCGCGGT
CTGCTTGAGCGGCGTTTCGACCAGCGTAACCCGTAGTTACGGCACCGGATGATGGCGCCA
- 4261 LeuAspValSerValIleProThrSerGlyAspValValValAlaThrAspAlaLeu
CTTGACGTGTCGTCATCCCGACAGCGCGATGTGTGTCGTGGCAACCGATGCCCTC
GAACTGCACAGGCAGTAGGGCTGGTCCCGCTACAACAGCAGCACCCTTGGCTACGGGAG
- 4321 MetThrGlyTyrThrGlyAspPheAspSerValIleAspCysAsnThrCysValThrGln
ATGACCGGCTATACCGGGCACTTCGACTCGGTGATAGACTGCAATACGTGTGTCACCCAG
TACTGGCCGATATGGCCGCTGAAGCTGAGCCACTATCTGACGTTATGCAACAGTGGGTCT
- 4381 ThrValAspPheSerLeuAspProThrPheThrIleGluThrIleThrLeuProGlnAsp
ACAGTCGATTTACAGCCTTGACCCTACCTTCACCATTGAGACAATCACGCTCCCCAGGAT
TGTCAGCTAAAGTCGGAAGTGGGATGGAAGTGGTAACCTCTGTTAGTGCGAGGGGGTCTTA
- 4441 AlaValSerArgThrGlnArgArgGlyArgThrGlyArgGlyLysProGlyIleTyrArg
GCTGTCTCCCGCACTCAACGTCGGGGCAGGACTGGCAGGGGGAAGCCAGGCATCTACAGA
CGACAGAGGGCGTGAGTTGCAGCCCCGTCTGACCGTCCCCCTTCCGTCGGTAGATGTCT
- 4501 PheValAlaProGlyGluArgProSerGlyMetPheAspSerSerValLeuCysGluCys
TTTGTGGCACCGGGGGAGCGCCCTCCGGCATGTTGACTCGTCCGTCTCTGTGAGTGC
AAACACCGTGGCCCCCTCGCGGGGAGGCCGTACAAGCTGAGCAGGCAGGAGACACTCAGC

SUBSTITUTE SHEET

30 / 40

FIG. 17-6

4561 TyrAspAlaGlyCysAlaTrpTyrGluLeuThrProAlaGluThrThrValArgLeuArg
TATGACGCAGGCTGTGCTTGGTATGAGCTACGCCCGGAGACTACAGTTAGGCTACGA
ATACTGCGTCCGACACGAACCATACTCGAGTGC GGCGGCTCTGATGTCAATCCGATGCT

4621 AlaTyrMetAsnThrProGlyLeuProValCysGlnAspHisLeuGluPheTrpGluGly
GCGTACATGAACACCCCGGGGCTTCCCGTGTGCCAGGACCATCTTGAATTTGGGAGGGC
CGCATGTACTTGTGGGGCCCCGAAGGGCACACGGTCCCTGGTAGAACTTAAACCCCTCCG

4681 ValPheThrGlyLeuThrHisIleAspAlaHisPheLeuSerGlnThrLysGlnSerGly
GTCTTTACAGGCCCTCACTCATATAGATGCCCACTTTCTATCCCAGACAAAGCAGAGTGGG
CAGAAATGTCGGAGTGAGTATATCTACGGGTGAAAGATAGGGTCTGTTTCGTCTCACC

4741 GluAsnLeuProTyrLeuValAlaTyrGlnAlaThrValCysAlaArgAlaGlnAlaPro
GAGAACCCTCCTTACCTGGTAGCGTACCAAGCCACCGTGTGGCTAGGGCTCAAGCCCCT
CTCTTGAAGGAATGGACCATCGCATGGTTCGGTGGCACACGGATCCCGAGTTCGGGGA

4801 ProProSerTrpAspGlnMetTrpLysCysLeuIleArgLeuLysProThrLeuHisGly
CCCCATCGTGGGACAGATGTGGAGTGTGATTGCTCAAGCCCACCTCCATGGG
GGGGTAGCACCCCTGGTCTACACCTTCAAACTAAGCGGAGTTCGGGTGGGAGGTACCC

4861 ProThrProLeuLeuTyrArgLeuGlyAlaValGlnAsnGluIleThrLeuThrHisPro
CCAAACCCCTGCTATACAGACTGGGCGCTGTTCAGAAATGAAATCACCTGACGCACCCA
GGTTGTGGGGACGATATGTCTGACCCGCGACAAGTCTTACTTTAGTGGGACTCGTGGGT

4921 ValThrLysTyrIleMetThrCysMetSerAlaAspLeuGluValValThrSerThrTrp
GTCACCAAATACATCATGACATGCATGTGGCCGACCTGGAGGTCTGTCAGACACCTGG
CAGTGGTTTATGTAGTACTGTAGTACAGCGGCTGGACCTCCAGCAGTCTCGTGGACC

4981 ValLeuValGlyGlyValLeuAlaAlaLeuAlaTyrCysLeuSerThrGlyCysVal
GTGCTCGTTGGCGGCTCCTGGCTGCTTTGGCCGCTATTGCCTGTCAACAGGCTGCGTG
CAGAGCAACCGCCGAGACCGACGAAACCGGCGCATAACGGACAGTTGTCCGACGCAC

5041 ValIleValGlyArgValValLeuSerGlyLysProAlaIleIleProAspArgGluVal
GTCATAGTGGGCAGGGTCTCTGTGTCGGGAAGCCGCAATCATACCTGACAGGAAGTC
CAGTATCACCCGTCCAGCAGAACAGGCCCTTCGGCCGTAGTATGGAAGTCTCCCTTCAG

5101 LeuTyrArgGluPheAspGluMetGluGluCysSerGlnHisLeuProTyrIleGluGln
CTCTACGAGAGTTCTGATGAGATGGAAGAGTGCTCTCAGCACTTACCGTACATCGAGCAA
GAGATGGCTCTCAAGCTACTCTACCTTCTCAGAGAGTCTGTAATGGCATGTAGCTCGTT

5161 GlyMetMetLeuAlaGluGlnPheLysGlnLysAlaLeuGlyLeuLeuGlnThrAlaSer
GGGATGATGCTCGCCGAGCAGTTCAAGCAGAGGCCCTCGGCCTCTGCAGACCGCGTCC
CCCTACTACGAGCGGCTCGTCAAGTCTCTTCCGGGAGCCGGAGAGCTCTGGCGCAGG

5221 ArgGlnAlaGluValIleAlaProAlaValGlnThrAsnTrpGlnLysLeuGluThrPhe
CGTCAGGCAGAGGTATCGCCCCTGCTGTCCAGACCAACTGGCAAACTCGAGACCTTC
GCAGTCCGTCTCCAATAGCGGGGACGACAGGTCTGTTGACGTTTTTGTAGCTCTGGAAG

5281 TrpAlaLysHisMetTrpAsnPheIleSerGlyIleGlnTyrLeuAlaGlyLeuSerThr
TGGGCGAAGCATATGTGGAAGTTCATCAGTGGGATACAATACTTGGCGGGCTTGTCAACG
ACCCGCTTCGTATACACCTTGAAGTAGTCAACCTATGTTATGAACCGCCCGAACAGTTGC

5341 LeuProGlyAsnProAlaIleAlaSerLeuMetAlaPheThrAlaAlaValThrSerPro
CTGCCTGGTAACCCCGCCATTGCTTCATGATGGCTTTTACAGCTGCTGTACACGCCCCA
GACGGACCATTGGGGCGGTAACGAAGTAACTACGAAAATGTCAGACAGTGGTCCGGT

5401 LeuThrThrSerGlnThrLeuLeuPheAsnIleLeuGlyGlyTrpValAlaAlaGlnLeu
CTAACCACTAGCCAAACCTCTTCAACATATTGGGGGGGTGGGTGGCTGCCAGCTC
GATTGGTGATCGGTTTGGGAGGAGAAGTTGTATAACCCCCCACCACCGAGGGTTCGAG

5461 AlaAlaProGlyAlaAlaThrAlaPheValGlyAlaGlyLeuAlaGlyAlaAlaIleGly
GCCGCCCCGGTGCCGCTACTGCTTGTGGGGCGCTGGCTAGCTGGCGCCGCTATCGGC
CGCGGGGGCCACGGCGATGACGGAAACACCGCGACCGAATCGACCGCGGGGTAGCCG

SerValGlyLeuGlyLysValLeuIleAspIleLeuAlaGlyTyrGlyAlaGlyValAla

SUBSTITUTE SHEET

31 / 40

FIG. 17-7

- 5521 AGTGTGGACTGGGGAAGGTCTCATAGACATCCTTGCAGGGTATGGCGCGGGCGTGGCG
TCACAACCTGACCCCTTCCAGGAGTATCTGTAGGAACGTCCCATAACGCGCCCGCACCGC
- 5581 GlyAlaLeuValAlaPheLysIleMetSerGlyGluValProSerThrGluAspLeuVal
GGAGCTCTTGTGGCATTCAAGATCATGAGCGGTGAGGTCCCCTCCACGGAGGACCTGGTC
CCTCGAGAACACCGTAAGTTCTAGTACTCGCCACTCCAGGGGAGGTGCCTCTTGACCAG
- 5641 AsnLeuLeuProAlaIleLeuSerProGlyAlaLeuValValGlyValValCysAlaAla
AATCTACTGCCCCCATCTCTCGCCCGGAGCCCTCGTAGTCGGCGTGGTCTGTGCAGCA
TTAGATGACGGGCGGTAGGAGAGCGGGCTCGGGAGCATCAGCCGACCAGACACGTCGT
- 5701 IleLeuArgArgHisValGlyProGlyGluGlyAlaValGlnTrpMetAsnArgLeuIle
ATACTGCGCCGCGACGTGGCCCGGGCGAGGGGCGAGTG CAGTGGATGAACCGGCTGATA
TATGACGCGGCCGTGCAACCGGCGCCGCTCCCCCGTCACGTCACCTACTTGCCCGACTAT
- 5761 AlaPheAlaSerArgGlyAsnHisValSerProThrHisTyrValProGluSerAspAla
GCCTTCGCCTCCCGGGGAACCATGTTTCCCCACGCACTACGTGCCGAGAGCGATGCA
CGGAAGCGGAGGGCCCCCTTGGTACAAAGGGGTGCGTGATGCACGCCCTCTCGCTACGT
- 5821 HisCys
AlaAlaArgValThrAlaIleLeuSerSerLeuThrValThrGlnLeuLeuArgArgLeu
GCTGCCCGCGTCACTGCCATACTCAGCAGCCTCACTGTAAACCCAGCTCCTGAGGCGACTG
CGACGGGCGCAGTGACGGTATGAGTCGTGGAGTGACATTGGGTGAGGACTCCGCTGAC
- 5881 HisGlnTrpIleSerSerGluCysThrThrProCysSerGlySerTrpLeuArgAspIle
CACCAGTGGATAAGCTCGGAGTGTACCACTCCATGCTCCGGTTCCTGGCTAAGGGACATC
GTGGTCACCTATTTCGAGCCTCACATGGTGAGGTACGAGGCCAAGGACCGATTCCCTGTAG
- 5941 TrpAspTrpIleCysGluValLeuSerAspPheLysThrTrpLeuLysAlaLysLeuMet
TGGGACTGGATATGCGAGGTGTTGAGCGACTTTAAGACCTGGCTAAAAGCTAAGCTCATG
ACCCTGACCTATACGCTCCACAACTCGCTGAAATTCTGGACCGATTTTCGATTTCGAGTAC
- 6001 ProGlnLeuProGlyIleProPheValSerCysGlnArgGlyTyrLysGlyValTrpArg
CCACAGCTGCCTGGGATCCCCTTTGTGTCTGCCAGCGCGGTATAAGGGGGTCTGGCGA
GGTGTGACGGACCCCTAGGGGAACACAGGACGGTCCGCCCATATTCCCCCAGACCGCT
- 6061 Gly
ValAspGlyIleMetHisThrArgCysHisCysGlyAlaGluIleThrGlyHisValLys
GTGGACGGCATCATGCACACTCGCTGCCACTGTGGAGCTGAGATCACTGGACATGTCAA
CACCTGCCGTAGTACGTGTGAGCGACGGTGACACCTCGACTCTAGTGACCTGTACAGTTT
- 6121 AsnGlyThrMetArgIleValGlyProArgThrCysArgAsnMetTrpSerGlyThrPhe
AACGGGACGATGAGGATCGTCGGTCTTAGGACCTGCAGGAACATGTGGAGTGGGACCTTC
TTGCCCTGCTACTCTAGCAGCCAGGATCCTGGACGCTCTTGTACACCTCACCTTGGAAG
- 6181 ProIleAsnAlaTyrThrThrGlyProCysThrProLeuProAlaProAsnTyrThrPhe
CCCATTAAATGCCTACACACGGGCCCTGTACCCCCCTTCTGCGCCGAACACAGCTTC
GGGTAATTACGGATGTGGTGCCCGGGGACATGGGGGAAGGACGCGGCTTGATGTGCAAG
- 6241 AlaLeuTrpArgValSerAlaGluGluTyrValGluIleArgGlnValGlyAspPheHis
GCGCTATGGAGGTGTCTGCAGAGGAATATGTGGAGATAAGGCAGGTGGGGGACTTCCAC
CGCGATACCTCCCACAGACGTCTCCTTATACACCTCTATTCCGTCCACCCCTGAAGGTG
- 6301 TyrValThrGlyMetThrThrAspAsnLeuLysCysProCysGlnValProSerProGlu
TACGTGACGGGTATGACTACTGACAATCTCAATGCCCCGTGCCAGGTCCCATCGCCCGAA
ATGCACTGCCCATACTGATGACTGTTAGAGTTTACGGGCACGGTCCAGGTTAGCGGGCTT
- 6361 PhePheThrGluLeuAspGlyValArgLeuHisArgPheAlaProProCysLysProLeu
TTTTTACAGAATTGGACGGGGTGCGCTACATAGGTTTGGCGCCCCCTGCAAGCCCTTG
AAAAGTGTCTTAACCTGCCCCACGCGGATGTATCCAAACGCGGGGGACGTTCCGGGAAC
- 6421 LeuArgGluGluValSerPheArgValGlyLeuHisGluTyrProValGlySerGlnLeu
CTGCGGGAGGAGGTATCATTCAGAGTAGGACTCCACGAATACCCGTTAGGGTTCGCAATTA
GACGCCCTCCTCCATAGTAAGTCTCATCTGAGGTGCTTATGGGCCATCCAGCGTTAAT

SUBSTITUTE SHEET

FIG. 17-8

6481 ProCysGluProGluProAspValAlaValLeuThrSerMetLeuThrAspProSerHis
CCTTGGCAGCCCGAACCGGACGTGGCCGTGTGACGTCCATGCTCACTGATCCCCCCCAT
GGAACGCTCGGGCTTGGCCTGCACCGGCACAACGTCAGGTACGAGTGACTAGGGAGGGTA

6541 IleThrAlaGluAlaAlaGlyArgArgLeuAlaArgGlySerProProSerValAlaSer
ATAACAGCAGAGGCGCGGGCGAAGGTGGCGAGGGGATACCCCCCTCTGTGCCAGC
TATTGTCTGCTCCGCCGCGCCGCTTCCAACCGCTCCCCCTAGTGGGGGAGACACCGGTG

6601 SerSerAlaSerGlnLeuSerAlaProSerLeuLysAlaThrCysThrAlaAsnHisAsp
TCCTCGGCTAGCCAGCTATCCGCTCCATCTCTCAAGGCAACTTGCACCGCTAACCATGAC
AGGAGCCGATCGGTGATAGGCGAGGTAGAGAGTTCCGTTGAACGTGGCGATTGGTACTG

6661 SerProAspAlaGluLeuIleGluAlaAsnLeuLeuTrpArgGlnGluMetGlyGlyAsn
TCCCCGTGATGCTGAGCTCATAGAGGCCAACCTCCTATGGAGGCAGGAGATGGGCGGCAAC
AGGGGACTACGACTCGAGTATCTCCGGTTGGAGGATACCTCCGTCTCTACCCGCCGTTG

6721 IleThrArgValGluSerGluAsnLysValValIleLeuAspSerPheAspProLeuVal
ATCACCAGGGTTGAGTCAGAAAACAAAGTGGTGATTCTGGACTCCTTCGATCCGCTTGTG
TAGTGGTCCCAACTCAGTCTTTTGTTCACCACTAAGACCTGAGGAAGCTAGGCGAACAC

6781 AlaGluGluAspGluArgGluIleSerValProAlaGluIleLeuArgLysSerArgArg
GCGAGGAGGACGAGCGGGAGATCTCCGTACCCGCGAGAAATCCTGCGGAAGTCTCGGAGA
CGCTCCTCTGCTCGCCCTCTAGAGGCATGGGCGTCTTTAGGACGCTTCAGAGCCTCT

6841 PheAlaGlnAlaLeuProValTrpAlaArgProAspTyrAsnProProLeuValGluThr
TTCGCCCAGGCCCTGCCGTTTGGGCGCGGCCGACTATAACCCCCGCTAGTGGAGACG
AAGCGGGTCCGGGACGGGCAACCCGCGCGGCTGATATTGGGGGGCGATCACCTCTGC

6901 TrpLysLysProAspTyrGluProProValValHisGlyCysProLeuProProProLys
TGGAAAAAGCCCGACTACGAACCACTGTGGTCCATGGCTGTCCGCTTCCACCTCCAAAG
ACCTTTTTCGGGCTGATGCTTGGTGGACACAGGTACCGACAGGCGAAGGTGGAGGTTTC

6961 SerProProValProProProArgLysLysArgThrValValLeuThrGluSerThrLeu
TCCCCTCTGTGCTCCGCCCTCGGAAGAAGCGGACGGTGGTCCCTCACTGAATCAACCCTA
AGGGGAGGACACGGAGGCGGAGCCTTCTTCGCTGCCACCAGGAGTGACTTAGTTGGGAT

7021 Ser
SerThrAlaLeuAlaGluLeuAlaThrArgSerPheGlySerSerSerThrSerGlyIle
TCTACTGCCTTGGCCGAGCTCGCCACCAGAAGCTTTGGCAGCTCCTCAACTTCCGGCATT
AGATGACGGAACCGGCTCGAGCGGTGGTCTTCGAAACCGTTCGAGGAGTTGAAGGCCGTAA

7081 ThrGlyAspAsnThrThrThrSerSerGluProAlaProSerGlyCysProProAspSer
ACGGGCGACAATACGACAACATCCTCTGAGCCCGCCCCCTTCTGGCTGCCCCCCGACTCC
TGCCCGCTGTTATGCTGTTGTAGGAGACTCGGGCGGGGAAGACCGACGGGGGGGCTGAGG

7141 PheAla
AspAlaGluSerTyrSerSerMetProProLeuGluGlyGluProGlyAspProAspLeu
GACGCTGAGTCCTATTCTCCATGCCCCCCCTGGAGGGGGAGCCTGGGGATCCGGATCTT
CTGCGACTCAGGATAAGGAGGTACGGGGGGGACCTCCCCCTCGGACCCCTAGGCCTAGAA

7201 SerAspGlySerTrpSerThrValSerSerGluAlaAsnAlaGluAspValValCysCys
AGCGACGGGTGATGGTCAACGGTCAGTAGTGAGGCCAACGCGGAGGATGTCGTGTGCTGC
TCGCTGCCAGTACAGTTGCCAGTCATCACTCCGGTTGCGCTCCTACAGCACACGACG

7261 SerMetSerTyrSerTrpThrGlyAlaLeuValThrProCysAlaAlaGluGluGlnLys
TCAATGTCTTACTCTTGGACAGGCGCACTCGTCACCCCGTGCGCCGCGAAGAACAGAAA
AGTTACAGAATGAGAACCTGTCCGCTGAGCAGTGGGGCACGCGGCGCTTCTTGTCTTT

7321 LeuProIleAsnAlaLeuSerAsnSerLeuLeuArgHisHisAsnLeuValTyrSerThr
CTGCCCATCAATGCACTAAGCAACTCGTTGCTACGTCACCACAATTTGGTGTATTCCACC
GACGGGTAGTTACGTGATTCTGTGAGCAACGATGCAGTGGTGTAAACCACATAAGGTGG

7381 ThrSerArgSerAlaCysGlnArgGlnLysLysValThrPheAspArgLeuGlnValLeu
ACCTCAGCAGTGCTTGCCAAAGGCAGAAAGTCACATTTGACAGACTGCAAGTTCTG
TGGAGTGCCTACGAACGGTTCCGTCTTCTTTTCACTGTAAACTGTCTGACGTTCAAGAC

SUBSTITUTE SHEET

33 / 40

FIG. 17-9

- 7441 AspSerHisTyrGlnAspValLeuLysGluValLysAlaAlaAlaSerLysValLysAla
GACAGCCATTACCAGGACGTACTCAAGGAGGTTAAAGCAGCGCGTCAAAAGTGAAGGCT
CTGTCGGTAATGGTCCTGCATGAGTTCCTCCAATTCGTGCGCCAGTTCCTCACTTCCGA
- Phe
- 7501 AsnLeuLeuSerValGluGluAlaCysSerLeuThrProProHisSerAlaLysSerLys
AACTTGCTATCCGTAGAGGAGCTTGACGCTGACGCCCCCACTCAGCCAAATCCAAG
TTGAACGATAGGCATCTCCTCGAACGTCGGACTGCGGGGTGTGAGTCGGTTTAGGTTCT
- 7561 PheGlyTyrGlyAlaLysAspValArgCysHisAlaArgLysAlaValThrHisIleAsn
TTTGTTATGGGGCAAAGACGTCCGTTGCCATGCCAGAAAGGCCGTAAACCCATCAAC
AAACCAATACCCCGTTTCTGCAGGCAACGGTACGGTCTTCCGGCATTTGGGTGTAGTTG
- 7621 SerValTrpLysAspLeuLeuGluAspAsnValThrProIleAspThrThrIleMetAla
TCCGTGTGGAAGACCTTCTGGAAGACAATGTAACACCAATAGACACTACCATCATGGCT
AGGCACACCTTTCTGGAAGACCTTCTGTTACATTGTGGTTATCTGTGATGGTAGTACCGA
- 7681 LysAsnGluValPheCysValGlnProGluLysGlyGlyArgLysProAlaArgLeuIle
AAGAACGAGGTTTCTGCGTTTCAGCCTGAGAAGGGGGTTCGTAAGCCAGCTCGTCTCATC
TTCTTGCTCCAAAGACGCAAGTCGGACTCTTCCCCCAGCATTGGTTCGACAGAGTAG
- 7741 ValPheProAspLeuGlyValArgValCysGluLysMetAlaLeuTyrAspValValThr
GTGTTCCCCGATCTGGGCGTGCCTGTGCGAAAAGATGGCTTTGTACGACGTGGTTACA
CACAAGGGGTAGACCCGCACGCGCACACGCTTTTCTACCGAAACATGCTGCACCAATGT
- 7801 LysLeuProLeuAlaValMetGlySerSerTyrGlyPheGlnTyrSerProGlyGlnArg
AAGCTCCCTTGGCCGTGATGGGAAGCTCCTACGGATTCCAATACTCACCAGGACAGCGG
TTCGAGGGGAACCGGCACTACCTTCGAGGATGCCTAAGGTTATGAGTGGTCTGTGCGC
- 7861 ValGluPheLeuValGlnAlaTrpLysSerLysLysThrProMetGlyPheSerTyrAsp
GTTGAATTCCTCGTGCAAGCGTGAAGTCCAAGAAAACCCCAATGGGGTTCTCGTATGAT
CAACTTAAGGAGCACGTTTCGCACCTTCAGGTTCTTTTGGGGTTACCCCAAGAGCATACTA
- 7921 ThrArgCysPheAspSerThrValThrGluSerAspIleArgThrGluGluAlaIleTyr
ACCGCTGCTTTGACTCCACAGTCACTGAGAGCGACATCCGTACGGAGGAGGCAATCTAC
TGGGCGACGAAACTGAGGTGTCAGTGACTCTCGCTGTAGGCATGCCTCCTCCGTTAGATG
- 7981 GlnCysCysAspLeuAspProGlnAlaArgValAlaIleLysSerLeuThrGluArgLeu
CAATGTTGTGACCTCGACCCCAAGCCCGCTGGCCATCAAGTCCCTCACGAGAGGCTT
GTTACAACACTGGAGCTGGGGGTTCGGGCGCACCGGTAGTTCAGGGAGTGGCTCTCCGAA
- Gly
- 8041 TyrValGlyGlyProLeuThrAsnSerArgGlyGluAsnCysGlyTyrArgArgCysArg
TATGTTGGGGGCCCTTACCAATTCAAGGGGGGAGAACTGCGGCTATCGCAGGTGCCGC
ATACAACCCCGGAGAAATGGTTAAGTTCCCCCTCTTGACGCCGATAGCGTCCACGGCG
- 8101 AlaSerGlyValLeuThrThrSerCysGlyAsnThrLeuThrCysTyrIleLysAlaArg
GCGAGCGGCGTACTGACAAGTCTGTGGTAACACCTCACTTGCTACATCAAGGCCCGG
CGCTCGCCGATGACTGTTGATCGACACCATTTGTGGGAGTGAACGATGTAGTTCCGGGCC
- 8161 AlaAlaCysArgAlaAlaGlyLeuGlnAspCysThrMetLeuValCysGlyAspAspLeu
GCAGCCTGTGAGCGCGCAGGGCTCCAGGACTGCACCATGCTCGTGTGTGGCGACGACTTA
CGTCGGACAGCTCGGGGTCCCGAGGTCTGACGTGGTACGAGCACACACCGCTGCTGAAT
- 8221 ValValIleCysGluSerAlaGlyValGlnGluAspAlaAlaSerLeuArgAlaPheThr
GTCGTTATCTGTGAAAGCGCGGGGTCCAGGAGGACGCGGCGAGCCTGAGAGCCTTCACG
CAGCAATAGACACTTTTCGCGCCCCCAGGTCTCTGCGCGGCTCGGACTCTCGGAAGTGC
- 8281 GluAlaMetThrArgTyrSerAlaProProGlyAspProProGlnProGluTyrAspLeu
GAGGCTATGACCAGGTACTCCGCCCCCTGGGGACCCCCACAACAGAATACGACTTG
CTCCGATACTGGTCCATGAGGCGGGGGGACCCCTGGGGGTGTTGGTCTTATGCTGAAC
- 8341 GluLeuIleThrSerCysSerSerAsnValSerValAlaHisAspGlyAlaGlyLysArg
GAGCTCATAACATCATGCTCCTCCAACGTGTCAGTCGCCACGACGGCGCTGGAAAGAGG
CTCGAGTATTGTAGTACGAGGAGGTTGCACAGTCAGCGGGTGTGCGCGACCTTTCTCC

SUBSTITUTE SHEET

34 / 40

8401 ValTyrTyrLeuThrArgAspProThrThrProLeuAlaArgAlaAlaTrpGluThrAla
GTCTACTACCTCACCCGTGACCCTACAACCCCTCGCGAGAGCTGCGTGGGAGACAGCA
CAGATGATGGAGTGGGCACCTGGGATGTTGGGGGAGCGCTCTCGACGCACCCTCTGTCTGT

8461 ArgHisThrProValAsnSerTrpLeuGlyAsnIleIleMetPheAlaProThrLeuTrp
AGACACACTCCAGTCAATTCTGGCTAGGCAACATAATCATGTTTGCCCCACACTGTGG
TCTGTGTGAGGTCAGTTAAGGACCGATCCGTTGTATTAGTACAAACGGGGGTGTGACACC

8521 AlaArgMetIleLeuMetThrHisPhePheSerValLeuIleAlaArgAspGlnLeuGlu
GCGAGGATGATACTGATGACCCATTTCTTTAGCGTCCTTATAGCCAGGGACCAGCTTGAA
CGCTCCTACTATGACTACTGGGTAAAGAAATCGCAGGAATATCGGTCCCTGGTCAACTT

8581 GlnAlaLeuAspCysGluIleTyrGlyAlaCysTyrSerIleGluProLeuAspLeuPro
CAGGCCCTCGATTGCGAGATCTACGGGGCTGCTACTCCATAGAACCCTTGATCTACCT
GTCCGGGAGCTAACGCTCTAGATGCCCCGACGATGAGGTATCTTGGTGAACCTAGATGGA

8641 ProIleIleGlnArgLeuHisGlyLeuSerAlaPheSerLeuHisSerTyrSerProGly
CCAATCATTCAAAGACTCCATGGCCTCAGCGCATTTTCACTCCACAGTTACTCTCCAGGT
GGTTAGTAAGTTTCTGAGGTACCGAGTCGCGTAAAGTGAGGTGTCAATGAGAGGTCCA

8701 GluIleAsnArgValAlaAlaCysLeuArgLysLeuGlyValProProLeuArgAlaTrp
GAAATTAATAGGGTGGCCGCATGCCTCAGAAAACCTGGGGTACCGCCCTTGCGAGCTTGG
CTTTAATTATCCCACCGCGTACGGAGTCTTTTGAACCCCATGGCGGGAACGCTCGAACC

8761 Gly
ArgHisArgAlaArgSerValArgAlaArgLeuLeuAlaArgGlyGlyArgAlaAlaIle
AGACACCGGGCCCGGAGCGTCCGCGCTAGGCTTCTGGCCAGAGGAGGCAGGGCTGCCATA
TCTGTGGCCCGGGCCTCGCAGGCGGATCCGAAGACCGGTCTCCTCCGTCCCCACGGTAT

8821 CysGlyLysTyrLeuPheAsnTrpAlaValArgThrLysLeuLys
TGTGGCAAGTACCTCTTCAACTGGGCAGTAAGAACAAAGCTCAAAC
ACACCGTTCATGGAGAAGTTGACCCGTCATTCTTGTTCGAGTTTG

FIG. 17-10


SUBSTITUTE SHEET

35 / 40


IMMUNOLOGICAL SCREENING IN BACTERIA

FIG. 18

Transform E coli with Recombinant Plasmids


(Blot Bacteria on
Nitrocellulose Filter)

IPTG Plate




Lyse with Chloroform


BSA absorbtion/DNAse/Lysozyme

Incubate with primary
antibody

Wash

Incubate with
 125 I secondary antibody

Wash



Autoradiograph

SUBSTITUTE SHEET

36 / 40

FIG. 19-1




FIG. 19-2




FIG. 19-3




FIG. 19-4







FIG. 19-5



SUBSTITUTE SHEET



FIG. 20-1



ANTIG INDEX

HCV (1 - 500)

FIG. 20-2




ANTIG INDEX


HCV (501 - 1000)

SUBSTITUTE SHEET


HYDROPHILIC FIG. 20-3



HYDROPHILIC FIG. 20-4




HYDROPHILIC FIG. 20-5



ANTIGEN INDEX

HCV (2001 - 2500)

HYDROPHILIC FIG. 20-6



ANTIGEN INDEX

HCV (2501 - 2955)


Some conserved co-linear peptides in HCV & Flaviviruses

FIG. 21

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US90/01348

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC(5): A61K 39/12; C07H 21/04; C07K 7/06, 08, 10, 13/00, 15/28;		
US: 536/27; 435/6, 7, 69.1, 320, 240.1; 530/324-327, 350, 387, 416; 424/89		
II. FIELDS SEARCHED		
Minimum Documentation Searched ?		
Classification System	Classification Symbols	
U.S.	536/27; 435/6, 7, 69.1, 320, 240.1 530/324-327, 350, 387, 416; 424/89	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
Automated Patent Search, Chemical Abstract Service		
III. DOCUMENTS CONSIDERED TO BE RELEVANT *		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A, P	US, A, 4,870,026 WANDS ET AL. 26 September 1989	1-34
A	US, A, 4,702,909 VILLAREJOS ET AL. 27 October 1987	1-34
A	US, A, 4,542,016 TREPO 17 September 1985	1-34
X, P	Science, "Isolation of a c DNA clone derived from a blood-borne non-A, non-B viral hepatitis genome" volume 244, pp 359-362. CHOO ET AL. 21 April 1989 see abstract & figs.	1-12, 15, 16, 18-20, 22, 28-31, 33
X, P	Science, "An assay for circulating antibodies to a major ehologic virus of human nonA, nonB Hepatitis" volume 244, pp 362-364. KUO ET AL 21 April 1989 see abstract.	18, 19, 22
<p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
28 MAY 1990	13 JUL 1990	
International Searching Authority	Signature of Authorized Officer	
ISA/US	NINA COSSANNA	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Continuation of Classification:

IPC(5) C12N 1/11, 15/02, 51 C12P 21/02; C12Q 1/68;
G01N 33/53

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the International application that do not comply with the prescribed requirements to such an extent that no meaningful International search can be carried out¹³, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING²

This International Searching Authority found multiple inventions in this International application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International search report covers all searchable claims of the International application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this International search report covers only those claims of the International application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.